

# Knowledge about emergency procedure in case of dental trauma among paediatricians in Croatia

---

**Nikolić, Harry; Ivančić Jokić, Nataša; Bakarčić, Danko; Hrvatin, Sandra; Jakljević, Nika**

*Source / Izvornik:* **European Journal of Paediatric Dentistry, 2018, 19, 277 - 281**

**Journal article, Published version**

**Rad u časopisu, Objavljena verzija rada (izdavačev PDF)**

<https://doi.org/10.23804/ejpd.2018.19.04.5>

*Permanent link / Trajna poveznica:* <https://urn.nsk.hr/urn:nbn:hr:184:477121>

*Rights / Prava:* [Attribution-NonCommercial-NoDerivatives 4.0 International/Imenovanje-Nekomercijalno-Bez prerada 4.0 međunarodna](#)

*Download date / Datum preuzimanja:* **2025-03-28**



*Repository / Repozitorij:*

[Repository of the University of Rijeka, Faculty of Medicine - FMRI Repository](#)



H. Nikolic\*, N. Ivancic Jokic\*\*, D. Bakarcic\*\*,  
S. Hrvatin\*\*, N. Jakljevic\*\*\*

Faculty of Medicine, University of Rijeka, Rijeka, Croatia

\*Department of Surgery

\*\*Department of Paediatric Dentistry

\*\*\*Private practice, Rijeka, Croatia

email: danko.bakarcic@medri.uniri.hr

DOI: 10.23804/ejpd.2018.19.04.5

# Knowledge about emergency procedure in case of dental trauma among paediatricians in Croatia

## ABSTRACT

**Aim** The aim of this study was to investigate paediatricians' actual attitudes towards emergency management of dental injuries, need for potential improvement in their knowledge and motivation for further education.

**Materials and Methods** This study was a questionnaire-based cross-sectional survey on a sample of 84 respondents from four Croatian counties. A questionnaire of 24 questions was divided into four parts: personal and professional profile; self-assessed perceived knowledge and experience with dental injuries; specific questions on the management of dental injuries and specific questions regarding education about dental injuries. The questionnaires were collected during a 2-month period.

**Results** A larger percentage (95%) of surveyed paediatricians have at least once encountered trauma in the practice and in this case 90% of them contacted the dentist. Three quarters of participants (76%) would correctly hold completely extruded tooth. In case of tooth or fragment preservation 60% of the subjects would place the tooth in the saline solution, and only 12% of the respondents would use milk as a transport medium. The 19% of the paediatricians

considered that the permanent tooth should not be replanted. Two thirds (62%) of participants did not get any information regarding dental trauma during their past education, but most of them (90%) consider that education about traumatic dental injuries is important, and 86% of the respondents are willing to be educated on the subject in the future. There is no difference in procedures (recommended vs. non-recommended) undertaken during management of dental injuries, according to the participants working place, experience in emergency department or their personal experience of dental trauma.

**Conclusion** The results revealed insufficient paediatricians' knowledge about emergency procedures in case of dental trauma, but they also point out high awareness of the importance of dental trauma, and willingness for further education.

**Keywords** Child, Dental trauma, Knowledge, Paediatricians.

## Introduction

All injuries involving the orofacial area are considered a serious public health problem, they should be treated as an emergency, and their timely and appropriate emergency management can enhance favourable prognosis and outcome for the injured tissues [Schwartz-Arad et al., 2004; Schwartz-Arad and Levin, 2004; Unal et al., 2014; Oleszkiewicz and Emerich, 2015].

In order to prevent aesthetic, functional, psychological and economic consequences for patients and their families, it is necessary for the injured child to visit the dental office immediately after injury [Andreasen and Andreasen, 2000; Rocha and Cardoso, 2001; Altay and Gungör, 2001; Nik-Hussein, 2001; Levin et al., 2003; Ulusoy et al., 2012].

When dental trauma occurs, parents, guardians, teachers, coaches or persons in a child's vicinity are responsible for helping the injured child, and children are referred to paediatric dentists or paediatric surgeons but also to doctors of various specialties, most often paediatricians [Ulusoy et al., 2012]. In order for paediatricians to act properly in these cases, it is necessary for them to have specific knowledge about the urgent procedure and the importance of the immediate beginning of treatment. Adequate paediatricians' knowledge about the treatment of dental injuries can play a significant role in the outcome of dental trauma in children [Škrinjari et al., 2010; Emerich and Gazda, 2010; Emerich and Wyszowski, 2010; Zampogna et al., 2014; Chanchala et al., 2016]. Prognosis of traumatised teeth highly depends on

immediate and appropriate emergency management of the injured tissues [Ulusoy et al., 2012; Škrinjarić et al., 2010; Zampogna et al., 2014]. Therefore, it is of particular importance that paediatricians are prepared to provide prompt and proper sequence of actions, thus reducing or preventing the possible serious consequences of dental trauma for the patient and his family [Ulusoy et al., 2012; Chanchala et al., 2016; Lin et al., 2006; Da Silva et al., 2004].

As recent literature is not abundant with studies regarding this subject especially from Croatia, the aim of this study was to investigate Croatian paediatricians' actual attitudes towards emergency management of dental injuries and need for potential improvement in their knowledge related to the topic of dental trauma emergency procedures. The further aim was to investigate if their place of work, experience in emergency department (ED) and personal experience of dental trauma influenced their action in case of patient with dental injuries. The aim was also to investigate if they are motivated for further education on the matter if improvement is necessary. Then the results of this report could provide information that can assist in creating instructive guidelines for paediatricians in their continuing education.

## Materials and method

This study was conducted as a questionnaire-based cross-sectional survey on a sample of 84 respondents from four Croatian counties: Primorsko-goranska, Istarska, Ličko-senjska and Splitsko-dalmatinska county.

For research purposes, a questionnaire of 24 questions was used. The questionnaire was designed and developed by the authors, and included: general information on respondents, information on education and employment, knowledge of emergency procedures, questions about possible prior education and wishes for further education on the management of dental trauma. To ensure the participants' self-approach in answering the questions and exclude any side intervention or consultation in the course of filling out the questionnaire forms, the investigators were individually present during distribution of the surveys and until the respondents completed the questionnaires. The examination was anonymous and carried out solely with voluntary consent of the respondent. In addition to the survey, an informed consent that described the purpose of the research and guaranteed confidentiality of data and their use for scientific purposes only, was provided. The study was approved by the Ethical Committee of the Faculty of Medicine, University of Rijeka, Croatia.

A total of 86 questionnaires were collected during a 2-month period, starting from January through March, 2014. Of these, 84 questionnaires were properly filled out,

whereas 2 were excluded due to incomplete answers.

Statistical analysis was performed using MedCalc statistical software (MedCalc Software ver. 18, Mariakerke, Belgium). Number and percentage of answer frequency were calculated. Chi-square analysis was performed to test differences in the management

Question	Answer	n (%)
Familiarity with the term tooth avulsion	Yes	74 (88%)
	No	10 (12%)
Personal experience with dental injuries	Yes	24 (29%)
	No	60 (71%)
Have you ever witness the dental injury observed a dental injury in your practice?	Yes, once	22 (26%)
	Yes, several times	58 (69%)
	No	4 (5%)

TABLE 1 Knowledge and experience with dental injuries (N=84).

Question	Answer	n (%)
How would you hold a completely extruded tooth	By the crown*	64 (76%)
	Would hold the whole tooth (it is not important how)	4 (5%)
	Would not touch it	16 (19%)
What would you do in case of a completely extruded tooth or with the parts of a broken tooth?	Place the tooth:	
	in saline solution*	50 (60%)
	in milk*	10 (12%)
	on clean handkerchief	12 (14%)
	in water	4 (5%)
	in disinfectant	4 (5%)
	in child's saliva*	2 (2%)
Do nothing	2 (2%)	
Would you replant the permanent tooth?	Yes*	68 (81%)
	No	16 (19%)
Would you replant the primary tooth?	Yes	34 (40%)
	No*	50 (60%)
What would you do in case of dental fracture?	Contact the dentist*	60 (72%)
	Advise parents to contact the dentist	22 (26%)
	Do nothing	2 (2%)
What would you do in case of tooth mobility?	Contact the dentist*	50 (59%)
	Advise parents to contact the dentist	30 (36%)
	Do nothing	4 (5%)

\* recommended procedure

TABLE 1 Management of dental injuries (N=84).

Dental trauma management		Working place			ED		Stat.	Personal experience		Stat.
	Procedure	Primary care (n)	Hospital (n)		No (n)	Yes (n)		No (n)	Yes (n)	
Tooth handling	Recommended (n)	48	16	$X^2=2,01$	34	30	$X^2=3,23$	46	18	$X^2=0,03$
	Non recommended (n)	18	2	$P=0.156$	6	14	$P=0.072$	14	6	$P=0.872$
Tooth transport	Recommended (n)	49	13	$X^2=0.03$	29	33	$X^2=0,07$	44	18	$X^2=0,02$
	Non recommended (n)	17	5	$P=0.864$	11	11	$P=0.796$	16	6	$P=0.876$
Primary teeth replantation	Recommended (n)	38	12	$X^2=0.48$	24	26	$X^2=0,01$	38	12	$X^2=1,25$
	Non recommended (n)	28	6	$P=0.489$	16	18	$P=0.933$	22	12	$P=0.264$
Permanent teeth replantation	Recommended (n)	55	13	$X^2=1,12$	32	36	$X^2=0,04$	48	20	$X^2=0,12$
	Non recommended (n)	11	5	$P=0.290$	8	8	$P=0.833$	12	4	$P=0.727$
Tooth fracture procedure	Recommended (n)	44	16	$X^2=3,38$	32	28	$X^2=2,72$	40	20	$X^2=2,31$
	Non recommended (n)	22	2	$P=0.066$	8	16	$P=0.099$	20	4	$P=0.129$
Tooth mobility procedure	Recommended (n)	42	8	$X^2=2,14$	28	22	$X^2=3,44$	30	20	$X^2=7,81$
	Non recommended (n)	24	10	$P=0.144$	12	22	$P=0.064$	30	4	$P=0.005$

TABLE 3 Differences in the management of dental trauma according to participants' working characteristics.

of dental trauma according to participants' characteristics e.g. working place, experience in emergency department (ED) and personal experience in dental trauma.

## Results

The questionnaire was completed by 84 of 86 respondents. Results were expressed as a number and percentage of question's response. The results regarding: knowledge and experience with dental injuries are shown in Table 1; management of dental trauma in Table 2, differences in the management of dental trauma according to participants' characteristics in Table 3 and education about dental injuries in Table 4.

The median age of the respondents was 51,5, ranging from 25 to 64 years. The study included 72 female (86%) and 12 male (14%) paediatricians. The median of working experience with children was 20 years, range 0,5-40 years. During their professional career 44 (52%) of respondents had the opportunity to work in ED, while 40 (48%) did not. Answers are summarised in Tables 1, 2 and 4 according to three groups of questions (knowledge, management and education regarding dental trauma), and in Table 3 differences in the management of dental trauma according to participants characteristics are presented.

## Discussion

### Limitations

This cross-sectional study has some limitations; it

Question	Answer	n (%)
Have you ever been educated about dental injuries?	Yes	32 (38%)
	No	52 (62%)
Are you well informed about dental injuries?	Completely	6 (7%)
	Partially	56 (67%)
	Not informed at all	22 (26%)
Is education about dental injuries important?	Yes	76 (90%)
	Neither important nor unimportant	8 (10%)
Are you interested in education about dental trauma?	Yes	72 (86%)
	No	12 (14%)

TABLE 4 Education about dental injuries (N=84).

was conducted on a small sample of paediatricians from only 4 out of 21 counties of Croatia, which limits the possibility to generalise the findings to the whole Croatian paediatricians' population. But we consider this sample representative enough to emphasise the fact clearly revealed from the survey; paediatricians lack education in dental injuries and serious actions need to be undertaken to change attitudes and procedures usually done by majority of paediatricians.

Timely and adequate emergency procedures for traumatic injuries of orofacial structures are of crucial importance for optimal therapy success, while untreated traumas can have severe consequences [Andreasen and Andreasen, 2008; Zampogna et al., 2014; Chanchala et al., 2016; Lin et al., 2006; Da Silva et al., 2004].

Recent literature is not abundant with studies

regarding paediatricians' actual attitudes and knowledge towards emergency management of dental injuries. That was one of the strongest motives to conduct this study, to investigate attitudes among paediatricians. We assumed that paediatricians working in hospital rather than those in primary care clinics, those with experience in emergency department and those with personal experience with dental trauma would more often know the recommended procedures in case of dental injury. However, there were no difference between all aforementioned categories, with the exception that paediatricians with personal experience with dental trauma will more likely know recommended procedure in a case of dental mobility than those without personal experience.

Current research point to the importance of education and training of health professionals, including paediatricians, in order to learn how to react promptly in case of dental trauma and prevent losing the precious time that plays a decisive role in dental trauma therapy [Ulusoy et al., 2012; Zampogna et al., 2014; Chanchala et al., 2016; Lin et al., 2006].

The median of working experience with children was 20 years, and during their working life 52% of respondents had the opportunity to work in ED, while 48% did not.

Familiarity with the term tooth avulsion was showed by 88% of the respondents, and this percentage is higher than that found in the studies of Bahammam [2018] and Dali et al. [2014] of 69% and 57%, respectively. A larger percentage of as many as 95% of surveyed paediatricians have at least once encountered trauma in their practice, larger than that the 65% reported by Chanchala et al. [2016].

Seventy-six percent of respondents would correctly hold a completely extruded tooth for the crown, and this percentage is higher than those found in the studies of Bahammam [2018] and Dali et al. [2014], and similar to that of Yothi et al. [2015]. Disappointing is the fact that 19% would not dare to touch the tooth.

In case of tooth or fragment preservation, half of the subjects would place the tooth in saline solution (53.6%), less than in the Chanchala et al. study (90%) [2016], but more than in the Bahammam study (21%) [2018]. Only 15.5% of the respondents would use milk as a transport medium, similarly to Chanchala et al. (10%) [2016], but lesser than in the Bahammam (31%) [2018] and Ulusoy et al. [2012] studies. Others (26%) would do something that generally is not recommended and that immediately reduces the chances of long-term survival of the tooth and periodontal tissues (i.e. wrap the tooth in a clean handkerchief, wash with water, place the tooth in the disinfectant). About 2% of respondents would not do anything. This is indeed a small percentage, nevertheless it is disappointing and unacceptable. Disappointing is also that 19% of the paediatricians considered that the permanent tooth

should not be replanted, a higher percentage than that found by Chanchala et al. (5%) [2016]. Nearly half (40%) of the respondents think that avulsed primary teeth should be replanted. This points out that a child who has experienced a dental trauma has higher chances to lose the permanent teeth or have some complications (e.g. permanent tooth dilaceration) due to replantation of the primary teeth because of inadequate treatment caused by paediatricians' poor knowledge about teeth replantation.

In other types of trauma involving fractures and injuries to periodontal tissues, paediatricians would most likely contact a dentist or advise the parents to contact their dentist. According to the data obtained in the study, 72% of the respondents would contact a dentist in case of dental fractures and will rarely advise parents to contact their dentist, and in the case of tooth mobility 59% of the respondents would contact a dentist. A small percentage of paediatricians would completely ignore this type of trauma (2%) and do nothing (5%). This alerts that some paediatricians as well as other non-dental medical professionals do not recognise the severity of such trauma and do not know how to provide appropriate first-aid in case of dental trauma [Holan and Shmueli, 2003; Abu-Dawoud et al., 2007; Subhashraj, 2009; Qazi and Nasir, 2009; Trivedy et al., 2012].

As far as the differences in the management of dental trauma according to participants characteristics (working place, experience in ED and personal experience in dental trauma) showed no statistically significant difference between recommended and non-recommended procedures in dental trauma management such as: tooth handling and transportation, tooth replantation, tooth fracture and mobility management regarding: working place, ED experience and personal experience with dental trauma. Re et al. [2014] in their study show a slightly better knowledge in professionals working in hospitals but this was the case of dentists employed in hospitals. Only in the case of personal experience and tooth mobility management the statistically significant difference is revealed, and in this case the respondents with personal experience of dental trauma are more likely to apply the correct recommended procedures. Thus, no matter whether they work in primary care or had worked in ED, or they had their own experience with dental trauma, there is no difference between recommended and non recommended procedures that paediatricians would perform in dental trauma management.

Paediatricians' education about dental trauma show some disappointing results. About two thirds of the respondents (62%) did not receive any information regarding dental trauma within their past education. The largest number of paediatricians considered to be partially informed (67%) while more than a quarter was not informed (26%), so the majority of paediatricians

were not adequately educated, similarly to the results of the Bahammam study [Bahammam, 2018].

A positive fact is that most of the respondents (90%) consider education about traumatic dental injuries important and that 86% of them are willing to be educated on the subject in the future, similarly to the findings of previous studies [Chanchala et al., 2016; Bahammam, 2018; Ulusoy et al., 2012; Aren et al., 2018]. The remaining 14% believe that such information is unnecessary.

## Conclusion

The results of this research showed insufficient paediatricians' knowledge about emergency procedures in case of dental trauma even though they meet such cases, and they do not recognise its severity. The results also show paediatricians' awareness of the consequences of dental trauma, and their willingness for further education.. Thus, special attention should be given to provide paediatricians additional education on this subject during pre- and post-graduate programmes.

## References

- › Abu-Dawoud M, Al-Enezi B, Andersson L. Knowledge of emergency management of avulsed teeth among young physicians and dentists. *Dent Traumatol* 2007;23:348–55.
- › Altay N, Güngör HC. A retrospective study of dento-alveolar injuries of children in Ankara, Turkey. *Dent Traumatol* 2001;17:201-4.
- › Andreasen JO, Andreasen FM. *Textbook and color atlas of traumatic Injuries to the Teeth*. Munksgaard: Copenhagen; 2008.
- › Aren A, Erdem AP, Aren G, Şahin ZD, Güney Tolgay C, Çayırıcı M, et al. Importance of knowledge of the management of traumatic dental injuries in emergency departments. *Ulus Travma Acil Cerrahi Derg* 2018;24:136–144
- › Bahammam LA. Knowledge and attitude of emergency physician about the emergency management of tooth avulsion. *BMC Oral Health*. 2018;18(1):57
- › Chanchala HP, Shanbhog R, Ravi MD, Raju V. Paediatricians' perspectives on dental trauma management: A cross-sectional survey. *J Indian Assoc Public Health Dent* 2016;14:419-23.
- › Dali M, Naulakha D, Rajbanshi L. Practice in emergency management of avulsed tooth among medical doctors in Nobel medical college, Biratnagar, Nepal: A cross sectional survey. *Int J Dent Heal Sci* 2014;1(1):3–12.
- › Da Silva AC, Passeri LA, Mazzone R, De Moraes M, Moreira RWF. Incidence of dental trauma associated with facial trauma in Brazil: a 1-year evaluation. *Dent Traumatol* 2004;20:6-11.
- › Emerich K, Gazda E. Review of recommendations for the management of dental trauma presented in first-aid textbooks and manuals. *Dent Traumatol* 2010;26: 212-6.
- › Emerich K, Wyszowski J. Clinical practice: dental trauma. *Eur J Pediatr* 2010;169: 1045-50.
- › Holan G, Shmueli Y. Knowledge of physicians in hospital emergency rooms in Israel on their role in cases of avulsion of permanent incisors. *Int J Paediatr Dent* 2003;13:13–9.
- › Levin L, Ashkenazi M, Schwartz-Arad D. Preservation of alveolar bone of un-restorable traumatized maxillary incisors for future implantation. *J Israel Dent Assoc* 2004;21:54-9.
- › Levin L, Friedlander LD, Geiger SB. Dental and oral trauma and mouthguard use during sport activities in Israel. *Dent Traumatol* 2003;19:237-42.
- › Lin S, Levin L, Emodi O, Fuss Z, Peled M. Physician and emergency medical technicians' knowledge and experience regarding dental trauma. *Dent Traumatol* 2006;22:124-6.
- › Nik-Hussein NN. Traumatic injuries to anterior teeth among schoolchildren in Malaysia. *Dent Traumatol* 2001;17:149-52.
- › Oleszkiewicz I, Emerich K. How to proceed in case of tooth avulsion: state of student knowledge. *Eur J Paediatr Dent* 2015;16(2):103-6.
- › Qazi SR, Nasir KS. First-aid knowledge about tooth avulsion among dentists, doctors and lay people. *Dent Traumatol* 2009;25:295–9.20.
- › Rocha MJ, Cardoso M. Traumatized permanent teeth in Brazilian children assisted at the Federal University of Santa Catarina, Brazil. *Dent Traumatol* 2001;17:245-9.
- › Re D, Augusti D, Paglia G, Augusti G, Cotti E. Treatment of traumatic dental injuries: evaluation of knowledge among Italian dentists. *Eur J Paediatr Dent* 2014;15(1):23-8.
- › Schwartz-Arad D, Levin L, Ashkenazi M. Treatment options of untreatable traumatized anterior maxillary teeth for future use of dental implantation. *Implant Dent* 2004;13:120–8.
- › Schwartz-Arad D, Levin L. Post-traumatic use of dental implants to rehabilitate anterior maxillary teeth. *Dent Traumatol* 2004;20:344–7.10.
- › Škrinjaric I, Škrinjaric T, Gorseta K, Čuković-Bagic I, Verzak Ž. Hitni i preventivni postupci kod trauma zuba u djece. *Paediatr Croat* 2010;54:154-162.
- › Trivedy C, Kodate N, Ross A, Al-Rawi H, Jaiganesh T, Harris T, et al. The attitudes and awareness of emergency department (ED) physicians towards the management of common dentofacial emergencies. *Dent Traumatol* 2012;28:121–6.
- › Ulusoy AT, Onder H, Cetin B, Kaya S. Knowledge of medical hospital emergency physicians about the first-aid management of traumatic tooth avulsion. *Int J Paediatr Dent* 2012;22:211-6.
- › Unal M, Oznurhan F, Kapdan A, Aksoy S, Dürer A. Traumatic dental injuries in children. Experience of a hospital in the central Anatolia region of Turkey. *Eur J Paediatr Dent* 2014;15(1):17-22.
- › Jyothi KN, Venugopal P, Nanda S, Shah MK. Knowledge and attitude of medical doctors towards emergency management of avulsed tooth-a cross sectional survey. *J Dent Sci Res* 2015;2:156–67.
- › Zampogna S, De Filippo S, Talarico V, Aloe M, Severini N, Pizzi S et al. First aid in dental trauma in pediatric age. *Italian Journal Pediatrics* 2014;40:A72.