

Lesions of Left Basal Ganglia and Insula Structures Impair Executive Functions but not Emotion Recognition: A Case Report

Švegar, Domagoj; Antulov, Ronald; Tkalčić, Mladenka; Antončić, Igor

Source / Izvornik: **Brain Impairment**, 2016, 17, 233 - 241

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.1017/BrImp.2016.23>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:184:684712>

Rights / Prava: [Attribution-NonCommercial-NoDerivatives 4.0 International](#)/[Imenovanje-Nekomercijalno-Bez prerada 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2025-02-07**



Repository / Repozitorij:

[Repository of the University of Rijeka, Faculty of Medicine - FMRI Repository](#)



Lesions of Left Basal Ganglia and Insula Structures Impair Executive Functions but not Emotion Recognition: A Case Report

Domagoj Švegar,¹ Ronald Antulov,² Mladenka Tkalčić¹ and Igor Antončić²

¹ Department of Psychology, Faculty of Humanities and Social Sciences, University of Rijeka, Rijeka, Croatia

² Department of Radiology, Clinical Hospital Center Rijeka, Rijeka, Croatia

Increasing research evidence suggests that basal ganglia are an important part of frontal-subcortical circuit which is involved not only in motor control but also in affective, cognitive and executive functions. In this article, we describe the ability of facial emotion recognition and cognitive functioning in a patient with left basal ganglia and insula damage. The patient's ability to recognise facial emotional expressions was intact in spite of unilateral injury of the left insula and basal ganglia. He showed preserved intellectual function in general, but experienced difficulties on subsets of the executive functions: set-shifting and ability to activate or generate cognitive strategies, commonly found in patients with caudate lesions. This case contributes to evidence that striatal structures are important for executive functions.

Keywords: basal ganglia, insula, emotion recognition, executive functions, infarction

Introduction

Numerous brain structures participate in recognising facial emotional expressions: the occipitotemporal cortices, amygdala, parietal cortices, orbitofrontal cortex, insula and basal ganglia (Adolphs, Tranel, Koenigs, & Damasio, 2005). In prior research, it has been established that the basal ganglia and specifically the insula have an important role in the recognition of facial disgust (Johnson et al., 2007; Phillips et al., 1997). Nevertheless, there is no consensus regarding the different role of the left and right side of these brain structures in the process of emotion recognition. Several authors reported that exposure to disgusted facial stimuli elicits activation in the right insula and right basal ganglia (Adolphs, Damasio, Tranel, Cooper, & Damasio, 2000; Anderson, Christoff, Panitz, De Rosa, & Gabrieli, 2003; Phillips et al., 1997, 2004;

Ruiz et al., 2013; Schroeder et al., 2004). According to their results, it can be concluded that patients with unilateral lesions of the right insula are expected to experience serious problems with recognition of disgusted facial stimuli, whilst patients with unilateral lesions of the left insula should not experience such deficits. Partly in contrast to their reports, Sprengelmeyer, Rausch, Eysel and Przuntek (1998) demonstrated that disgusted facial expressions activated the right putamen and left insular cortex. Furthermore, Calder, Keane, Manes, Antoun and Young (2000) presented a patient with infarction in the left hemisphere, involving insula and basal ganglia, who showed a deficit in recognition of facial expression of disgust, even though the right basal ganglia and the right insula of this patient remained intact. Finally, some other studies demonstrated that bilateral insula lesions resulted

Address for correspondence: Domagoj Švegar, Department of Psychology, Faculty of Humanities and Social Sciences, University of Rijeka, Filozofski fakultet u Rijeci, Odsjek za psihologiju, Sveučilišna avenija 4, 51000 Rijeka, Croatia (HR). Phone: +385 51 265 770. E-mail: dsvegar@ffri.hr

in an impaired ability to recognise facial disgust (e.g., Adolphs et al., 2005).

Additionally, basal ganglia are a part of frontal-subcortical circuit which is involved not only in motor control and emotion processing but also in cognitive and executive functions (Alexander, DeLong, & Strick, 1986; Lichter & Cummings, 2001; Middleton & Strick, 2002; Salmon, Heindel, & Hamilton, 2001). Evidence that those structures are particularly important for executive functions emerges from neuropsychological studies of neurological disorders such as Huntington's and Parkinson's diseases as well as from studies of basal ganglia damage caused by stroke or traumatic brain injury (Backman, Robins-Wahlin, Lundin, Ginovart, & Farde, 1997; Elliott, 2003; Salmon et al., 2001). These findings have led to the suggestion that executive function depends not on the prefrontal cortex (PFC) in isolation, but on the intact functioning of corticostriatal circuitry (Bonelli & Cummings, 2007; Elliott, 2003; Salmon et al., 2001). The executive functions consist of those capacities that enable an individual to engage in independent, purposive, self-directed and self-serving behaviour (Lezak, Howieson, Bigler, & Tranel, 2012). Thus, executive functions relate to various complex cognitive processes and sub-processes, such as task-switching, planning, working memory, solving novel problems, modifying behaviour, generating strategies, sequencing complex actions, etc. (Elliott, 2003).

The first objective of this study was to assess if the lesions of the left insula and the left part of the basal ganglia are related to impaired recognition of facial expression of emotion, especially disgust. Since the injury of our patient was almost identical to Calder et al.'s (2000) patient's injury, we hypothesised that he would experience the same deficits in recognition of disgusted facial expressions. The second objective was to test the patient's cognitive and executive functioning. We hypothesised that due to impaired corticostriatal connectivity our patient would show deficiency in certain cognitive and executive functions primarily related to frontal lobes.

Materials and Methods

Case Description


MS is a 21-year-old right-handed male attending the final year of his undergraduate study. Two years ago, MS presented to the Emergency Department with right extremities numbness and later developed choreoathetoid movements of the right arm. His mental status was normal. Brain MRI done at the time of his admission to the hospital showed an

acute left cerebral hemisphere infarction involving part of the basal ganglia (putamen, caudate nucleus head, globus pallidus and claustrum), as well as left insula (Figure 1). After a month, his motor function recovered completely. On a follow-up visit done 3 months after hospital discharge his neurological examination was unremarkable. MS returned to all of his everyday activities which he used to do prior to his illness.

Tests

Emotion identification test. In order to assess MS's ability to recognise emotional expressions in general and expression of disgust in particular, the Karolinska Directed Emotional Faces (KDEF) (Lundqvist, Flykt & Öhman, 1998) stimuli set was used. That dataset is comprised of 4900 photographs of human facial expressions, displaying anger, disgust, fear, happiness, sadness and surprise, as well as neutral expressions. The KDEF database is equivalent to Ekman and Friesen's (1976) Pictures of facial affect (PFA) database, which contains facial expressions of same six emotions. For the purpose of the present research, Calvo and Lundqvist's (2008) adaptations of the KDEF photographs were used. Stimuli were selected according to normative data regarding identification accuracy obtained on Spanish (Calvo & Lundqvist, 2008) and Croatian (Milovanović, Švegar, & Kardum, 2013) populations. Recognition of emotional expressions was examined via forced choice labelling procedure, using equivalent procedure as in Milovanović et al.'s (2013) study. MS was presented with a total of 70 photographs: 60 emotional (10 per each of six emotions) and 10 neutral. His task was to identify each of successively presented expressions by selecting one label from a list of seven possible expressions: anger, disgust, fear, happiness, sadness, surprise and neutral. In order to ensure higher reliability, the procedure was repeated after 14 days, with the same stimuli, displayed in a randomly selected serial order.

Neuropsychological tests. In order to assess cognitive and executive functions, we applied standard neuropsychological tests listed and described in detail below. All neuropsychological tests were translated to Croatian language and standardised for use on a Croatian population (Galić, 2002). MS's performance on all neuropsychological tests, except the Stroop test and the Tower of Hanoi test, were compared with a normative results obtained on Croatian (Galić, 2002) and international (Lezak et al., 2012; Strauss, Sherman & Spreen, 2006) samples of healthy adults stratified for age and

**FIGURE 1**

(a) Axial T2-weighted brain MRI shows left cerebral hemisphere post-infarction encephalomalacia involving part of the basal ganglia (putamen, caudate nucleus head, globus pallidus and claustrum), as well as left insula. (b) Coronal T2-weighted brain MRI also demonstrates post-infarction encephalomalacia of a part of the left basal ganglia and left insula.

years of education. In order to evaluate the performance of MS on the Stroop and the Tower of Hanoi tests, we formed two control samples, each comprised of 20 university students. The detailed description of these samples is provided later in the Method section.

Tests for cognitive functions.

Rey-Osterrieth's Complex Figure test (R-OCF) (Meyers & Meyers, 1995; Strauss et al., 2006) was given in two parts: copy and memory. The subject was given a blank sheet of paper, and another sheet of paper on which the R-OCF was printed. He was instructed to copy the figure as accurately as possible, with no time restrictions. Each time a portion of the drawing was completed, the subject was given a different coloured pencil whilst the examiner was noting the order of colour used. Switching colours affords an adequate record of the subject's strategy. Following completion, the examiner removed both the original and the copied figure. Completion time was recorded. After a retention interval of 30 minutes during which the Verbal Fluency tests, Auditory Verbal Learning test

and Clock Drawing test (CDT) were completed, the subject was again given a blank sheet of paper, and asked to reproduce the R-OCF from memory.

Rey-Osterrieth's Auditory Verbal Learning Test (RAVLT) (Strauss et al., 2006) was given to assess verbal learning and memory. The most commonly used version consists of 15 nouns (List A) that are read aloud (with a 1-s interval between words) for five consecutive trials, each trial followed by a free-recall test. On completion of Trial 5, an interference list of 15 words (List B) is presented, followed by a free-recall test of that list. Immediately after this, delayed recall of the first list is tested (Trial 6) without further presentation of those words. After a 20-minute delay period during which the CDT and R-OCF (memory) were completed, the subject is again required to recall words from List A.

Wechsler Memory Scale (WMS) (Strauss et al., 2006) – The Digits Forward and Digits Backward subtest was used to assess attention and working memory. The number sequences of the WMS are two to eight and two to seven digits, respectively. On hearing them, the subject's task is to repeat them in the exact order in the

‘forward situation’ and in exactly reversed order in the ‘backward situation’. The normal raw score difference between digits forward and digits reversed tends to range a little above 1 (Lezak et al., 2012). The reversed digit span is a measure of working memory and is distinct from the more passive span of apprehension measured by digits forward (Lezak et al., 2012).

Boston Naming Test (BNT) (Kaplan, Goodglass, & Weintraub, 1983). This test consists of 60 large ink drawings of items ranging in familiarity from common (e.g., tree) at the beginning of the test to uncommon (e.g., sphinx) near its end. The subject begins with item 30 and proceeds forward. This test effectively elicits naming deficits.

CDT (Shulman, 2000; Strauss et al., 2006). This test is aimed to measure visual–spatial, constructional and executive function. The subject is asked to draw a clock face with all the numbers and hands and then to state the time ‘ten after eleven’. The strength and the weakness of this test lies in the number of cognitive, motor and perceptual functions required simultaneously for successful completion.

Tests for executive functions.

Wisconsin Card Sorting Test (WCST) (Heaton, Chelune, Talley, Kay, & Curtiss, 1993). The subjects is given one pack of 64 cards in the WCST. One to four instances of each of four symbols in four different colours are printed on each card (e.g., one red triangle, two green stars, three yellow crosses and four blue circles). The subject’s task is to sort the cards according to a principle inferred from the examiner’s responses to the subject’s placement of the cards. The test involves three principles for sorting: colour, symbol or number. The examiner responds only ‘right’ or ‘wrong’ according to the subject’s placement of the card, and does not answer any questions. After 10 consecutive correct placements, the examiner shifts the principle. The test continues until the subject makes six runs of ten correct placements, or when all 64 cards are placed.

Verbal Fluency Test (Galić, 2002). A number of verbal fluency tests, modelled on Thurstone’s Word Fluency Test (Lezak et al., 2012) have been developed to assess more ‘executive’ aspects of verbal behaviour: the ability to think flexibly, switch response sets, self-regulate and self-monitor. Two variants of verbal fluency test were applied: phonemic/letter fluency and semantic/category fluency. In the first test, the subject was asked to produce as many words as possible beginning with the letters K, L and P (Croatian version of the test, according to Galić, 2002) with the exception of proper

names, numbers and words with the same suffix. One minute was allowed for each letter. In the second test, the category fluency test, the subject was asked to name all the animals he can think of in a period of 1 minute.

Design Fluency Test (Jones-Gotman & Milner, 1977). This test was developed as a non-verbal counterpart of Word Fluency Test. The subject is asked to invent drawings that represent neither actual objects nor nameable abstract forms and that are not merely scribbles. After being shown examples of acceptable and unacceptable drawings made by the examiner, the subject is given 5 minutes to make up as many different kinds of drawings as he or she can. Design Fluency Test has been used to assess various aspects of executive functioning.

Stroop Test. A numerical variation of the classical Stroop task paradigm was designed for the present study. Each of the 60 trials consisted of a certain number of digits (between 2 and 6), and subject’s task was to quickly answer how many digits were presented. Answers were collected via keyboard. In congruent tasks, the number and the content of stimuli was equal (e.g., 4 4 4 4), whilst in incongruent trials it differed (e.g., 2 2 2 2 2). The performance of MS was compared to the results of 20 healthy control subjects, undergraduate psychology students (mean age: 20.25 years; $SD = 1.37$) who went through the same procedure. This test is a measure of cognitive control, specifically the ease with which a person can maintain a goal in mind and suppress a habitual response (Strauss et al., 2006).

Tower of Hanoi (computerised version). This test has been considered an ‘executive function’ test focused partly on planning and mostly on inhibiting a prepotent response (the goal–subgoal conflict). The object of the Tower of Hanoi test is to move all the rings from the first column to the third column. To arrive at the best solution (most direct, fewest moves) of the test, the subject must look ahead to determine the order of moves necessary to rearrange five rings of varying sizes. The rings should be moved one by one and a larger ring cannot be placed on top of a smaller ring. This test was also administered to the group of 20 healthy control subjects, male university students (mean age: 21.65 years; $SD = 2.98$), comprised for the present study.

Procedure.

The study was conducted at the Laboratory for Experimental Psychology of the Faculty of Humanities and Social Sciences in Rijeka, Croatia. Neuropsychological assessment was

TABLE 1

Recognition of Emotional Facial Expressions: Percentages of Correct Answers of MS, Calvo and Lundqvist's (2008) Spanish Sample of Participants and Milovanović et al.'s (2013) Croatian Sample

Emotional Expression	MS Initial Assessment*	MS Follow-up Assessment*	Calvo and Lundqvist's (2008) Sample (N = 160)**	Milovanović et al.'s (2013) Sample*** (N = 40)
Anger	90	100	89–98	100–100
Disgust	90	100	86–100	100–100
Fear	80	70	79–98	60–100
Happiness	100	100	97–100	100–100
Sadness	100	100	90–100	80–80
Surprise	100	100	87–98	80–100
Neutral	90	100	98–100	100–100

*Percentage of correct recognitions.

**Average percentage of correct recognitions per stimulus.

***Interquartile range of correct recognitions.

carried out over three meetings lasting about 45 minutes each, 2 years and 6 months after hospitalisation. On the first meeting, the Emotion identification test, R-OCF: copy and memory, Verbal fluency tests, Auditory Verbal Learning Test and CDT were applied. On the second meeting after 2 weeks, the Emotion identification test, Tower of Hanoi, Design Fluency Test and BNT were applied and, on the last meeting a week later, the WMS, WCST and Stroop Test were applied. Over the period of testing, the patient was cooperative and motivated to complete the tasks.

The study was granted approval by the Research Ethics Committee of the Faculty of Humanities and Social Sciences at the University of Rijeka. The patient expressed his willingness to participate in the study by signing informed consent.

Results

Emotion Recognition

On the facial emotion recognition test no impairment was found, except some difficulties that MS had with the recognition of fearful faces, several of which he inaccurately labelled as surprised. The results of initial and follow-up assessments were similar (Table 1). However, since healthy subjects also most often misclassify fearful faces as surprised and vice versa (Calvo & Lundqvist, 2008; Huang, Tang, Helmeste, Shioiri, & Someya, 2001; Palermo & Coltheart, 2004), MS's ability to recognise facial emotional expressions in general and expression of disgust in particular seems to be intact. MS's results on emotion recognition test were compared to Calvo and Lundqvist's (2008) and Milovanović et al.'s (2013) participants. Calvo and Lundqvist's (2008) sample was comprised of

160 Spanish students (mean age: 22.2 years; $SD = 3.1$), whilst Milovanović et al.'s (2013) consisted of 58 Croatian students (mean age: 23.05; $SD = 3.68$) from the same university that MS attends. When the results of anxious and depressed participants are excluded from Milovanović et al.'s (2013) sample, 40 of 58 subjects remain, and their results together with Calvo and Lundqvist's (2008) subjects are compared to MS's scores in Table 1.

Cognitive Functioning

The patient successfully copied a complex drawing (R-OCF) and used an appropriate strategy for organising the sequence of the drawing process. Also, drawing from memory was successful, which reflects normal non-verbal memory and constructional abilities.

He showed normal (average and above average) performance on verbal learning and memory tasks (RAVLT), as well as on language (BNT) and visual-spatial and constructive tasks (CDT). Since these results did not differ from normal expectations we will not describe them in detail.

He showed slight difficulties in maintaining information in working memory because on the Digit Backwards subtest of the WMS, his raw score was on the lower border (four digits). According to Lezak et al. (2012), scores 4 to 5 can be considered 'within normal limits', but we should expect a better result from someone with the patient's educational background. Of course, we must be careful when interpreting a borderline result.

Executive Functions

On the WCST, MS clearly showed *failure to maintain set* which resulted in zero categories completed. In other words, the patient sorted correctly

five cards in the first category (colour) but failed to complete this category and passed over to another sorting principle. This type of error can reflect an inability to continue using a strategy that is working successfully, or loss of train of thought (Lezak et al., 2012). The failure to maintain set rarely occurs but is often linked to brain lesion (Galić, 2002). Our patient obviously had some difficulties in forming and applying a principle and then continuing to use a successful strategy. This means that he could generally have problems with abstraction and flexible thinking. This inference is also supported by his results on verbal and non-verbal fluency tests.

MS expressed impaired or defective phonemic verbal fluency. His result of a total score of 22 generated words was significantly below the average result obtained by comparison group of the same age and educational level (53 words on average for healthy subjects, age range 15 to 35). His total score on the semantic/category fluency test was almost within the average range for healthy subjects. Generating the names of words beginning with particular letter is more difficult than animal naming. The former task relates to strategies of self-retrieving information from explicit memory and the latter to semantic memory. To put it more clearly, patient MS has more problems with the ability to generate words and keep track of earlier responses simultaneously, which all stand for executive aspects of verbal behaviour, than with his retrieval capacities. Although impaired verbal fluency may occur with left hemisphere damage from a variety of etiologies, imaging studies have shown that frontal damage disproportionately impairs letter fluency whilst temporal lobe damage has a greater effect on semantic fluency (Birn et al., 2010). Patient MS expresses a similar problem with letter fluency as patients with frontal lobe damage (Lezak et al., 2012).

On the Design Fluency test, he achieved a low score. He successfully drew only three drawings in a 5-minute period (compared to 15 drawings in average for healthy subjects, age range 14 to 55). It has been found that patients with frontal lobe damage tended to have reduced output relative to healthy subjects and patients with posterior lesions (Lezak et al., 2012). Design Fluency was used to assess various aspects of executive functioning, thus MS has again shown difficulties in some aspects of executive functions similar to patients with frontal lobe damage (e.g., productivity, speed, self-monitoring and self-control of own performance, remembering rules, strategy use and creative imagination).

In order to evaluate MS's performance on the Stroop test, his speed and accuracy of respond-

ing were analysed and compared to the performance of the control group. When the results of MS are transformed to Z standard values, his performance on Stroop tasks is clearly impaired. MS was slower ($Z = -2.81$) and made more errors ($Z = -2.02$) than control participants. According to Strauss et al. (2006), patients with brain injury are typically slower to respond on all tasks and more prone to Stroop interference. Thus, we could conclude that our patient had some difficulties with cognitive control.

Performance of MS on the Tower of Hanoi test was also compared to a control group and expressed in Z standard values. Although MS did not require a lot of moves to complete the tasks ($Z = 0.15$), he required more time than control subjects ($Z = -1.92$). This could mean that he had some difficulties with response inhibition and working memory.

Results Summary

To summarise, MS's ability to recognise facial emotional expressions in general and expression of disgust in particular seems to be intact, but he has clearly showed impaired performance on subsets of the executive functions related to the elaboration of plans of actions (rule-finding, set-shifting and self-activation of cognitive strategies/cognitive control). In other cognitive tasks, he showed average or above average performance in accordance with his age and educational level.

Discussion

Analysis of MS's performance on the emotional recognition test revealed that his ability to perceive facial emotions in general and expression of disgust in particular remained intact after unilateral injury of the left insula and basal ganglia. This finding is contrary to the central finding of Calder et al. (2000) who demonstrated that unilateral lesions of the left insula and putamen result in deficits in recognition of facial disgust. However, the findings of the present study are in confluence with several reports that activation of the right insula is important for the recognition of disgusted faces (Anderson et al., 2003; Phillips et al., 1997; Phillips et al., 2004; Ruiz et al., 2013; Schroeder et al., 2004). Our finding is not in contradiction to the report of Adolphs et al. (2005) who claimed that bilateral insula lesions resulted in impaired ability to recognise facial disgust. When the results of Calder et al. (2000) are re-analysed in detail, their subject did not lose the ability to recognise disgust, since his performance on the Ekman and Friesen's (1976) PFA and Matsumoto and

Ekman's (1988) Japanese and Caucasian Facial Expressions of Emotion (JACFEE) stimuli was above guessing by chance. On both tests, he managed to correctly recognise facial disgust in 50% of trials, whilst the expected accuracy in the case of guessing by chance would be only around 14.3–16.7% since there were six–seven possible answers in the forced-choice procedure used by Calder et al. (2000). Moreover, his Z-score on recognition of disgust in Ekman and Friesen's (1976) PFA stimuli equaled $Z = -1.68$, which means that his performance was better than about 10% of controls. Therefore, our findings support the hypothesis that the insula-striatal system is involved in recognising signals of disgust in others and that the right insula is crucial in this process.

On the other hand, MS showed preserved intellectual function in general, but impairment on subsets of the executive functions. According to Elliott (2003), the executive dysfunctions associated with basal ganglia disorders provide neuropsychological evidence that frontostriatal circuitry, rather than discrete prefrontal regions, may be important in mediating executive function. This case contributes to evidence that striatal structures are important for the executive functions. Our patient has shown the similar pattern of impairment of executive functions, with emphasis on set-shifting and ability to activate or generate cognitive strategies, to that found in patients with caudate lesions (Levy & Dubois, 2006). Executive function deficits, therefore, appear to be a genuine concomitant of basal ganglia damage (Elliott, 2003). The dorsolateral PFC is tightly connected with the caudate nucleus. Electrophysiological studies focusing on the head of caudate nuclei have demonstrated patterns of neural activation similar to those observed in the dorsolateral PFC during working memory or sequencing tasks (Levy & Dubois, 2006). Also, in functional imaging, activation in the dorsal portion of the head of caudate nucleus was found during working memory tasks and during planning tasks (Levy & Dubois, 2006; Postuma & Dagher, 2006). The obtained results are in line with Levy and Dubois (2006) and Bonelli and Cummings' (2007) explanations of cognitive processing related to the dorsolateral prefrontal circuit and impairment in specific subsets of the executive functions associated with cognitive inertia, such as self-generation of rules and set-shifting as well as strategies to retrieve and organise information. Furthermore, Levy and Dubois (2006) suggested that specific cognitive tasks such as the WCST (rule-finding, maintenance and set-shifting), the Tower of Hanoi task (planning) or the letter fluency task (self-activation of cognitive strategies) can be used to detect this cognitive inertia (man-

ifested as weakened incentive or apathy). Patients with dorsolateral PFC dysfunction have difficulty focusing and sustaining attention, generating hypotheses, and maintaining or shifting sets in response to changing task demands. According to this, we could conclude that our patient, as we assumed, had impaired corticostriatal connectivity due to his brain injury.

Conclusion

MS's ability to perceive facial emotions in general and expression of disgust in particular remained intact in spite of unilateral injury of the left insula and basal ganglia. Also, MS showed preserved intellectual function in general, but he experienced difficulties on subsets of the executive functions. Impaired executive functions could lead to very serious problems such as impaired capacity to initiate activity, decreased motivation and defects in planning and carrying out activity sequences as a part of goal-directed behaviours (Lezak et al., 2012). Because executive functions are often impaired but overlooked in patients with basal ganglia damage, it is of crucial importance to attentively define the possible impairment of executive functions and to help the patient to cope with such disabilities.

Acknowledgements

We would like to thank the participant for his time and willingness to cooperate.

Financial Support

This research was supported by the University of Rijeka grant as a part of the research project: Personality, emotions and social processes as determinants of health outcomes (Research project number: 13.04.1.2.01).

Conflict of Interest

None.

Ethical Standards

The authors assert that all procedures contributing to this work comply with the ethical standards of the relevant national and institutional committees on human experimentation and with the Helsinki Declaration of 1975, as revised in 2008.

References

- Adolphs, R., Damasio, H., Tranel, D., Cooper, G., & Damasio, A.R. (2000). A role for somatosen-

- sory cortices in the visual recognition of emotion as revealed by three-dimensional lesion mapping. *The Journal of Neuroscience*, 20(7), 2683–2690.
- Adolphs, R., Tranel, D., Koenigs, M., & Damasio, A.R. (2005). Preferring one taste over another without recognizing either. *Nature Neuroscience*, 8(7), 860–861.
- Alexander, G.E., DeLong, M.R., & Strick, P.L. (1986). Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annual Review of Neuroscience*, 9(1), 357–381.
- Anderson, A.K., Christoff, K., Panitz, D., De Rosa, E., & Gabrieli, J.D. (2003). Neural correlates of the automatic processing of threat facial signals. *The Journal of Neuroscience*, 23(13), 5627–5633.
- Backman, L., Robins-Wahlin, T.B., Lundin, A., Ginovart, N., & Farde, L. (1997). Cognitive deficits in Huntington's disease are predicted by dopaminergic PET markers and brain volumes. *Brain*, 120(12), 2207–2217.
- Birn, R.M., Kenworthy, L., Case, L., Caravella, R., Jones, T.B., Bandettini, P.A., & Martin, A. (2010). Neural systems supporting lexical search guided by letter and semantic category cues: A self-paced overt response fMRI study of verbal fluency. *Neuroimage*, 49(1), 1099–1107.
- Bonelli, R.M., & Cummings, J.L. (2007). Frontal-subcortical circuitry and behavior. *Dialogues in Clinical Neuroscience*, 9(2), 141.
- Calder, A.J., Keane, J., Manes, F., Antoun, N., & Young, A.W. (2000). Impaired recognition and experience of disgust following brain injury. *Nature Neuroscience*, 3(11), 1077–1078.
- Calvo, M.G., & Lundqvist, D. (2008). Facial expressions of emotion (KDEF): Identification under different display-duration conditions. *Behavior Research Methods*, 40(1), 109–115.
- Ekman, P., & Friesen, W.V. (1976). *Pictures of facial affect*. Palo Alto, CA: Consulting Psychologists Press.
- Elliott, R. (2003). Executive functions and their disorders. *British Medical Bulletin*, 65(1), 49–59.
- Galić, S. (2002). *Neuropsihološka procjena: Testovi i tehnike*. Jastrebarsko: Naklada Slap.
- Heaton, R.K., Chelune, G.J., Talley, J.L., Kay, G.G., & Curtiss, J. (1993). *Wisconsin card sorting test manual*. Odessa, FL: Psychological Assessment Resources.
- Huang, Y., Tang, S., Helmeste, D., Shioiri, T., & Someya, T. (2001). Differential judgement of static facial expressions of emotions in three cultures. *Psychiatry and Clinical Neurosciences*, 55(5), 479–483.
- Johnson, S.A., Stout, J.C., Solomon, A.C., Langbehn, D.R., Aylward, E.H., Cruce, C.B., ... & Predict-HD Investigators of the Huntington Study Group. (2007). Beyond disgust: Impaired recognition of negative emotions prior to diagnosis in Huntington's disease. *Brain*, 130(7), 1732–1744.
- Jones-Gotman, M., & Milner, B. (1977). Design fluency: The invention of nonsense drawings after focal cortical lesions. *Neuropsychologia*, 15(4–5), 653–674.
- Kaplan, E., Goodglass, H., & Weintraub, S. (1983). *The Boston Naming Test*. Philadelphia: Lea & Febiger.
- Levy, R., & Dubois, B. (2006). Apathy and the functional anatomy of the prefrontal cortex-basal ganglia circuits. *Cerebral Cortex*, 16(7), 916–928.
- Lezak, M.D., Howieson, D.B., Bigler, E.D., & Tranel, D. (2012). *Neuropsychological assessment*. New York: Oxford University Press.
- Lichter, D.G., & Cummings, J.L. (Eds.) (2001). *Frontal-subcortical circuits in psychiatric and neurological disorders*. New York: The Guilford University Press.
- Lundqvist, D., Flykt, A., & Öhman, A. (1998). *The Karolinska Directed Emotional Faces - KDEF*, CD ROM from Department of Clinical Neuroscience, Psychology section, Karolinska Institutet, ISBN 91-630-7164-9.
- Matsumoto, D., & Ekman, P. (1988). *Japanese and Caucasian facial expressions of emotion (JACFEE) and neutral faces*. San Francisco: Department of Psychiatry, University of California.
- Meyers, J.E. & Meyers, K.R. (1995). *Rey complex figure test and recognition trial : RCFT : professional manual*. Odessa: PAR.
- Middleton, F.A., & Strick, P.L. (2002). Basal-ganglia 'projections' to the prefrontal cortex of the primate. *Cerebral Cortex*, 12(9), 926–935.
- Milovanović, T., Švegar, D., & Kardum, I. (2013). Efekti anksioznosti i depresivnosti na prepoznavanje emocionalnih izraza lica. *Psihološki teme*, 22(1), 29–49.
- Palermo, R., & Coltheart, M. (2004). Photographs of facial expression: Accuracy, response times, and ratings of intensity. *Behavior Research Methods, Instruments, & Computers*, 36(4), 634–638.
- Phillips, M.L., Williams, L.M., Heining, M., Herba, C.M., Russell, T., Andrew, C., ... Gray, J.A. (2004). Differential neural responses to overt and covert presentations of facial expressions of fear and disgust. *Neuroimage*, 21(4), 1484–1496.
- Phillips, M.L., Young, A.W., Senior, C., Brammer, M., Andrew, C., Calder, A.J., ... David, A.S. (1997). A specific neural substrate for perceiving facial expressions of disgust. *Nature*, 389(6650), 495–498.
- Postuma, R.B., & Dagher, A. (2006). Basal ganglia functional connectivity based on a meta-analysis of 126 positron emission tomography and functional magnetic resonance imaging publications. *Cerebral Cortex*, 16(10), 1508–1521.
- Ruiz, S., Lee, S., Soekadar, S.R., Caria, A., Veit, R., Kircher, T., ... Sitaram, R. (2013). Acquired self-control of insula cortex modulates emotion recognition and brain network connectivity in schizophrenia. *Human Brain Mapping*, 34(1), 200–212.
- Salmon, D.P., Heindel, W.C., & Hamilton, J.M. (2001). Cognitive abilities mediated by frontal-subcortical

- circuits. In D.G. Lichter and J.L. Cummings (Eds.), *Frontal-subcortical circuits in psychiatric and neurological disorders* (pp. 114–150). New York: The Guilford University Press.
- Schroeder, U., Hennenlotter, A., Erhard, P., Haslinger, B., Stahl, R., Lange, K.W., & Ceballos-Baumann, A.O. (2004). Functional neuroanatomy of perceiving surprised faces. *Human Brain Mapping, 23*(4), 181–187.
- Shulman, K.I. (2000). Clock-drawing: is it the ideal cognitive screening test?. *International Journal of Geriatric Psychiatry, 15*(6), 548–561.
- Sprengelmeyer, R., Rausch, M., Eysel, U.T., & Przuntek, H. (1998). Neural structures associated with recognition of facial expressions of basic emotions. *Proceedings of the Royal Society of London B: Biological Sciences, 265*(1409), 1927–1931.
- Strauss, E., Sherman, E.M.S., & Spreen, O. (2006). *A compendium of neuropsychological tests. Administration, norms, and commentary*. Oxford: Oxford University Press.