

Ambulatory Eye Surgery and Antithrombotic Therapy - New Approaches

Šimurina, Tatjana; Danilović Luketić, Marija; Graf Župčić, Sandra;
Mraović, Boris

Source / Izvornik: **Acta clinica Croatica, 2021, 60., 749 - 756**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.20471/acc.2021.60.04.23>

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:184:228305>

Rights / Prava: [Attribution-NonCommercial-NoDerivatives 4.0 International/Imenovanje-Nekomercijalno-Bez prerada 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2024-06-29**



Repository / Repozitorij:

[Repository of the University of Rijeka, Faculty of
Medicine - FMRI Repository](#)





AMBULATORY EYE SURGERY AND ANTITHROMBOTIC THERAPY – NEW APPROACHES

Tatjana Šimurina^{1,2,3}, Marija Danilović Luketić^{1,4}, Sandra Graf Župčić^{5,6} and Boris Mraović⁷

¹Department of Anesthesiology, Resuscitation, Intensive Medicine and Pain Management, Faculty of Medicine, Josip Juraj Strossmayer University of Osijek, Osijek, Croatia;

²Department of Health Studies, University of Zadar, Zadar, Croatia;

³Department of Anesthesiology, Resuscitation and Intensive Care Medicine, Zadar General Hospital, Zadar, Croatia;

⁴Department of Anesthesiology, Resuscitation and Intensive Care Medicine, Požega General County Hospital, Požega, Croatia;

⁵School of Medicine, University of Rijeka, Rijeka, Croatia;

⁶Department of Neurology, Rijeka University Hospital Center, Rijeka, Croatia;

⁷Department of Anesthesiology and Perioperative Medicine, School of Medicine, University of Missouri, Columbia, USA

SUMMARY – One of the most common surgeries in elderly patients is eye surgery. An increasing number of patients undergoing ambulatory eye surgery are on antithrombotic therapy. These drugs may increase the risk of perioperative bleeding associated with ophthalmic needle blocks and/or eye surgery. Intraoperative bleeding and postoperative hemorrhagic complications may lead to the loss of vision or even eyes. On the other hand, stopping anticoagulants and antiplatelets before the surgery may increase the risk of thrombotic events with potentially life-threatening complications. The aim of this narrative review is to provide a systematic review of the published evidence for the perioperative antithrombotic management of patients undergoing different types of eye surgery in ambulatory settings. A comprehensive review of the English-language medical literature search utilizing PubMed, Ovid Medline® and Google Scholar from January 2015 to December 2018 was performed. The database searches included studies providing evidence relevant to ambulatory eye surgery and perioperative antiplatelet medications and anticoagulants. Updated recommendations will be given for continuation, discontinuation, and modification of antithrombotic agents in order to optimize the management of antithrombotic therapies in outpatients scheduled for eye surgery.

Key words: *Surgical procedure, ophthalmologic; Ambulatory surgery; Anesthesia, regional; Anticoagulants; Agents, antiplatelet*

Introduction

Eye surgery is increasingly performed as outpatient surgery over the last couple of decades. Phacoemulsification has contributed greatly to the transition from inpatient to outpatient ophthalmic surgery¹. Other fac-

tors are introduction of minimally invasive surgery, laser, ultrasound and robotic techniques, improved pain control, greater attention on patient outcomes, broader indications for regional anesthesia that has almost completely replaced general anesthesia, and clearly defined discharge criteria^{2,3}. Longer eye surgery procedures with a high risk of perioperative complications are not appropriate for ophthalmic ambulatory anesthesia.

Patients undergoing eye surgery are predominantly elderly with significant comorbidities and often take

Correspondence to: *Marija Danilović Luketić, MD*, Požega General County Hospital, Osječka 107, HR-34000 Požega, Croatia
E-mail: mdanilov11@gmail.com

Received August 19, 2019, accepted October 9, 2019

multiple medications including antithrombotic drugs. The use of antithrombotic drugs has significantly increased as the prevalence of cardiovascular diseases among aging people has increased. Surgical bleeding and hemorrhagic complications of ophthalmic needle blocks could be catastrophic for visual function. Discontinuation of antithrombotic agents, dose reduction, and/or low molecular weight heparin (LMWH) bridging may reduce the risk of bleeding related to eye surgery and needle based ophthalmic blocks, but at the expense of an increased risk of life-threatening thromboembolic events.

Guidelines have been published to address the issues of the use of antithrombotic drugs in the broader context of general and regional anesthesia, as well as specific recommendations for ophthalmic regional anesthesia^{4,7}. Recommendations for ophthalmic patients on antithrombotic therapy undergoing eye surgery are based on observational studies, retrospective and prospective cohort studies, case series, case reports, and expert reports. Prospective randomized trials are lacking because such studies would be considered unethical. There are a few factors that have recently emerged in deciding on discontinuation of antithrombotic drugs prior to eye surgery in order to minimize perioperative bleeding, i.e. an increasing number of patients on new-generation antithrombotic agents, advances in regional ophthalmic anesthesia and surgical techniques, changing demographic characteristics of ophthalmic patients, and increased awareness of patient outcomes. Moreover, there is sufficient evidence for the prothrombotic phenomenon, the so-called 'rebound' effect following antithrombotic withdrawal, that may be responsible for systemic adverse events⁸. However, there is paucity of studies examining the balance between the risk of perioperative hemorrhage and the risk of thromboembolic complications in association with newer oral antithrombotic agents in outpatients undergoing eye surgery.

We hypothesized that current recommendations would need updating in accordance to newly published clinically relevant studies. The objective of this narrative review is to summarize current knowledge based on the recently published peer-reviewed medical articles with respect to continuation, discontinuation, and modification of antithrombotic agents in order to achieve an optimal approach for patients on anti-

thrombotic treatment who are scheduled to undergo eye surgery in ambulatory settings.

Material and Methods

Two independent reviewers searched for medical literature written in English on PubMed, Ovid Medline® and Google Scholar. Peer-reviewed articles, systematic or narrative review articles, meta-analyses, prospective randomized studies, retrospective cohort studies, observational studies, case reports and case series were systematically searched for, through the time period from January 2015 to December 2018. Other types of publications such as editorials, letters to editors, animal experiments, studies in children, and data available only in abstracts were excluded. The search of databases included older and newer anticoagulants and antiplatelet drugs, the risk of intraoperative surgical bleeding during different types of elective same-day ophthalmic procedures, and hemorrhagic complications of ophthalmic regional anesthesia (topical anesthesia with intracameral local anesthetic, needle blocks [intraconal or extraconal] or cannula sub-Tenon's [episcleral] blocks) in outpatients on antithrombotic agents. Further, we searched for thromboembolic events associated with cessation or modification of the antithrombotic treatment regimens. We searched the following Medical Subject Headings (MeSH) terms: Surgical Procedure, Ophthalmologic; Surgery, Outpatient; Ambulatory Surgery Procedures; Anesthesia, Regional; Anticoagulants; Agents, Antiplatelet; Aspirin; Warfarin; Clopidogrel; Heparin, Low-Molecular-Weight; Thrombin Inhibitors, Direct; Direct Factor Xa Inhibitors; Thrombosis and Embolism.

Results

We identified 20 articles (9 review articles, 2 systematic reviews with meta-analysis of randomized clinical trials (RCTs), 1 prospective randomized study, 3 retrospective analyses, 1 observational study, 2 case reports, and 2 case series) with relevant information on significant ocular hemorrhagic complications and thromboembolic adverse events in patients with anticoagulant (Acoag) and antiplatelet (Aptl) therapy. Out of 20 articles, 11 studies met the eligibility criteria (8 review articles, 1 retrospective analysis, 1 prospective randomized

study, and 1 case report) and were selected to summarize updated information on intraoperative bleeding, postoperative hemorrhagic complications, and arterial or venous thromboembolic events. Critical appraisal was performed based on the selected studies and summarized updated recommendations are presented.

Discussion

We reviewed and analyzed new evidence from recently published literature and summarized optimal approaches to ophthalmic patients on antithrombotic treatment that are scheduled for eye surgery under local ophthalmic anesthesia in ambulatory settings.

Eye surgery is currently the most common procedure among elderly patients and is often performed as outpatient surgery in ambulatory surgical centers. The most common procedures are cataract extraction, strabismus repair, glaucoma surgery, simple vitrectomy, minimal plastic surgery, nasolacrimal duct probing, chalazion excision, and eye examinations such as tonometry, etc. The vast majority of ocular surgeries are procedures under topical anesthesia with intracameral application of local anesthetics and needle or cannula based ophthalmic blocks⁹. When local application or regional blocks are not feasible, e.g., uncooperative or anxious patients, fast-track general anesthesia has been a safe choice even in frail elderly patients who are often polymedicated and usually on antithrombotic therapy. Often indications for antithrombotic drugs are stroke prevention in atrial fibrillation (AF), management and prevention of thromboembolism, *in situ* prosthetic heart valves, treatment of acute coronary syndrome, and secondary prevention of cardiovascular disease. More than 28% of ophthalmic patients take aspirin, 2% take clopidogrel, and around 5% take various anticoagulants¹⁰. In general, eye surgery is considered a low-risk surgery with a cardiac risk less than 1%. However, many complications may occur in ophthalmic patients undergoing eye surgery who are on chronic antiplatelet and anticoagulant therapy. Surgical bleeding and hemorrhagic complications related to sharp needle (intraconal or extraconal) or cannula (episcleral) based blocks may have sight-threatening complications if antithrombotic therapy is continued. Even small intraoperative and postoperative bleeding may present a severe risk due to limited and not ex-

pandable space in the orbit and compartments of the eye globe.

Risk of sight-threatening bleeding in eye surgery

Predisposing factors to perioperative hemorrhage include increasing age, comorbidities (liver failure, renal failure, anemia, cardiac stent, uncontrolled hypertension), history of bleeding disorders or thromboembolic events, family history of bleeding and clotting disorders, medications used (antithrombotic drugs, steroids) and herbal treatments, eye characteristics (myopic eye with staphyloma), and type of eye surgery. Decision on cessation or continuation of antithrombotic drugs, or the use of bridging therapy with unfractionated heparin (UFH) or LMWH should be based on bleeding risk prediction and stratification of the risk of sight-threatening hemorrhagic complications.

Type of regional ophthalmic blocks

The type of regional ophthalmic blocks and patient-related risk factors are used to stratify the risk of sight-threatening bleeding to low, moderate and high risk if antithrombotic therapy is continued perioperatively¹¹ (Table 1). According to the recommendation shown in Table 1, there are no significant bleeding risks in peribulbar and retrobulbar anesthesia for most ophthalmic patients on Aptl and/or Acoag treatment except for the case of mechanical heart valves or a combination of Aptl and Acoag therapy in a patient who has only one eye. Moreover, the risk of bleeding may be reduced by single shot inferonasal puncture into poor vascular tissue with a narrow and short needle through a small incision. Local anesthesia with peribulbar, retrobulbar or episcleral block can be performed safely not only in otherwise healthy ophthalmic patients without antithrombotic therapy but even in those with renal or liver failure or coagulopathy while antithrombotic treatment is maintained.

Hemorrhagic complications related to ophthalmic blocks are usually assessed on the 4-grade scale proposed by Kallio *et al.*¹² as mild (grade 1 [spot ecchymosis] or grade 2 [lid ecchymosis involving half of the lid surface area or less]), moderate (grade 3 [lid ecchymosis all around the eye]), and severe (grade 4 [retrobulbar hemorrhage with increased intraocular pressure]).

Three non-randomized controlled trials by Calenda *et al.*¹³⁻¹⁵ and two prospective cohorts by Kallio *et al.*¹² and Katz *et al.*¹⁶ from a recent systematic review by

Table 1. Risk stratification for sight-threatening bleeding in ophthalmic needle and cannula blocks

Risk			Regional ophthalmic blocks	
High	Comorbidity, anticoagulant /antiplatelet therapy		Peribulbar/retrobulbar	Episcleral (sub-Tenon's block)
Moderate				
Low				
	ASA I, no therapy		Low	Low
	Liver/renal failure, coagulopathy		Moderate	Low
Aptl	Prophylaxis (single Aptl)	Primary	Low	Low
		Secondary	Low	Low
	Dual Aptl	Stop one Aptl	Moderate	Low
		Dual Aptl	Moderate	Low
Acoag (VKA: target INR 2.5)	AF		Moderate	Low
	PE, DVT		Moderate	Low
	Long-term for recurrent PE/DVT /high risk of stroke		Moderate	Low
Acoag +/- Aptl t (VKA: target INR 3.5)	MHV/Acoag+Aptl		Moderate	Moderate
	MHV/Acoag+Aptl + only one eye		High	Moderate

ASA = American Society of Anesthesiologists; INR = international normalized ratio; Aptl = antiplatelet drugs; Acoag = anticoagulant drugs; VKA = vitamin K antagonist; AF = atrial fibrillation; PE = pulmonary embolism; DVT = deep vein thrombosis; MHV = mechanical heart valve (data in Table 1 are based on the results from reference 11).

Takaschima *et al.*¹⁷ did not find severe bleeding related to needle blocks in association with the use of anti-thrombotic drugs. There were no differences regarding mild to moderate hemorrhage (grades 1 to 3) between patients who underwent ophthalmic regional anesthesia and were still taking aspirin, clopidogrel or warfarin, and control group patients who did not take any antithrombotic drug. In a retrospective cohort study by Katz *et al.*¹⁶, the incidence of retrobulbar hemorrhage was only 0.04% (95% confidence interval [CI]: 0.001-0.10) among 14,823 patients with peribulbar or retrobulbar blocks for cataract surgery.

Type of ophthalmic procedures

Antiplatelet medications can be continued for most of ophthalmic surgeries in ambulatory settings. Vigilance is needed regarding a new stronger Aptl, e.g., prasugrel or ticagrelor. A recently published article has reported a case of a 62-year-old female with proliferative diabetic retinopathy on dual Aptl therapy with aspirin and prasugrel for preventing coronary stent thrombosis. She presented for panretinal photocoagulation under retrobulbar block but she developed retrobulbar hemorrhage. Newer, more potent Aptl drugs might carry higher risks of severe bleeding com-

plications in susceptible patients¹⁸. Anesthesiologists and ophthalmologists should consider delay of the surgery until dual Aptl therapy is no longer needed, to hold prasugrel or avoid retrobulbar blocks and choose different regional ophthalmic blocks such as peribulbar block with the use of ultrasound.

In most ophthalmic outpatients, anticoagulants do not significantly increase the risk of severe ocular bleeding if anticoagulation markers are within the therapeutic range¹⁹.

The types of ophthalmic procedures involving a high risk of hemorrhage are orbital surgery and major oculoplastic surgery. An intermediate risk of hemorrhage is present in vitreoretinal surgery, glaucoma surgery, and corneal transplant. A high hemorrhagic risk of eye surgery in combination with sharp needle ophthalmic blocks in patients with serious comorbidities and on antithrombotic therapy demands careful patient evaluation, proper preparation, and it would be reasonable to keep them in hospital overnight for monitoring and evaluation.

Cataract surgery and intravitreal injections

These procedures have a low risk of ocular hemorrhage and cessation of anticoagulation is not necessary

in most patients. All antithrombotic drugs can be continued if the cataract surgery is performed under topical or sub-Tenon's (episcleral) block²⁰. Even if local anesthesia is performed with sharp needles, Aptl and Acoag therapy can be continued provided that the international normalized ratio (INR) for vitamin K antagonists (VKA) is in the therapeutic range. Dual Aptl therapy should be avoided, the new P2Y12 inhibitors should be stopped but acetylsalicylic acid (ASA) could be continued. Direct oral anticoagulants (DOACs) at therapeutic doses should be stopped before and restarted after the cataract surgery under sharp needle blocks. A recent review by Grzybowski *et al.*²¹ included case series by Barequet *et al.*^{22,23}, Salam *et al.*²⁴ and Kobayashi²⁵, and Cataract National Dataset electronic multicenter audit by Benzimira *et al.*¹⁰, and showed that phacoemulsification of cataracts under local anesthesia is not associated with higher risks of severe intraoperative bleeding and postoperative hemorrhage in outpatients taking Aptl and Acoag agents. The risk of sight-threatening bleeding is intermediate in case of anticoagulation in patients with mechanical heart valves (MHV) who take VKA with INR target values of 3.5 undergoing cataract surgery, or who have both Acoag and Aptl therapy and have only one functional eye¹¹. For cataract surgery, DOACs should be continued.

Bridging therapy with UFH or LMWH is unnecessary for most ophthalmic procedures in ambulatory settings. A gradual decrease of the anticoagulation effect is predictable during short-term perioperative cessation of DOACs, but it is influenced by renal function¹⁹. If resuming DOACs carries a risk of postoperative hemorrhage that outweighs the thromboembolic risk, then bridging therapy with reduced prophylactic doses of LMWH may be an option²⁶.

The risk of intraocular bleeding among new users of antithrombotic therapy was tested in a recent large retrospective cohort study with two parallel analyses of data from national insurance claim database by Uyhazi *et al.*²⁷. They compared patients who were on dabigatran or rivaroxaban with those who were on warfarin, and patients with new use of prasugrel with those on clopidogrel. The main outcomes were hazard ratios (HR) of developing intraocular hemorrhage at 90 and 365 days. The novel oral antithrombotic drugs showed a significant decrease in hemorrhage compared with warfarin at 365 days (HR=0.75; 95% CI 0.58-0.97, p=0.03) but not at 90 days (HR=0.73; 95% CI 0.22-

2.63, p=0.13). There were no differences between prasugrel and clopidogrel at 90 days (HR=0.75; 95% CI 0.29-1.92, p=0.55) or 365 days (HR=1.19; 95% CI 0.69-2.04, p=0.53). However, more evidence is needed for safety of novel oral anticoagulants in ophthalmic patients.

Trabeculectomy is the most common glaucoma surgery in ambulatory settings. Overall, glaucoma filtration surgery has an intermediate risk of bleeding. However, the risk is higher in patients on Acoag therapy for AF, pulmonary embolism (PE) or deep venous thrombosis (DVT), as well as in patients with MHV or on both Acoag and Aptl therapy. There is no unique approach to maintain or to hold Acoag and Aptl therapy prior to glaucoma surgery. The majority of eye surgeons do not stop therapy especially in cases with a high thromboembolic risk without increasing the risk of surgical complications. Current recommendations from the literature suggest that ASA should be stopped in cases where the Aptl is the only drug for primary prevention of cardiovascular disease, but they could be continued for secondary prevention. In patients on clopidogrel, it should be stopped, but ASA should be continued or introduced. New oral Aptl should be stopped before glaucoma surgery but ASA continued. Traditional anticoagulants and DOACs should be stopped before glaucoma surgery and bridging therapy should be considered according to patient thrombotic risk^{11,19,28}.

For strabismus repair surgery, the recommendations are similar to those for glaucoma surgery.

A growing number of vitreoretinal surgery (VRS) is performed as ambulatory surgery. The risk of ocular bleeding is higher in patients with liver failure, renal failure, diabetes and coagulopathy than in otherwise healthy patients²⁹. The risk of bleeding is classified as intermediate in case of dual Aptl therapy for coronary stent or Acoag therapy for AF, and anticoagulation with therapeutic range for recurrent PE or DVT, as well as in case of MHV or both Acoag and Aptl therapy. The perioperative risk of ocular bleeding is high in patients with MHV (for VKA, INR target value of 3.5) or those who have both Acoag and Aptl therapy, and in patients who have only one functional eye. Studies of VRS showed controversial results. In general, Aptl drugs could be continued in many patients undergoing VRS in ambulatory settings³⁰. Perioperative management is similar to glaucoma surgery except

for the interruption of clopidogrel being required. Hemorrhagic complications such as retrobulbar, sub-retinal, suprachoroidal or vitreous cavity hemorrhage still can occur. Vigilance is needed in the presence of neovascular retinal diseases²⁸. Acoag therapy may be continued unless the surgeon requests discontinuation. DOACs at therapeutic doses should be stopped before VRS and the duration of withholding depends on renal function. A retrospective cohort study by Grand and Walia³¹ found a low risk of hemorrhagic complications or need for reoperation in patients undergoing VRS while maintaining therapy with rivaroxaban, apixaban, dabigatran or prasugrel.

Oculoplastic surgery that is performed anterior to the orbital septum has a low risk. Minor oculoplastic surgeries with a low risk of hemorrhage such as chalazion, eyelid cyst removal and eyelid lesion removal could be performed in ambulatory settings, and Acoag and Aptl agents may be continued perioperatively. Patients scheduled for oculoplastic surgeries with a high risk of sight-threatening hemorrhage, such as deep orbital surgery, postseptal eyelid surgery and dacryocystorhinostomy should have antithrombotic drugs withheld providing that it is safe to do so. Acoag and Aptl therapy should be withheld in patients who have a low risk of thromboembolism (less than 5% annual risk). If patients are at a high risk of thromboembolism, they should have bridging therapy or continue the anticoagulation to avoid bleeding complications associated with bridging therapy³².

For most ophthalmic day-surgeries, the risk of stopping antithrombotic therapy is higher than the risk of continuing antithrombotic drugs. If interruption of antithrombotic treatment is needed, a modification of therapy should be discussed with a cardiologist, hematologist, neurologist, ophthalmologist, and other specialists³³. The BRIDGE trial, a randomized, double-blind, placebo-controlled study by Douketis *et al.*³⁴, included 1,884 patients with AF on warfarin therapy who underwent elective invasive gastrointestinal, cardiothoracic and orthopedic surgery. The results showed that in patients who had warfarin therapy interrupted before the surgery, no bridging therapy was inferior to bridging therapy with LMWH for prevention of perioperative arterial thromboembolism [G_{NB} , No bridging group=4/918 (0.4%) *vs.* G_B , Bridging group=3/895 (0.3%)]. Further, no bridging therapy was superior to bridging therapy with respect to major

perioperative bleeding [G_{NB} group=12/ 918 (1.3%) *vs.* G_B 29/895 (3.2%), $p<0.005$] and minor bleeding preoperatively [G_{NB} group=110/918 (12.0%) *vs.* G_B 187/895 (20.9%), $p<0.001$]. The bridging therapy increased the risk of bleeding but did not provide any benefit for stroke prevention [G_{NB} =2/918 (0.2%) *vs.* G_B =3/895 (0.3%)]³⁴.

Thrombotic risk

During the preoperative visit, anesthesiologists and ophthalmologists should identify the risk factors for ocular bleeding and the risk of thromboembolic events. The approach to antithrombotic therapy should be individualized for a particular patient scheduled for the specific type of eye surgery¹¹. It is important to be familiar with the indications for antithrombotic therapy, to know when it may be stopped without unnecessary risks. There is substantial evidence that cessation of anticoagulation even for a short perioperative period is associated with significant thrombotic events such as stroke, myocardial infarction, or even death. High-risk patients are those with advanced age, severe comorbidities, thrombophilia, cerebrovascular, coronary artery and peripheral vascular diseases, and those who have mitral prosthetic valves or more than one prosthetic valve. The risks and benefits of continuing or stopping antithrombotic drugs should be discussed with other specialists involved in patient management to make an optimal, individualized approach for perioperative antithrombotic therapy in ophthalmic surgery.

Conclusion

In this review, we discuss the balance between systemic thrombotic risks related to modification of antithrombotic therapy before ophthalmic day-surgery under regional anesthesia and the risk of intraoperative and postoperative bleeding associated with continuation of antithrombotic therapy. Ophthalmic patients require a slightly different approach than other surgical patients depending on the type of eye surgery and antithrombotic treatment. Ophthalmic surgeries are stratified according to the risk of hemorrhagic complications in the perioperative period. Many ophthalmic procedures can be safely performed as outpatient surgery while antithrombotic therapy is continued. For some ophthalmic outpatients, the risk of

stopping Aprotin and Acoag medications may outweigh the risk of perioperative hemorrhage.

There is no unique approach to eye surgery and perioperative use of antithrombotic drugs. The best way is a tailored approach depending on the type of anesthesia and surgical technique and assessment of renal function. The 'one size fits all' approach is not acceptable and individualized approach is highly recommended. Local protocols for ambulatory centers may help minimize cessation of antithrombotic drugs and thrombotic complications while limiting the risk of perioperative hemorrhage and maximize patient satisfaction and safety. Economic pressures for more ambulatory surgeries, increased indications for surgery in elderly people, increased prevalence of antithrombotic therapy, and greater focus on patient outcomes determine the important role of anesthesiologists. Anesthesiologists have become team leaders with a responsibility for perioperative management of ophthalmic patients. Full communication between the patients and prescribing physicians is the key to ensure an optimal outcome for ophthalmic patients on antithrombotic treatment undergoing eye surgery under regional anesthesia in ambulatory settings.

References

1. Stanić R, Bućan K, Stanić-Jurašin K, Kovačić Ž. Phacoemulsification in eyes with posterior polar cataract. *Acta Clin Croat* [Internet]. 2012 [cited 2019 June 01];51(1):55-8. Available from: <https://hrcak.srce.hr/85736>
2. Lukenda A, Karaman Martinović Ž, Kalauz M. Excimer laser correction of hyperopia, hyperopic and mixed astigmatism: past, present, and future. *Acta Clin Croat* [Internet]. 2012 [cited 2019 June 01];51(2):299-304. Available from: <https://hrcak.srce.hr/106658>
3. Kovačević D, Mance T, Mišljenović T. Combined sutureless procedure: phacoemulsification with intraocular lens implantation and 25 gauge pars plana vitrectomy. *Acta Clin Croat* [Internet]. 2008 [cited 2019 June 01];47(Suppl 1):31-4. Available from: <https://hrcak.srce.hr/22762>
4. McCaughan B, Marr C. Clinical Excellence Commission, 2018, Guidelines on Perioperative Management of Anticoagulant and Antiplatelet Agents [Internet]. Sydney: Clinical Excellence Commission. 2018 [cited 2019 June 02]; Available from: http://www.cec.health.nsw.gov.au/_data/assets/pdf_file/0006/458988/Guidelines-on-perioperative-management-of-anticoagulant-and-antiplatelet-agents.pdf
5. Horlocker TT, Vandermeulen E, Kopp SL, Gogarten W, Lefert LR, Benzon HT. Regional Anesthesia in the Patient Receiving Antithrombotic or Thrombolytic Therapy: American Society of Regional Anesthesia and Pain Medicine Evidence-Based Guidelines (Fourth Edition). *Reg Anesth Pain Med*. 2018;43(3):263-309. doi: 10.1097/AAP.0000000000000763
6. Kumar CM, Eke T, Dodds C, *et al.* Local anaesthesia for ophthalmic surgery – new guidelines from the Royal College of Anaesthetists and the Royal College of Ophthalmologists. *Eye (Lond)*. 2012;26(6):897-8. doi: 10.1038/eye.2012.82
7. Kong K-L, Budd J; Chapter Development Group Members. Chapter 13: Guidelines for the Provision of Anaesthesia Services (GPAS); Guidelines for the Provision of Ophthalmic Anaesthesia Services 2019 [Internet]. London: Royal College of Anaesthetists (RCoA). 2019 [cited 2019 June 02]; Available from: <https://www.rcoa.ac.uk/system/files/gpas-2019-13-ophthal.pdf>
8. Sambu N, Warner T, Curzen N. Clopidogrel withdrawal: is there a "rebound" phenomenon? *Thromb Haemost*. 2011;105(2):211-20. doi: 10.1160/TH10-08-0554
9. Palte HD. Ophthalmic regional blocks: management, challenges, and solutions. *Local Reg Anesth* 2015;8:57-70. doi: 10.2147/LRA.S64806 eCollection 2015.
10. Benzimra JD, Johnston RL, Jaycock P, *et al.*; the EPR User Group. The Cataract National Dataset electronic multicentre audit of 55 567 operations: antiplatelet and anticoagulant medications. *Eye*. 2009;23(1):10-6. doi: 10.1038/sj.eye.6703069
11. Kiire CA, Mukherjee R, Ruparelia N, Keeling D, Prendergast B, Norris JH. Managing antiplatelet and anticoagulant drugs in patients undergoing elective ophthalmic surgery. *Br J Ophthalmol*. 2014;98(10):1320-4. doi: 10.1136/bjophthalmol-2014-304902
12. Kallio H, Paloheimo M, Maunuksela EL. Haemorrhage and risk factors associated with retrobulbar/peribulbar block: a prospective study in 1383 patients. *Br J Anaesth*. 2000;85(5):708-11. doi: 10.1093/bja/85.5.708
13. Calenda E, Cardon-Guiton A, Genevois O, Gueudry J, Muraine M. Peribulbar block in 500 patients scheduled for eye procedures and treated with acetyl salicylic acid. *Acta Anaesthesiol Taiwan*. 2011;49(4):141-3. doi: 10.1016/j.aat.2011.11.003
14. Calenda E, Lamothe L, Genevois O, Cardon A, Muraine M. Peribulbar block in patients scheduled for eye procedures and treated with clopidogrel. *J Anesth*. 2012;26(5):779-82. doi: 10.1007/s00540-012-1406-6
15. Calenda E, Genevois O, Cardon A, Muraine M. Peribulbar anesthesia in 750 patients treated with oral anticoagulants. *Int J Ophthalmol*. 2014;7(1):110-3. doi: 10.3980/j.issn.2222-3959.2014.01.20
16. Katz J, Feldman MA, Bass EB, *et al.*; Study of Medical Testing for Cataract Surgery Team. Risks and benefits of anticoagulant and antiplatelet medication use before cataract surgery. *Ophthalmology*. 2003;110(9):1784-8. doi: 10.1016/S0161-6420(03)00785-1
17. Takaschima A, Marchioro P, Sakae TM, Porporatti AL, Mezzomo LA, De Luca Canto G. Risk of hemorrhage during

- needle-based ophthalmic regional anesthesia in patients taking antithrombotics: a systematic review. *PLoS One*. 2016;11(1): e0147227. doi: 10.1371/journal.pone.0147227
18. Matharu KS, Smith SV, Lee AG. Retrobulbar hemorrhage and prasugrel. *Can J Ophthalmol*. 2016;51(6):e189-e190. doi: 10.1016/j.jcjo.2016.07.003
 19. Bonhomme F, Hafezi F, Boehlen F, Habre W. Management of antithrombotic therapies in patients scheduled for eye surgery. *Eur J Anaesthesiol*. 2013;30:449-54. doi: 10.1097/EJA.0b013e328360c442
 20. Li Q, Qian Y, Zhang Y, Sun G, Zhou X, Wang Z. Continuation of aspirin therapy before cataract surgery with different incisions: safe or not? *J Ophthalmol*. 2018;2018:6543937. doi: 10.1155/2018/6543937. eCollection 2018.
 21. Grzybowski A, Ascaso FJ, Kupidura-Majewski K, Packer M. Continuation of anticoagulant and antiplatelet therapy during phacoemulsification cataract surgery. *Curr Opin Ophthalmol*. 2015;26(1):28-33. doi: 10.1097/ICU.0000000000000117
 22. Barequet IS, Sachs D, Priel A, *et al.* Phacoemulsification of cataract in patients receiving Coumadin therapy: ocular and hematologic risk assessment. *Am J Ophthalmol*. 2007;144(5):719-23. doi: 10.1016/j.ajo.2007.07.029
 23. Barequet IS, Sachs D, Shenkman B, *et al.* Risk assessment of simple phacoemulsification in patients on combined anticoagulant and antiplatelet therapy. *J Cataract Refract Surg*. 2011;37(8):1434-8. doi: 10.1016/j.jcrs.2011.02.035
 24. Salam T, Raines MF. Clopidogrel and intraocular surgery: complicated or uncomplicated? *Grand Rounds*. 2007;7:58-60. doi: 10.1102/1470-5206.2007.0017
 25. Kobayashi H. Evaluation of the need to discontinue antiplatelet and anticoagulant medications before cataract surgery. *J Cataract Refract Surg*. 2010;36(7):1115-9. doi: 10.1016/j.jcrs.2010.01.017
 26. Tan LT, Norris J, Prendergast B. The management of antiplatelets and anticoagulation in elective ophthalmic surgery. *Eye News* [Internet]. 2016 [cited 2019 June 03]; 22(6):1-3. Available from: <https://www.eyenews.uk.com/media/9179/eyeam16-toptips.pdf>
 27. Uyhazi KE, Miano T, Pan W, VanderBeek BL. Association of novel oral antithrombotics with the risk of intraocular bleeding. *JAMA Ophthalmol*. 2018;136(2):122-30. doi: 10.1001/jamaophthalmol.2017.5677
 28. Kong K-L, Khan J. Ophthalmic patients on antithrombotic drugs: a review and guide to perioperative management. *Br J Ophthalmol*. 2015;99(8):1025-30. doi: 10.1136/bjophthalmol-2014-306036
 29. Fabinyi DC, O'Neill EC, Connell PP, Clark JB. Vitreous cavity haemorrhage post-vitreectomy for diabetic eye disease: the effect of perioperative anticoagulation and antiplatelet agents. *Clin Exp Ophthalmol*. 2011;39(9):878-84. doi: 10.1111/j.1442-9071.2011.02575.x
 30. Mantopoulos D, Vavvas DG, Fine HF. Perioperative risks of antiplatelet and anticoagulant drugs in vitreoretinal procedures. *Ophthalmic Surg Lasers Imaging Retina*. 2017;48(1):4-8. doi: 10.3928/23258160-20161219-01
 31. Grand MG, Walia HS. Hemorrhagic risk of vitreoretinal surgery in patients maintained on novel oral anticoagulant therapy. *Retina*. 2016;36(2):299-304. doi: 10.1097/IAE.00000000000000783
 32. Esparaz ES, Sobel RK. Perioperative management of anticoagulants and antiplatelet agents in oculoplastic surgery. *Curr Opin Ophthalmol*. 2015;26(5):422-8. doi: 10.1097/ICU.000000000000187
 33. Stuart A. Perioperative management of antithrombotics. *EyeNet Magazine* [Internet]. 2016 [cited 2019 June 03]; 35-7. Available from: <https://www.aao.org/eyenet/article/perioperative-management-of-antithrombotics>
 34. Douketis JD, Spyropoulos AC, Kaatz S, *et al.* Perioperative bridging anticoagulation in patients with atrial fibrillation. *N Engl J Med*. 2015;373(9):823-33. doi: 10.1056/NEJMoa1501035

Sažetak

JEDNODNEVNA OČNA KIRURGIJA I ANTIKOAGULANTNA TERAPIJA – NOVIJI PRISTUPI

T. Šimurina, M. Danilović Luketić, S. Graf Župčić i B. Mraović

U populaciji bolesnika starije dobi očni kirurški zahvati su jedni od najčešćih kirurških zahvata. Sve više bolesnika kojima je potreban kirurški zahvat na očima su starije dobi i većinom su na kroničnoj terapiji lijekovima uključujući antitrombocitne lijekove. Ti lijekovi mogu povećati rizik od perioperacijskog krvarenja prilikom izvođenja regionalnih očnih blokova ili kirurškog zahvata. Krvarenje tijekom operacije oka i hemoragijske komplikacije poslije zahvata mogu dovesti do gubitka vidne funkcije ili čak samog oka. S druge strane, prekidanje uzimanja antitrombocitnih i antikoagulacijskih lijekova prije kirurškog zahvata dovodi do povećanog rizika za nastanak ozbiljnih i za život opasnih tromboembolijskih komplikacija. Cilj ovoga narativnog preglednog članka je sustavni pregled objavljenih dokaza o perioperacijskom antitrombotskom liječenju očnih bolesnika planiranih za različite zahvate u dnevnoj očnoj kirurgiji. Pretražene su baze medicinskih podataka pomoću PubMed, Ovid Medline® i Google Scholar za razdoblje od siječnja 2015. godine do prosinca 2018. godine. Obuhvaćene su studije relevantne za planirane očne operacije u jednodnevnoj kirurgiji i perioperacijsko liječenje antitrombocitnim i antikoagulacijskim lijekovima s naglaskom na sadašnje stavove u pogledu nastavka, prekida ili modifikacije antitrombotske terapije kako bi se pospješila priprema bolesnika za očne zahvate.

Ključne riječi: *Kirurški zahvati, oftalmološki; Dnevna kirurgija; Anestezija, regionalna; Antikoagulacijski lijekovi; Antitrombocitni lijekovi*