

Molecular aspects of immunological evasion by cytomegalovirus

Crnković, Irena; Koszinowski, Ulrich

Source / Izvornik: **Periodicum biologorum, 1996, 98, 175 - 182**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:184:662925>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-01-07**



Repository / Repozitorij:

[Repository of the University of Rijeka, Faculty of Medicine - FMRI Repository](#)





Conference paper
INVITED LECTURE

Molecular aspects of immunological evasion by cytomegalovirus*

IRENA CRNKOVIĆ and ULRICH H. KOSZINOWSKI

Received May 21, 1996.

Immune surveillance of CMV infection

Cytomegaloviruses (CMVs) are species-specific herpesviruses that constitute the Betaherpesvirus subfamily. Like all herpesviruses, they establish life-long infection in their hosts despite the presence of an active immune system. Primary infection with human CMV (HCMV) in the immunocompetent host is usually asymptomatic and leads to the establishment of a latent state from which reactivation can occur. However, infection of the immunologically immature or immunodeficient host such as organ transplant recipients or AIDS patients, can cause severe and often fatal disease (21). Analysis of the pathogenesis of the HCMV infection is hampered by the species specificity of cytomegaloviruses. Therefore, infection of the mouse with murine CMV (MCMV) is widely used as a model for HCMV infection and biology (22).

CMV gene expression is regulated in a cascade fashion. Viral proteins expressed in the immediate-early (IE) phase of infection are required for the subsequent transcription of early (E) genes (26). E proteins regulate the viral DNA replication, which is followed by the synthesis of the structural proteins during the late phase of infection.

Although different natural immunity mechanisms, especially NK cells, are associated with host resistance to CMV infection (35), a specific immune response is necessary for the clearance of CMV. During infection specific antiviral antibodies are produced and specific T lymphocytes are activated. Antibodies are not essential for the control of the acute viral infection, but they prevent viral spread after reactivation of CMV from latency (25). The presence of T lymphocytes is required for the clearance of the acute infection. CD8⁺ T cells play a decisive role in the antiviral defence (32, 33). They recognise peptides derived from viral proteins synthesised in in-

fecting cells and presented at the cell surface by specialised molecules encoded in the major histocompatibility complex (MHC) (38). Despite their dominant role in the control of CMV infection, CD8⁺ T lymphocytes are not indispensable. In their absence the virus is cleared by CD4⁺ -subset-dependent mechanisms (24). The antiviral activity of CD4⁺ cells can be mediated either by the direct, MHC II restricted, cytotoxic potential of these cells or by the production of cytokines like IFN-gamma and TNF-alpha, that have an antiviral effect (28, 29).

To establish a persistent infection in an immunocompetent host viruses require sophisticated mechanisms to struggle or to escape from the host immune control. CMV commands several mechanisms that enable the evasion from the host immune response. There is strong evidence that HCMV and MCMV possess several genes whose products interfere with immune effector mechanisms (Figure 1). More than one strategy is used. Viral proteins are expressed that bind to the Fc fragment of host immunoglobulins (Ig), others interfere with the antigen processing and presentation process in the MHC class I pathway. Additionally, HCMV encodes a homologue of the MHC class I heavy chain (13) and functional beta chemokine receptor (17). Both proteins are thought to intervene with natural immunity mechanisms.

The potential role of the CMV Fc receptor in immune evasion

The CMV protein with IgG Fc binding properties is certainly one of the viral products that has an obvious potential to interfere with the humoral immunity. The binding of human IgG to HCMV infected cells has been described long ago (27) but the gene responsible for this phenomenon is still not localised. The MCMV gene coding for the Fc receptor (FcR)

* Presented at the First Annual Meeting of the Croatian Immunological Society, Zagreb, November 23-24, 1995.

and the degradation of the newly synthesised molecules is enhanced (39, 42). The availability of HCMV deletion mutants revealed that a terminal genomic region contained more than one gene which independently down-regulate MHC class I heavy chain expression (23). The HCMV US11 gene product down-regulates class I molecules expression by dislocating MHC heavy chains from the endoplasmic reticulum (ER) to the cytosol, where they are rapidly destroyed by the cytosolic enzymes (41). A stable MHC complex is not formed. The second locus containing genes that affect MHC class I molecules is composed of the US2-US5 ORFs. The mechanism by which this gene(s) down-regulate MHC class I expression is not known.

Further, HCMV inhibits the TAP-dependent-transport of peptides from the cytosol into the ER. In the HCMV mutant ts9, in which the region between US1-US14 is deleted, TAP functions appear normal (19). Therefore, the gene coding for this function is probably located in this genomic region. Remarkably, the inactivation of TAP was first described as a mechanism used by the HSV-1 ICP47 protein (15, 20). By blocking the transport of cytosolic peptides into the lumen of the ER, ICP47 prevents the generation of stable MHC class I molecules and thus their ER export and expression at the surface of the infected cell. Given the essential function of TAP in translocation of antigenic peptides in the ER, it is not unexpected that it is a target for more than one member of the herpesvirus family. The unknown HCMV gene affecting TAP certainly differs from the HSV-1 ICP47 because no ICP47 homolog has been detected in the sequence of the HCMV genome.

MCMV interacts with the class I antigen presentation pathway by a different mechanism. The dominant MCMV antigen, recognised by the majority of antiviral cytotoxic T lymphocytes (CTL), is a nonapeptide derived from the IE protein pp89 and presented by L^d MHC class I molecules. It was observed that L^d-restricted pp89 specific CTL clone recognises pp89 during the IE phase of infection, but not during the early phase (7, 31). Lack of CTL recognition after expression of the E genes suggested the presence of a viral gene(s) that interact with antigen presentation. The rate of synthesis, stability and nuclear transport of pp89 was found no to be altered during the E phase. Neither were the formation and the stability of the trimolecular complex of MHC class I heavy chain, β_2 -microglobulin and peptide affected. Studies on the glycosylation pattern of MHC class I molecules revealed that they remained in an Endo H-sensitive form. This observation suggested that the complex is not degraded as in HCMV infected cells, but is retained in the endoplasmic reticulum (ER)/cis-Golgi compartment (9) (Figure 5).

The first step toward the identification of the genes whose products are responsible for retention of MHC class I molecules, was the construction of a recombinant MCMV deletion mutant that differs from wild type virus in its effect on antigen presentation. In the design of the deletion mutant we were directed by the sequence analysis of the MCMV genome. The




FIGURE 5. The MCMV early gene product retains MHC class I molecules in the ER/cis-Golgi compartment. During the immediate early phase class I molecules are transported to the cell surface. In the early phase class I molecules are retained in the ER/cis-Golgi cellular compartment. The putative viral gene product responsible for this effect is symbolised by the closed oval. In the Δ MS94.5 recombinant virus this gene is deleted, and transport of the MHC molecules to the cell surface is restored.

comparison between herpesvirus genomes has revealed conserved gene blocks shared between alpha-, beta- and gammaherpesviruses. The conserved gene blocks are located in the central portion of the genome and probably encode essential functions. The genes in the termini of the genomes are not conserved between different herpesviruses and are probably not essential for replication *in vitro*. It was reasoned that gene(s) involved in retention of a MHC class I molecules are members of one of the gene families located in the termini of the viral genome, similar to those encoding the HCMV US families. Therefore, the MCMV deletion mutant Δ MS94.5 was constructed with a large deletion (15,8 kb) in the genomic area represented by the *Hind*III E fragment of the MCMV genome (Figure 3) (37). The replication kinetics of the recombinant virus under *in vitro* conditions was indistinguishable from that of wild type virus. The next step was to verify if the viral genes that cause the block of MHC class I molecules transport are deleted in mutant Δ MS94.5. Therefore we compared the glycosylation pattern of MHC class I molecules in cells infected with wild type and recombinant virus. After infection with wild type virus and expression of E genes, MHC molecules remain Endo H-sensitive, typical for ER/cis-Golgi glycoproteins. In contrast, after infection with the Δ MS94.5 deletion mutant under the same conditions, the glycosylation pattern of newly synthesised MHC class I molecules was identical to that of noninfected cells (Figure 5).

Also, the presentation of pp89 to specific CTLs in the E phase was restored in the Δ MS94.5 infected cells. Using conditions permissive for E-gene ex-

pression, in contrast to wild type MCMV, pp89 presentation in the E phase and lysis by specific CD8⁺ T cells was not abolished after infection with Δ MS94.5 (Figure 6).




FIGURE 6. Restored antigen presentation during the E phase in MS94.5-infected cells. Mouse embryonal fibroblast target cells were infected with wild type MCMV or with the MS94.5 recombinant virus. Antigen presentation of pp89 was tested by pp89-specific CTLs in a standard 4 h cytolytic assay under the infection conditions with selective expression of IE or IE and E viral genes.

The results of the cytolytic assay as well as the glycosylation pattern of MHC class I molecules in cells infected with the Δ MS94.5 suggested that viral gene(s) responsible for retention were indeed deleted in recombinant Δ MS94.5. The next step in our experiments is the detailed analysis of the responsible gene that we have already defined by the gene microinjection technique. This will allow us to study the molecular mechanism of the MHC class I molecules retention.

There is evidence for more than one early-gene function of MCMV, which affect the MHC class I pathway of antigen presentation. Namely, within 3-6 hours after MCMV infection surface class I molecules disappear from the membrane of the infected cells. The export block of nascent MHC I complexes alone cannot explain this rapid reduction because the half life of resident class I molecules is much longer. It seems that, in addition to the retention of nascent class I molecules in ER/cis-Golgi, plasma membrane resident molecules are actively removed from the surface of the infected cells. The gene(s) that carry out this function are still unknown. They must be located outside the 15,8 kb gene region that is deleted in Δ MS94.5 and expressed in the E phase later than genes from this gene region because surface reduction of surface MHC complexes is present also in Δ MS94.5 mutant infected cells at later stages of infection. Viral E proteins that reside at the plasma membrane and coprecipitate with MHC class I molecules have been identified as potential candidates for this function. Whether they are really involved in the

MCMV effect on the class I molecules still has to be proven. Moreover, a reduction of the class I heavy chain synthesis has been proposed as additional mechanism of MCMV to affect antigen presentation (4).

The presence of multiple genes that can interact with the MHC class I antigen presentation pathway is so far unique for CMV. They may be descendants of a unique ancestor gene, and may have resulted from duplication and divergence during the evolution process. This hypothesis is supported by the presence of the tandemly arranged glycoprotein gene families in the CMV genome (5). However, a common origin of HCMV and MCMV MHC class I interacting genes is not likely, because beyond the phenotype of MHC class I reduction there is no significant homology between the genes and the products probably act by quite different mechanism.

The interference of MCMV with the transport of newly assembled MHC complexes combined with the removal of preexisting surface MHC class I proteins, represents a very effective evasion mechanism from CD8⁺ T lymphocyte control. If we consider the role CD8⁺ T cells still play in the control of the CMV infection, it is not surprising that MCMV has developed more than one function to evade CTL mediated control in order to persist in its host and be effectively transmitted despite the presence of an active immune system.

At the same time the still important role of CD8⁺ T cells indicates that the viral mechanism of immune evasion do operate *in vivo* only with a limited efficacy. The host immune system has found ways to recognise and to eliminate infected cells via CD8-dependent mechanisms, despite viral functions that downregulate antigen presentation by MHC class I molecules. One of the mechanisms involved is represented by IFN-gamma. IFN-gamma upregulates expression of the MHC class I molecules and the excess of the MHC molecules can escape the transport block and present antigen at the plasma membrane (18). CD8⁺ T cells also release IFN-gamma and restore antigen presentation in a paracrine fashion. This observation reconciles findings of altered *in vitro* presentation of viral antigens by class I molecules in parallel with CD8⁺ T cell mediated, MHC class I restricted protection *in vivo*. To what extent different viral mechanisms really contribute to CMV immunological evasion, can only be answered after *in vivo* studies of virus mutants that lack each of the MHC-reactive genes alone or in combination. This is a task that is possible to achieve only with MCMV recombinants.

Protection of the NK cells mediated lysis - a role for the viral class I homolog?

Although the reduction in MHC class I surface expression clearly results in a reduced T cell recognition of infected cells and probably has a positive effect for the multiplication and survival of CMV, it may also have some drawbacks for the virus. Namely, re-

duced MHC class I expression is balanced by enhanced NK sensitivity. It is known that NK cells are inhibited by the recognition of peptide filled class I molecules (6). This may explain the significant role of NK cells in MCMV clearance (40). The potential disadvantage of increased NK sensitivity due to the loss of class I molecules may be compensated by other viral mechanisms that counteract the NK-function. HCMV gene UL18 codes for a MHC class I homolog that binds cellular β_2 -m and can present endogenous peptides (13). It has been assumed that the peptide filled viral class I homolog which is expressed at the cell surface might engage NK cells inhibitory receptors and thus prevent the NK-mediated lysis of the infected cell. The UL18 gene product may represent the principle by which the virus balances the negative effects of MHC class I molecules down-regulation.

The HCMV chemokine receptor homolog an additional mechanism of immune evasion?

Chemokines are small proinflammatory peptides with leukocyte-chemoattractant activity. The chemokine receptors are members of the G protein-coupled receptor superfamily. A data base search has revealed that the HCMV US28 gene product is 33% identical to the mammalian leukocyte receptors for alpha and beta chemokines (16). There is also functional homology. The binding of the chemokine to the US28 gene product leads to the intracellular calcium mobilisation, but, distinct from that of known mammalian receptors, the binding is selective for beta chemokines (17). It can be expected that expression of this protein may interfere with the host antiviral response but the definitive active principle is still enigmatic.

Concluding remarks

CMV is another example of a virus which has evolved multiple mechanisms to evade hosts immune control. Remarkably the strategies appear different from those employed by another DNA virus, Pox virus (34). Although the viral interference with expression of MHC class I molecules is the best documented, CMV also modulates the humoral immune response as well as mechanisms of natural immunity. The CMV genome contains numerous genes whose functions are probably not essential for viral growth in fibroblasts *in vitro*. These genes demonstrate their effect only *in vivo*. Therefore we anticipate that all deletion mutants will show some degree of attenuation, either generally or only in some tissues of the host. The characterisation of the attenuation mechanism may cause formidable difficulties. Deletion of the genes coding for immune evasion mechanism as well as the genes coding for functions that participate in the infection of certain target cells, virus spread, regulation of gene expression or establishment of latency, will result in a similar phenotype: limited virus growth after infection.

These genes remain to be discovered. The dissection and analysis of their functions will provide us informations on new principles in cell biology and moreover, they hold the promise for new therapeutic approaches.

Acknowledgements: Ulrich H. Koszinowski is supported by grants of the Deutsche Forschungsgemeinschaft. We thank Martin Messerle for the critical reading of the manuscript.

REFERENCES

1. BEERSMA M F C, BIJLMAKERS M J E, PLOEGH H L 1993 Human cytomegalovirus down-regulates HLA class I expression by reducing the stability of class I heavy chains. *J Immunol* 151: 4455-4464
2. BROWNE H, SMITH G, BECK S, MINSON T 1990 A complex between the MHC class I homologue encoded by human cytomegalovirus and β_2 microglobulin. *Nature (London)* 347: 770-772
3. BURGERT H G, KVIST S 1985 An adenovirus type 2 glycoprotein blocks cell surface expression of human histocompatibility class I antigens. *Cell* 41: 987-997
4. CAMPBELL A E, SLATER J S 1994 Down-regulation of major histocompatibility complex class I synthesis by murine cytomegalovirus early gene expression. *J Virol* 68: 1805-1811
5. CHEE M S, BANKIER A T, CECK S, BJOHNI R, BROWN C M, CERNY R, HORSNELL T, HUTCHISON C A, KOUZARIDES T, MARTIGNETTI T A, PREDDIE E, SATCHWELL S C, TOMLINSON P, WESTON K M, BARELL B G 1990 Analysis of the protein-coding content of the sequence of the human cytomegalovirus strain AD 169. *Curr Top Microbiol Immunol* 154: 125-169
6. CORREA I, RAULET D H 1995 Binding of the diverse peptides to MHC class I molecules inhibits target cell lysis by activated natural killer cells. *Immunity* 2: 61-71
7. DEL VAL M, VOLKMER H, ROTHBARD J B, JONJIC S, MESSERLE M, SCHICKEDANZ J, REDDEHASE M J, KOSZINOWSKI U H 1988 Molecular basis for cytolytic T lymphocyte recognition of murine cytomegalovirus immediate-early protein pp89. *J Virol* 62: 3965-3972
8. DEL VAL M, MÜNCH K, REDDEHASE M J, KOSZINOWSKI U H 1989 Presentation of CMV immediate-early antigen to cytolytic T lymphocytes is selectively prevented by viral genes expressed in the early phase. *Cell* 56: 305-315
9. DEL VAL M, HENGEL H, HÄCKER H, HARTLAUB U, RUPPERT T, LUČIN P, KOSZINOWSKI U H 1992 Cytomegalovirus prevents antigen presentation by blocking the transport of peptide-loaded major histocompatibility complex class I molecules into the medial-transport of peptide-loaded major histocompatibility complex class I molecules into the medial Golgi compartment. *J Exp Med* 176: 729-738
10. DINGWELL K S, BRUNETTI C R, HENDRICKS R L, TANG Q, TANG M, RAINBOW A J, JOHNSON D C 1994 Herpes simplex virus glycoproteins E and I facilitate cell-to-cell spread *in vivo* and across junctions of cultured cells. *J Virol* 68: 834-845
11. DINGWELL K S, DOERING L C, JOHNSON D C 1995 Glycoproteins E and I facilitate neuron-to-neuron spread of herpes simplex virus. *J Virol* 69: 7087-7098
12. DUBIN G, SOCOLOF E, FRANK I, FRIEDMAN H M 1991 Herpes simplex virus type I Fc receptor protect infected cells from antibody-dependent cellular cytotoxicity. *J Virol* 65: 7046-7050

13. FAHNESTOCK M, JOHNSON J L, FELDMAN R M R, NEVEU J M, LANE W S, BJORKMAN P J 1995 The MHC class I homolog encoded by human cytomegalovirus binds endogenous peptides. *Immunity* 3: 583-590
14. FRANK I, FRIEDMAN H M 1989 A novel function of the herpes simplex virus type I Fc receptor: participation in bipolar bridging of antiviral immunoglobulin G. *J Virol* 63: 4479-4488
15. FRÜH K, AHN K, DJABALLAH H, SEMPE P, VAN ENDERT P M, TAMPE R, PETERSON R A, YANG Y 1995 A viral inhibitor of peptide transporters for antigen presentation. *Nature (London)* 375: 415-418
16. GAO J L, KUHNS D B, TIFFANY H L, McDERMOTT D, LI X, FRANCKE U, MURPHY P M 1993 Structure and functional expression of the human macrophage inflammatory protein 1 alpha/RANTES receptor. *J Exp Med* 177: 1421-1427
17. GAO J L, MURPHY P M 1994 Human cytomegalovirus open reading frame US28 encodes a functional beta chemokine receptor. *J Biol Chem* 269: 28539-28542
18. HENGEL H, LUČIN P, JONJIC S, RUPPERT T, KOSZINOWSKI U H 1994 Restoration of cytomegalovirus antigen presentation by gamma interferon combats viral escape. *J Virol* 68: 289-297
19. HENGEL H, FLOHR T, HÄMMERLING G J, KOSZINOWSKI U H, MOMBURG F 1996 Human cytomegalovirus inhibits peptide translocation into the endoplasmic reticulum for MHC class I assembly. *J Gen Virol (in press)*
20. HILL A, JUGOVIĆ P, YORK I, RUSS G, BENNIK J, YEWDELL J, PLOEGH H, JOHNSON D 1995 Herpes simplex virus turns off the TAP to evade host immunity. *Nature (London)* 375: 411-415
21. HO M 1994 Advances in understanding cytomegalovirus infection after transplantation. *Transplant Proc* 26: 7-11
22. HUDSON J B 1979 The murine cytomegalovirus as a model for the study of viral pathogenesis and persistent infection. *Arch Virol* 62: 1-29
23. JONES T R, HANSON L K, SUN L, SLATER J S, CAMPBELL A E 1995 Multiple independent loci within the human cytomegalovirus unique short region down-regulate expression of major histocompatibility complex class I heavy chains. *J Virol* 69: 4830-4841
24. JONJIC S, PAVIĆ I, LUČIN P, RUKAVINA D, KOSZINOWSKI U H 1990 Efficacious control of cytomegalovirus infection after long-term depletion of CD8⁺ T lymphocytes. *J Virol* 64: 5457-5464
25. JONJIC S, PAVIĆ I, POLIĆ B, CRNKOVIĆ I, LUČIN P, KOSZINOWSKI U H 1994 Antibodies are not essential for the resolution of primary cytomegalovirus infection but limit dissemination of recurrent virus. *J Exp Med* 179: 1713-1717
26. KEIL G M, EBELING-KEIL A, KOSZINOWSKI U H 1984 Temporal regulation of the murine cytomegalovirus transcription and mapping of viral RNA synthesised at immediate early times after infection. *J Virol* 50: 784-795
27. KELLER R, PEITCHEL R, GOLDMAN J N, GOLDMAN M 1976 An IgG-Fc receptor induced in cytomegalovirus infected human fibroblasts. *J Immunol* 116: 772-777
28. LUČIN P, PAVIĆ I, POLIĆ B, JONJIC S, KOSZINOWSKI U H 1992 Gamma interferon-dependent clearance of cytomegalovirus infection salivary gland. *J Virol* 66: 1977-1984

29. PAVIĆ I, POLIĆ B, CRNKOVIĆ I, LUČIN P, JONJIC S, KOSZINOWSKI U H 1993 Participation of endogenous tumour necrosis factor alpha in host resistance to cytomegalovirus infection. *J Gen Virol* 74: 2215-2223
30. PIERSCHABCHER M D, RUOSLAHTI E 1984 Cell attachment activity of fibronectin can be duplicated by small synthetic fragments of the molecule. *Nature (London)* 309: 30-33
31. REDEHASE M, FIBI M R, KEIL G M, KOSZINOWSKI U H 1986 Late-phase expression of a murine cytomegalovirus immediate-early antigen recognised by cytolytic T lymphocytes. *J Virol* 60: 1125-11129
32. REDDEHASE M J, MUTTER W, MUNCH , BUHRING H J, KOSZINOWSKI U H 1987 CD8 positive T lymphocytes specific for murine cytomegalovirus immediate-early genes mediate protective immunity. *J Virol* 61: 3102-3108
33. RIDDEL S, WATANABE K, GOODRICH J, LI C, AGHA C R, GREENBERG P 1992 Restoration of viral immunity in immunodeficient humans by the adoptive transfer of T cell clones. *Science* 257: 238-241
34. SMITH G L 1993 Vaccinia virus glycoproteins and immune evasion. *J Gen Virol* 74: 1725-1740
35. SHELLAM G R, ALLEN J E, PAPADIMITROU J M, BANCROFT G J 1981 Increased susceptibility to cytomegalovirus infection in beige mutant mice. *Proc Natl Acad Sci USA* 78: 5104-5113
36. THÅLE R, LUČIN P, SCHEIDER K, EGGERS M, KOSZINOWSKI U H 1994 Identification and expression of a murine cytomegalovirus early gene coding for and Fc receptor. *J Virol* 68: 7757-7765
37. THÅLE R, SZEPAŃ U, HENGEL H, GEGINAT G, LUČIN P, KOSZINOWSKI U H 1995 Identification of the mouse cytomegalovirus genomic region affecting major histocompatibility complex class I transport. *J Virol* 68: 6098-6105
38. TOWNSEND A, BODMER H 1989 Antigen recognition by class I-restricted T lymphocytes. *Annu Rev Immunol* 7: 601-624
39. WARREN A P, DUCROQ D H, LEHNER P J, BORYSIEWICZ L K 1994 Human cytomegalovirus-infected cells have unstable assembly of major histocompatibility complex class I complexes and are resistant to lysis by cytotoxic T lymphocytes. *J Virol* 68: 2822-2829
40. WELSH R M, BRUBAKER J O, VARGAS-CORTES M, O'DONELL C L 1991 Natural killer (NK) cell response to virus infections in mice with severe combined immunodeficiency. The stimulation of NK cells and NK cell dependent control of virus infection occur independently of T and B function. *J Exp Med* 173: 1053-1063
41. WIERTZ J H J, JONES T R, SUN L, BOGYO M, GEUZE H J, PLOEGH H L 1996 The human cytomegalovirus US11 gene product dislocates MHC class I heavy chains from the endoplasmic reticulum to the cytosol. *Cell* 84: 769-779
42. YAMASHITA Y, SHIMOKATA K, SAGA S, MIZUNO S, TSURMI T, NISHIYAMA Y 1994 Rapid degradation of the heavy chain of class I major histocompatibility complex antigens in the endoplasmic reticulum of human cytomegalovirus-infected cells. *J Virol* 68: 7933-7943
43. ZSAK L, ZUCKERMANN F, SUGG N, BEN-PORAT T 1992 Glycoprotein gI of pseudorabies virus promotes cell fusion and virus spread via direct cell-to-cell transmission. *J Virol* 66: 2316-2325

ABSTRACT***Molecular aspects of immunological evasion by cytomegalovirus***

Several viruses have developed strategies to modulate or evade the immune control of their hosts. Different ways of interference with distinct host mechanisms which contribute to the control of the viral infection are used: from the interaction with complement system and cytokines, to the inhibition of the specific humoral and cellular immune response. Here, we describe principles of immune evasion used by cytomegaloviruses. They include gene products that bind host immunoglobulins, interfere with antigen presentation in the MHC class I pathway, abrogate the recognition by NK cells or serve as receptors for chemokines.

¹Department of Virology, Institute for Hygiene, University of Heidelberg, 69120 Heidelberg, Germany

²Max von Pettenkofer-Institute for Hygiene and Medical Microbiology, Pettenkoferstraße 9a, 80336 München, Germany

Correspondence:
Ulrich H. Koszinowski
Max von Pettenkofer-Institute for Hygiene and Medical Microbiology
Pettenkoferstraße 9a
80336 München, Germany