

ENTERALNA PREHRANA NEKAD I SAD

Pešić, Mladen; Krznarić, Željko

Source / Izvornik: **Medicina, 2003, 42(39), 224 - 228**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:184:525944>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-07-08**



Repository / Repozitorij:

[Repository of the University of Rijeka, Faculty of
Medicine - FMRI Repository](#)





TEMATSKI BROJ

Enteralna prehrana

Gosti urednici: Mladen Peršić i Željko Krznarić

ENTERALNA PREHRANA NEKAD I SAD

Pravilna prehrana temeljni je uvjet opstanka živog bića, tj. preduvjet održanja energetskih i metaboličnih homeostatskih procesa. U suvremenoj ljudskoj zajednici uzimanje hrane nije samo ispunjavanje fiziološke potrebe nego postaje obilježjem kulture i statusa, zrcalom životne filozofije i religijskih nazora. Istodobno, značajan je rast interesa i brige za održanje zdravlja i potrebe da se dijetnim modifikacijama produlji životni vijek, spriječe ili izliječe kronične bolesti, osigura dugovječnost i zdrava starost.

Spoznaje o značenju prehrane u liječenju bolesnika prisutne su odavno i sastavni su dio svih medicinskih kultura i škola, uključivši našu sadašnju.

Pojam "klinička prehrana" obuhvaća sve oblike prehrane bolesnika, tj. uobičajenu peroralnu prehranu, dijetne modifikacije i pripravke te enteralnu i parenteralnu prehranu.

Osnovni koncept nutritivne potpore bolesnicima datira još iz vremena ranih civilizacija Egipta, Grčke i Rima, kada se postavljaju i temelji enteralne prehrane.

Enteralna prehrana važan je segment kliničke prehrane, a osnovni je preduvjet njezine primjene strukturalno i funkcionalno podoban probavni sustav, osobito tanko i debelo crijevo. U ovom obliku nutritivne podrške hrana i/ili komercijalne nutritivne otopine i pripravci, unose se različitim hranidbenim sondama u želudac, dvanaesnik ili jejunum.

Nutritivna potpora enteralnim putem u zadnja dva desetljeća doživljava upravo neviđen uzlet. Značajan napredak u razvoju posebnih enteralnih formula te različitih akcesorija (sonde, pumpe) doveo je do toga da je enteralna prehrana postala najprihvatljiviji i ekonomski najpovoljniji način nutritivne potpore u bolničkim i kućnim uvjetima liječenja.

Različiti oblici enteralnih pripravaka razvijaju se unatrag više stotina, ako ne i tisuća godina. Medicinski zapi-

si iz razdoblja Starog Egipta, 3000 godina prije Krista, opisuju primjenu nutritivnih klizmi (mješavina vina, mlijeka, surutke, pšenice, bujona) kao alternativnog načina hranjenja bolesnika i liječenja proljeva.

Zanimljivost je da modificirani koncept nutritivnih klizmi primjenom krantolančanih masnih kiselina (butiratne ili maslačne) u liječenju ulceroznog kolitisa predstavlja danas gotovo vrhunac razvoja nutritivne potpore i održanja integriteta sluznice debelog crijeva. Capivaccus (1598.), Aquapendente (1617.), Von Helmont (1646.) i Boerhave (1710.) svojim idejama otvaraju put razvoju nazogastričnog načina hranjenja primjenom primitivnih oblika sondi. John Hunter, rodonačelnik britanske kirurgije, opisuje 1790. godine, na način primjeren suvremenome medicinskom razmišljanju, primjenu nutritivne sonde za dopremanje hrane u želudac. Ta je sonda bila načinjena od jeguljine kože i kitove kosti. Sljedeći značajni koraci u razvoju enteralne prehrane zbili su se 1910. godine, kada je Max Einhorn opisao prvu primjenu duodenalne sonde, te 1958. godine, kada Barron i Fallis razvijaju prvu enteralnu pumpu i načine kontinuirane primjene enteralnih pripravaka.

Enteralne formule razvijaju se neprekidno od sredine dvadesetog stoljeća, a prekretnica je Rosov prototip elementarnog enteralnog pripravka koji se pojavio 1949. godine. Ruski astronaut German Titov (kolovoz, 1961.) i američki John Glenn (veljača, 1962.) prvi su ljudi koji su se hranili u svemiru, ali su i prvi ljudi koji su se suočili sa situacijama mikrogravitacije i hipogravitacije, te posljedica djelovanja na hranjenje. Osobitu važnost za razvoj enteralnih pripravaka ima NASA-in program svemirskih letova s ljudskom posadom i potreba za razvojem enteralnih pripravaka pogodnih za hranjenje astronauta, s naglaskom na smanjenju količine stolice. Iz tog je razdoblja tipičan prikaz Greensteina i Winitza koji

se ubraja u klasike nutricionističke literature, a objavljen u časopisu *Nature* 1965 godine.

Od tihog ulaska enteralnih formula u kliničku medicinu prije pedeset godina, do danas je učinjen značajan napredak. Sredinom prošlog stoljeća imali smo ograničene spoznaje o značenju hranjenja bolesnika koji nisu mogli uzimati normalnu hranu, o komplikacijama enteralne, ali i parenteralne nutritivne potpore, a razvoj enteralnih otopina bio je tek u začetku.

Klinička prehrana danas, osobito enteralna prehrana, spada u sam vrh medicinske misli i djelovanja. Rastu spoznaje o interakciji hrane i gena, a različiti nutritivni sastojci svojim farmakološkim djelovanjima afirmiraju imunonutriciju kao sasvim nov koncept u razvoju ovog segmenta liječenja.

Na europskom tržištu trenutačno je prisutno više od 150 različitih enteralnih pripravaka, a na listi Hrvatskog zavoda za zdravstveno osiguranje nalazi se 19 enteralnih pripravaka. To omogućuje različite oblike nutritivne podrške bolesnika ne samo u institucijama, već i u kućnom liječenju, što je iskorak gotovo istovjetan razvoju najsofisticiranijih formula ili sustava za aplikaciju.

Enteralna prehrana, kao temelj, te eventualna njezina nadopuna parenteralnim pripravcima do zadovoljenja potpunih nutritivnih potreba, vrh su nutritivne piramide.

Mladen Peršić

KBC Rijeka
Klinika za pedijatriju
"Kantrida"

Željko Krznarić

KBC Zagreb-Rebro
Interna Klinika
Zavod za gastroenterologiju

LITERATURA

- Harkness L. The history of enteral nutrition therapy: from raw eggs and nasal tubes to purified amino acids and early postoperative jejunal delivery. *J Am Diet Assoc* 2002; 102(3):399-404.
- DiSario JA, Baskin WN, Brown RD, DeLegge MH, Fang JC, Ginsberg GG et al. Endoscopic approaches to enteral nutritional support. *Gastrointest Endosc* 2002; 55(7):901-908.
- Kudsk KA. Current aspects of mucosal immunology and its influence by nutrition. *Am J Surg* 2002; 183(4):390-398.
- McClave SA, Marsano LS, Lukan JK. Enteral access for nutritional support: rationale for utilization. *J Clin Gastroenterol* 2002; 35(3):209-213.
- Jeejeebhoy KN. Enteral and parenteral nutrition: evidence-based approach. *Proc Nutr Soc* 2001; 60(3):399-402.
- American Gastroenterological Association Medical Position Statement: guidelines for the use of enteral nutrition. *Gastroenterology* 1995; 108(4):1280-1281.
- Klein S, Kinney J, Jeejeebhoy K, Alpers D, Hellerstein M, Murray M et al. Nutrition support in clinical practice: review of published data and recommendations for future research directions. National Institutes of Health, American Society for Parenteral and Enteral Nutrition, and American Society for Clinical Nutrition. *JPEN J Parenter Enteral Nutr* 1997; 21(3):133-156.
- Pearce CB, Duncan HD. Enteral feeding. Nasogastric, nasojejunal, percutaneous endoscopic gastrostomy, or jejunostomy: its indications and limitations. *Postgrad Med J* 2002; 78(918):198-204.
- Gabriel SA, Ackermann RJ, Castresana MR. A new technique for placement of nasoenteral feeding tubes using external magnetic guidance. *Crit Care Med* 1997; 25(4):641-645.
- Boivin M, Levy H, Hayes J. A multicenter, prospective study of the placement of transpyloric feeding tubes with assistance of a magnetic device. The Magnet-Guided Enteral Feeding Tube Study Group. *JPEN J Parenter Enteral Nutr* 2000; 24(5):304-307.
- Ozdemir B, Frost M, Hayes J, Sullivan DH. Placement of nasoenteral feeding tubes using magnetic guidance: retesting a new technique. *J Am Coll Nutr* 2000; 19(4):446-451.
- Kearns PJ, Donna C. A controlled comparison of traditional feeding tube verification methods to a bedside, electromagnetic technique. *JPEN J Parenter Enteral Nutr* 2001; 25(4):210-215.
- Grathwohl KW, Gibbons RV, Dillard TA, Horwhat JD, Roth BJ, Thompson JW et al. Bedside videoscopic placement of feeding tubes: development of fiberoptics through the tube. *Crit Care Med* 1997; 25(4):629-634.
- Gauderer MW. Percutaneous endoscopic gastrostomy and the evolution of contemporary long-term enteral access. *Clin Nutr* 2002; 21(2):103-110.
- Adler DG, Gostout CJ, Baron TH. Percutaneous transgastric placement of jejunal feeding tubes with an ultrathin endoscope. *Gastrointest Endosc* 2002; 55(1):106-110.
- Whelan K, Gibson GR, Judd PA, Taylor MA. The role of probiotics and prebiotics in the management of diarrhoea associated with enteral tube feeding. *J Hum Nutr Diet* 2001; 14(6):423-433.
- Afzal NA, Addai S, Fagbemi A, Murch S, Thomson M, Heuschkel R. Refeeding syndrome with enteral nutrition in children: a case report, literature review and clinical guidelines. *Clin Nutr* 2002; 21(6):515-520.
- Guidelines for the use of parenteral and enteral nutrition in adult and pediatric patients. *JPEN J Parenter Enteral Nutr* 2002; 26(1 Suppl):1SA-138SA.
- Nobel JJ. Enteral feeding pumps. *Pediatr Emerg Care* 1996; 12(2):128-136.
- Varella LD, Young RJ. New options for pumps and tubes: progress in enteral feeding techniques and devices. *Curr Opin Clin Nutr Metab Care* 1999; 2(4):271-275.
- Gomez CC, Cos Blanco AI, Iglesias RC, Planas VM, Castella M, Garcia Luna PP et al. Home enteral nutrition. Annual report 1999. NADYA-SENPE Group. *Nutr Hosp* 2002; 17(1):28-33.
- Marik PE, Zaloga GP. Early enteral nutrition in acutely ill patients: a systematic review. *Crit Care Med* 2001; 29(12):2264-2270.
- Davies AR, Froome PR, French CJ, Bellomo R, Gutteridge GA, Nyulasi L et al. Randomized comparison of nasojejunal and nasogastric feeding in critically ill patients. *Crit Care Med* 2002; 30(3):586-590.
- Montejo JC, et al. Multicenter, prospective, randomized, single blind study comparing the efficacy and gastroin-

- testinal complications of early jejunal feeding with early gastric feeding in critically ill patients. *Crit Care Med* 2002; 30(4):796.
- Caparros T, Lopez J, Grau T. Early enteral nutrition in critically ill patients with a high-protein diet enriched with arginine, fiber, and antioxidants compared with a standard high-protein diet. The effect on nosocomial infections and outcome. *JPEN J Parenter Enteral Nutr* 2001; 25(6):299-308.
- Bines JE, Taylor RG, Justice F, Paris MC, Sourial M, Nagy E et al. Influence of diet complexity on intestinal adaptation following massive small bowel resection in a preclinical model. *J Gastroenterol Hepatol* 2002; 17(11):1170-1179.
- Bozzetti F, Braga M, Gianotti L, Gavazzi C, Mariani L. Post-operative enteral versus parenteral nutrition in malnourished patients with gastrointestinal cancer: a randomised multicentre trial. *Lancet* 2001; 358(9292):1487-1492.
- Lewis SJ, Egger M, Sylvester PA, Thomas S. Early enteral feeding versus "nil by mouth" after gastrointestinal surgery: systematic review and meta-analysis of controlled trials. *BMJ* 2001; 323:1-5.
- Braga M, Gianotti L, Gentilini O, Liotta S, Di C, V. Feeding the gut early after digestive surgery: results of a nine-year experience. *Clin Nutr* 2002; 21(1):59-65.
- Braga M, Gianotti L, Nespoli L, Radaelli G, Di C, V. Nutritional approach in malnourished surgical patients: a prospective randomized study. *Arch Surg* 2002; 137(2):174-180.
- Rayes N, Hansen S, Seehofer D, Muller AR, Serke S, Bengmark S et al. Early enteral supply of fiber and Lactobacilli versus conventional nutrition: a controlled trial in patients with major abdominal surgery. *Nutrition* 2002; 18(7-8):609-615.
- (Gassull MA, Cabre E. Nutrition in inflammatory bowel disease. *Curr Opin Clin Nutr Metab Care* 2001; 4(6):561-569.
- Walker-Smith JA. Management of growth failure in Crohn's disease. *Arch Dis Child* 1996; 75(4):351-354.
- O'Morain C, Segal AW, Levi AJ. Elemental diet as primary treatment of acute Crohn's disease: a controlled trial. *Br Med J (Clin Res Ed)* 1984; 288(6434):1859-1862.
- Sanderson IR, Udeen S, Davies PS, Savage MO, Walker-Smith JA. Remission induced by an elemental diet in small bowel Crohn's disease. *Arch Dis Child* 1987; 62(2):123-127.
- Malchow H, Steinhardt HJ, Lorenz-Meyer H, Strohm WD, Rasmussen S, Sommer H et al. Feasibility and effectiveness of a defined-formula diet regimen in treating active Crohn's disease. European Cooperative Crohn's Disease Study III. *Scand J Gastroenterol* 1990; 25(3):235-244.
- Lochs H, Steinhardt HJ, Klaus-Wentz B, Zeitz M, Vogelsang H, Sommer H et al. Comparison of enteral nutrition and drug treatment in active Crohn's disease. Results of the European Cooperative Crohn's Disease Study. IV. *Gastroenterology* 1991; 101(4):881-888.
- Akobeng AK. Enteral nutritional supplementation and the maintenance of remission in Crohn's disease. *J Pediatr Gastroenterol Nutr* 2002; 34(5):572-573.
- Fernandez-Banares F, Cabre E, Esteve-Comas M, Gassull MA. How effective is enteral nutrition in inducing clinical remission in active Crohn's disease? A meta-analysis of the randomized clinical trials. *JPEN J Parenter Enteral Nutr* 1995; 19(5):356-364.
- Griffiths AM, Ohlsson A, Sherman PM, Sutherland LR. Meta-analysis of enteral nutrition as a primary treatment of active Crohn's disease. *Gastroenterology* 1995; 108(4):1056-1067.
- Heuschkel RB, Menache CC, Megerian JT, Baird AE. Enteral nutrition and corticosteroids in the treatment of acute Crohn's disease in children. *J Pediatr Gastroenterol Nutr* 2000; 31(1):8-15.
- Zachos M, Tondeur M, Griffiths AM. Enteral nutritional therapy for inducing remission of Crohn's disease. *Cochrane Database Syst Rev* 2001;(3):CD000542.
- Verma S, Brown S, Kirkwood B, Gjaffer MH. Polymeric versus elemental diet as primary treatment in active Crohn's disease: a randomized, double-blind trial. *Am J Gastroenterol* 2000; 95(3):735-739.
- Cameron EA, Middleton SJ. Elemental diet in the treatment of orofacial Crohn's disease. *Gut* 2003; 52(1):143.
- Beattie RM, Schifffrin EJ, Donnet-Hughes A, Huggett AC, Domizio P, MacDonald TT et al. Polymeric nutrition as the primary therapy in children with small bowel Crohn's disease. *Aliment Pharmacol Ther* 1994; 8(6):609-615.
- Gassull MA, Fernandez-Banares F, Cabre E, Papo M, Gjaffer MH, Sanchez-Lombrana JL et al. Fat composition may be a clue to explain the primary therapeutic effect of enteral nutrition in Crohn's disease: results of a double blind randomised multicentre European trial. *Gut* 2002; 51(2):164-168.
- Fell JM, Paintin M, Arnaud-Battandier F, Beattie RM, Hollis A, Kitching P et al. Mucosal healing and a fall in mucosal pro-inflammatory cytokine mRNA induced by a specific oral polymeric diet in paediatric Crohn's disease. *Aliment Pharmacol Ther* 2000; 14(3):281-289.
- Johansson E, Lonroth I, Lange S, Jonson I, Jennische E, Lonroth C. Molecular cloning and expression of a pituitary gland protein modulating intestinal fluid secretion. *J Biol Chem* 1995; 270(35):20615-20620.
- Bjorck S, Bosaeus I, Ek E, Jennische E, Lonroth I, Johansson E et al. Food induced stimulation of the antisecretory factor can improve symptoms in human inflammatory bowel disease: a study of a concept. *Gut* 2000; 46(6):824-829.
- Corey M, McLaughlin FJ, Williams M, Levison H. A comparison of survival, growth, and pulmonary function in patients with cystic fibrosis in Boston and Toronto. *J Clin Epidemiol* 1988; 41(6):583-591.
- Akobeng AK, Miller V, Thomas A. Percutaneous endoscopic gastrostomy feeding improves nutritional status and stabilizes pulmonary function in patients with cystic fibrosis. *J Pediatr Gastroenterol Nutr* 1999; 29(4):485-486.
- Walker SA, Gozal D. Pulmonary function correlates in the prediction of long-term weight gain in cystic fibrosis patients with gastrostomy tube feedings. *J Pediatr Gastroenterol Nutr* 1998; 27(1):53-56.
- Williams SG, Ashworth F, McAlweenie A, Poole S, Hodson ME, Westaby D. Percutaneous endoscopic gastrostomy feeding in patients with cystic fibrosis. *Gut* 1999; 44(1):87-90.
- Erskine JM, Lingard CD, Sontag MK, Accurso FJ. Enteral nutrition for patients with cystic fibrosis: comparison of a semi-elemental and nonelemental formula. *J Pediatr* 1998; 132(2):265-269.
- Gottrand F, Hankard R, Michaud L, Ategbo S, Dabadie A, Druon D et al. Effect of glucose to fat ratio on energy sub-

- strate disposal in children with cystic fibrosis fed enterally. *Clin Nutr* 1999; 18(5):297-300.
- Conway SP, Morton A, Wolfe S. Enteral tube feeding for cystic fibrosis. *Cochrane Database Syst Rev* 2000;(2):CD001198.
- Borowitz D, Baker RD, Stallings V. Consensus report on nutrition for pediatric patients with cystic fibrosis. *J Pediatr Gastroenterol Nutr* 2002; 35(3):246-259.
- Sinaasappel M, Stern M, Littlewood J, Wolfe S, Steinkamp G, Heijerman H et al. Nutrition in patients with cystic fibrosis: a European Consensus. *J Cystic Fibrosis* 2002; 1(2):51-75.
- Lobo DN, Memon MA, Allison SP, Rowlands BJ. Evolution of nutritional support in acute pancreatitis. *Br J Surg* 2000; 87(6):695-707.
- Kalfarentzos F, Kehagias J, Mead N, Kokkinis K, Gogos CA. Enteral nutrition is superior to parenteral nutrition in severe acute pancreatitis: results of a randomized prospective trial. *Br J Surg* 1997; 84(12):1665-1669.
- Imrie CW, Carter CR, McKay CJ. Enteral and parenteral nutrition in acute pancreatitis. *Best Pract Res Clin Gastroenterol* 2002; 16(3):391-397.
- Eckerwall G, Andersson R. Early enteral nutrition in severe acute pancreatitis: a way of providing nutrients, gut barrier protection, immunomodulation, or all of them? *Scand J Gastroenterol* 2001; 36(5):449-458.
- Windsor AC, Kanwar S, Li AG, Barnes E, Guthrie JA, Spark JI et al. Compared with parenteral nutrition, enteral feeding attenuates the acute phase response and improves disease severity in acute pancreatitis. *Gut* 1998; 42(3):431-435.
- Al Omran M, Groof A, Wilke D. Enteral versus parenteral nutrition for acute pancreatitis. *Cochrane Database Syst Rev* 2001;(2):CD002837.
- Olah A, Pardavi G, Belagyi T, Nagy A, Issekutz A, Mohamed GE. Early nasojejunal feeding in acute pancreatitis is associated with a lower complication rate. *Nutrition* 2002; 18(3):259-262.
- Olah A, Belagyi T, Issekutz A, Gamal ME, Bengmark S. Randomized clinical trial of specific lactobacillus and fibre supplement to early enteral nutrition in patients with acute pancreatitis. *Br J Surg* 2002; 89(9):1103-1107.
- Abou-Assi S, Craig K, O'Keefe SJ. Hypocaloric jejunal feeding is better than total parenteral nutrition in acute pancreatitis: results of a randomized comparative study. *Am J Gastroenterol* 2002; 97(9):2255-2262.
- McClave SA, Dryden GW. Issues of nutritional support for the patient with acute pancreatitis. *Semin Gastrointest Dis* 2002; 13(3):154-160.
- Meier R, Beglinger C, Layer P, Gullo L, Keim V, Laugier R et al. ESPEN guidelines on nutrition in acute pancreatitis. European Society of Parenteral and Enteral Nutrition. *Clin Nutr* 2002; 21(2):173-183.
- Sentongo T, Mascarenhas MR. Newer components of enteral formulas. *Pediatr Clin North Am* 2002; 49(1):113-125.
- Boelens PG, Houdijk AP, De Thouars HN, Teerlink T, Van Engeland MI, Haarman HJ et al. Plasma taurine concentrations increase after enteral glutamine supplementation in trauma patients and stressed rats. *Am J Clin Nutr* 2003; 77(1):250-256.
- Buchman AL. Glutamine for the gut: mystical properties or an ordinary amino acid? *Curr Gastroenterol Rep* 1999; 1(5):417-423.
- Buchman AL. Glutamine: commercially essential or conditionally essential? A critical appraisal of the human data. *Am J Clin Nutr* 2001; 74(1):25-32.

USPOREDBA ENTERALNE I PARENTERALNE PREHRANE

COMPARISON BETWEEN ENTERAL AND PARENTERAL NUTRITION

Željko Krznarić¹, Živka Dika²

SAŽETAK

Enteralna i parenteralna prehrana osnovni su oblici nutritivne podrške u kliničkoj medicini. Ciljevi primjene enteralne i parenteralne prehrane jesu prevencija ili korekcija nutritivnih deficita i sindroma malnutricije. Dobro poznavanje tih oblika kliničke prehrane omogućava valjan i pravodoban terapijski pristup. Enteralna prehrana ima brojne prednosti naspram parenteralne prehrane u bolesnika s očuvanom strukturom i funkcijom probavnog sustava. Parenteralna prehrana je način supstitucije organizma vodom, energetske supstratima, kao i svim drugim hranjivim tvarima krvožilnim putem u bolesnika s kontraindikacijama za primjenu enteralne prehrane ili u situacijama kada se primjenom enteralne prehrane ne mogu u potpunosti zadovoljiti sve nutritivne potrebe.

KLJUČNE RIJEČI: enteralna prehrana, parenteralna prehrana, indikacije, provođenje, komplikacije

Klinička prehrana obuhvaća sve oblike prehrane bolesnika, tj. uobičajenu peroralnu prehranu, dijetne modifikacije i pripravke te enteralnu i parenteralnu prehranu.

U užem smislu klinička prehrana obuhvaća enteralnu i parenteralnu prehranu.

Nakon procjene stanja uhranjenosti te uvida u stanje i funkciju probavnog sustava (osobito tankog i debelog crijeva), odlučujemo o obliku kliničke prehrane koji se može i koji je poželjno provoditi tijekom liječenja (tablica 1.).

ABSTRACT

Enteral and parenteral nutrition are the main methods of nutritional support in clinical medicine. Enteral and parenteral nutrition are administered in order to prevent or correct nutritional deficits and malnutrition syndroms. A good knowledge of these methods of clinical nutrition ensures a proper and timely therapeutic approach. Enteral nutrition has numerous advantages in comparison with parenteral nutrition in patients with the preserved structure and function of the digestive system. Parenteral nutrition is a way of substituting the organism with water, energetic substrates and all other nutrients using the intravenous route in patients with counterindications for enteral feeding or in situations in which the administration of enteral feeding cannot meet all nutritional needs.

KEY WORDS: enteral feeding, parenteral feeding, indications, administration, complications

UOBIČAJENA PERORALNA PREHRANA I DIJETNE MODIFIKACIJE

Razvoj znanosti o prehrani u prošlom stoljeću kretao se od uočavanja i prepoznavanja problema nutritivnih deficita, njihove prevencije i rješavanja do razvoja modela uranotežene prehrane, s ciljem promicanja zdravlja i smanjenja rizika za nastanak pojedinih bolesti. Na prijelazu u novi milenij razvija se model optimalne prehrane koja pojačavanjem fizioloških funkcija osigurava stanje maksimalnog zdravlja. Radi se o konceptu poznatijem kao *funkcionalna hrana*. Peroralna prehrana koja se razvijala na temeljnim postulatima nutricionizma, najbolji je način nutritivne podrške i treba je uvijek provoditi ako je bolesnik sposoban uzimati hranu.

Ovaj način prehrane uključuje standardnu dijetu i različite dijetne prilagođene pojedinim kliničkim zahtjevima. Modificirane ili prilagođene dijetne razlikuju se ili u konzistenciji ili sadržajem nutritivnih pripravaka.

Ustanova: ¹KBC i Medicinski fakultet – Zagreb, ²Medicinski fakultet – Zagreb

Prispjelo: 16. 7. 2003.

Prihvaćeno: 23. 7. 2003.

Adresa za dopisivanje: Doc. dr. sc. Željko Krznarić, KBC Zagreb, Klinika za unutarnje bolesti, Kišpatićeva 12, 10000 Zagreb