

Coronavirus epidemic in Croatia: case fatality decline during summer?

Kristić, Ivica; Pehlić, Marina; Pavlović, Mirjana; Kolarić, Branko; Kolčić, Ivana; Polašek, Ozren

Source / Izvornik: **Croatian Medical Journal, 2020, 61, 501 - 507**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.3325/cmj.2020.61.501>

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:184:633979>

Rights / Prava: [Attribution-ShareAlike 4.0 International/Imenovanje-Dijeli pod istim uvjetima 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2025-01-18**



Repository / Repozitorij:

[Repository of the University of Rijeka, Faculty of Medicine - FMRI Repository](#)



Croat Med J. 2020;61:501-7

<https://doi.org/10.3325/cmj.2020.61.501>

Coronavirus epidemic in Croatia: case fatality decline during summer?

Ivica Kristić¹, Marina Pehlić², Mirjana Pavlović³, Branko Kolarić^{4,5}, Ivana Kolčić⁶, Ozren Polašek⁶

¹Department of Cardiology, University Hospital Split, Split, Croatia

²Department of Gynecology, University Hospital Split, Split, Croatia

³Clinic for Tumors, Sestre Milosrdnice University Hospital, Zagreb, Croatia

⁴Department of Social Medicine and Epidemiology, Medical Faculty, University of Rijeka, Rijeka, Croatia

⁵Andrija Štampar Teaching Institute of Public Health, Zagreb, Croatia

⁶Department of Public Health, University of Split School of Medicine, Split, Croatia

*The first two authors contributed equally.

Aim To describe the SARS-CoV-2 epidemic pattern in Croatia during February-September 2020 and compare the case fatality ratio (CFR) between spring and summer.

Methods National data were used to calculate the weekly and monthly CFRs, stratified by three age groups: 0-64, 65-79, and 80+ years. We also calculated the standardized mortality ratios (SMR) to offset the differences in age composition.

Results The epidemic consisted of the initial wave, a trough in June, and two conjoined summer waves, yielding 17 206 coronavirus disease 2019 cases and 290 deaths. While the number of confirmed cases nearly quadrupled during summer, case fatality estimates decreased; CFR in spring was 4.81 (95% confidence interval 3.91-5.71), compared with 1.24 (1.06-1.42) in summer. The SMR for summer was 0.45 (0.37-0.55), suggesting that the case fatality risk halved compared with spring. Cardiovascular comorbidity was an important risk factor for case fatality (SMR 2.63 [2.20-3.13] during spring and 1.28 [1.02-1.59] during summer). The risk of death in ventilated patients remained unchanged (SMR 0.98 [0.77-1.24]).

Conclusions The epidemic dynamics suggests summer decline in case fatality, except in ventilated patients. While the effect of comorbidity also decreased, cardiovascular comorbidity remained an important risk factor for death even during summer. A plethora of possible confounders and an ever-changing landscape of SARS-CoV-2 epidemic in Croatia require constant monitoring and evaluation, with an aim to prevent the uncontrolled spread of the virus and a disruption of health care functioning.

Received: November 1, 2020

Accepted: December 6, 2020

Correspondence to:

Ozren Polašek
University of Split School of
Medicine
Šoltanska 22
21000 Split, Croatia
ozren.polasek@mefst.hr

The understanding of the seasonal pattern of SARS-CoV-2 spread and coronavirus disease 2019 (COVID-19) symptoms severity is not of only academic significance, but plays a central role in the epidemic preparedness (1-4). For the Northern hemisphere, the issues currently of the greatest importance are the prediction of the autumn and winter disease spread and mortality (5). Previous articles have most commonly confirmed the expected role of seasonality (6), often with an explanation that higher temperature reduces the virus spread during summer (7-9). On the other hand, some studies have shown that we may not expect a substantial seasonality pattern, suggesting that SARS-CoV-2 may not behave like a common seasonal respiratory pathogen (10-12). A recent systematic review confirmed the existence of some degree of seasonality, but also stated that it may not be the only or the critical variable that will define the pandemic patterns (13).

There are various possible causes underlying the seasonal pattern of the pandemic, including changes in temperature and humidity (14-17), or wind speed and air pollution (18). Additionally, unfavorable conditions were reported to affect the ability of respiratory mucosa to fend off (19), which suggests a possible synergistic effect between numerous factors. An important distinction needs to be made by separately addressing the viral spread and case fatality, since the two might not exhibit similar patterns. The recognition of such differences may shed more light on the disease pathogenesis, and possibly even steer the path toward mechanisms of epidemic control. However, studies that made this distinction often report the expectation that for Europe, an increase in humidity will lower the number of new cases and deaths, while a rise in temperature will increase these parameters (20). Although such predictions are very sensitive to numerous modeling input parameters and assumptions, none can be superior to the epidemiological *ex post-facto* analysis. Several such articles suggested lowering of the summer case fatality (9,21), but this finding was not replicated systematically across all studies (22). Therefore, the aim of this study was to explore the epidemic pattern in Croatia, with a special focus on understanding the difference in the disease spread and case fatality during the spring and summer of 2020.

MATERIAL AND METHODS

The main source of information were the databases of the Croatian Institute of Public Health and the Croatian National Health Insurance Fund. The final data set encompassed the data from the period February-Sep-

tember 2020 (ending with the data available on September 09, 2020), accompanied by the available information on the testing outcomes, disease progression, the use of hospital services, and the disease outcome. In order to obtain the comorbidity information, we merged the data set with the Croatian National Health Insurance usage data from the previous three years. For this period, we defined all the ICD-10 disease categories attributed to a single patient, and used this as the information on comorbidity status. In order to perform a pooled analysis, we classified all comorbid conditions by the first letter of the ICD-10 classification code, meaning that only the disease category was used as comorbidity information (since we had no means of harmonizing the data to obtain more refined estimates). All of the comorbidity categories were scored as binary variables, allowing us to calculate the overall comorbidity load, defined as the sum of all positive results of comorbid conditions. This variable was only assumed as a relative proxy of the comorbidity status, as it suffers from numerous methodological risks and biases; it was based only on registered and diagnosed conditions, each ICD-10 category was treated as having equal weight, the disease severity or duration was not estimated, to name just the most obvious ones. In addition, we used three more binary comorbidity-related variables: the presence of any comorbidity, cardiovascular comorbidity, or tumor-related comorbidity.

All the data were sorted according to the epidemic duration into weeks and months, enabling us to analyze the selected parameters as the epidemic progressed. Notably, the data for September were included without the ability to infer the outcomes for some patients, therefore any estimates from September, especially those related to case fatality, are only treated as preliminary and provisional. Two main analytic periods were loosely defined as spring (February to May) and summer (June to September). Although this breakdown does not strictly conform to the official start and end of the seasons, we considered the analyzed periods to be good proxies for spring and summer.

All the data were routinely collected and anonymized at the source, before the export to the analytic file was made. No personal information data were available to the authors during any part of the study execution; therefore, no ethical approval for this study was sought.

Statistical analysis

The principal focus of the study was the calculation of the case fatality ratio (CFR), defined as the number of the de-

ceased among the total number of confirmed cases. The CFRs were calculated for every week and month of the epidemic, as well as for spring vs summer. However, for the purposes of this study, we used the diagnosis date as the reference point, regardless of disease duration and outcome. In effect, this meant that every case of death was scaled according to the date of diagnosis, in order to provide a better estimate of seasonality and offset the differences in disease duration (which sometimes extended for longer than a month). This also meant that the CFRs were less biased, since the disease duration effect was counteracted (as opposed to the aggregated intra-epidemic daily CFRs, which suffer from bias due to the outcome lag). For every point estimate of CFR, we also calculated the 95% confidence intervals.

In order to offset the differences in the age composition between units of observation, we calculated the standardized mortality ratios (SMR). For this purpose, we used the five-year breakdown according to age, with spring as the referent and summer as the target group; this meant that the SMRs reflected the age-adjusted spring vs summer differences. We also used SMRs in several more situations where comparisons were assumed to be confounded by the age structure.

The calculation of the delay from diagnosis to hospitalization was restricted to the period of 0-7 days, in order to remove possible bias due to other causes of hospitalization in previously positive patients. For a few deceased patients in whom the disease duration was not evident, as the disease duration indicator we used the time from diagnosis to death.

Categorical data were analyzed with the χ^2 test, correlations with the Pearson correlation test, while numerical data were analyzed with ANOVA or the *t* test, depending on the number of analyzed categories. We also used logistic regression for case fatality prediction, with age, sex, season, and comorbidity load as covariates. All the data were analyzed with R, with significance level set at $P < 0.05$.

RESULTS

From the beginning of the epidemic to the end of the study period, there were 17 206 cases of COVID-19 in Croatia, distributed in three outbreaks: the first corresponded to spring, while the second and third were overlapping in summer (Figure 1, right axis; Supplementary Table 1). The epidemic pattern was marked with several important events, namely the lockdown, including school closings and a ban on inter-city travel (March 13-23), and the decision on the mandatory use of masks in public areas and transport (July 13). Additionally, there was a three-week period in June when the newly diagnosed cases and deaths neared zero (Figure 1). During the entire epidemic, the case fatality ratios varied, with the highest values among the oldest patients, and lower values during the summer months (Figure 1, left axis; Supplementary Table 1). The overall case fatality for spring was 4.8% (108 cases), while that for summer was 1.2% (182 cases; $P < 0.001$). Notably, the CFRs exhibited the highest decline in summer in the youngest age group and the lowest in the oldest age group (Supplementary Table 2).




FIGURE 1. Comprehensive scheme of the SARS-CoV-2 epidemic in Croatia. The left axis denotes the case fatality ratio (CFR), with cases of deaths being classified according to the date of diagnosis (mortality lag removed). The right axis denotes the daily number of newly diagnosed cases (gray bars). Case fatality ratios were stratified to three age groups (0-64 years, 65-79 years, ≥80 years old).

The standardized mortality ratio confirmed a reduction in case fatality, with SMR=0.45 (95% CI 0.37-0.55), which was similar in men (0.45 [0.34-0.58]) and women (0.45 [0.32-0.60]). The presence of any kind of comorbid conditions nearly doubled the SMR in spring (2.02 [1.66-2.44]), while it decreased to having no effect in summer (1.00 [0.46-1.89]). The SMRs for cardiovascular diseases decreased substantially (2.63 [2.20-3.13] in spring vs 1.28 [1.02-1.59] in summer). On the contrary, the SMR for tumors was 1.12 [0.83-1.48] in spring, while in summer it seemed to have a somewhat protective effect, with a SMR of 0.45 (0.36-0.56). Finally, the logistic regression model that investigated the risk factors for death outcome suggested that female sex and summer infection were protective, while older age and accumulating comorbidity were detrimental effects; notably, the odds ratio for the oldest age group was substantial (Table 1).

There were 57 patients requiring assisted ventilation in spring (2.5% out of confirmed cases), compared with 96 patients (0.6%) in summer ($P < 0.001$). The SMR for summer vs spring suggested that the ratio for ventilator use more than halved (0.41 [0.33-0.50]). However, the progression from ventilation to death did not show significant differences between the seasons; 37 (65%) of ventilated patients deceased in spring, compared with 70 (73%) in summer ($P = 0.297$). Accordingly, the SMR for dying after the ventilation use did not differ between summer and spring (0.98 [0.77-1.24]).

Delay of hospitalization from the time of diagnosis increased during the observed period, from 1.00 ± 0.82 days in February, 1.17 ± 1.40 in March, 0.93 ± 1.14 in April, 0.85 ± 0.68 in May, 1.94 ± 2.13 in June, 1.85 ± 2.18 in June, 2.20 ± 2.19 in August to 2.31 ± 2.36 in September (monthly comparison ANOVA $P < 0.001$; spring vs summer $P < 0.001$).

TABLE 1. Predictors of death outcomes of coronavirus 2019 in the Croatian population from March to September 2020, logistic regression

Predictor	Odds ratio	
	[95% confidence interval]	<i>P</i>
Sex		
male (Ref.)	1.00	<0.001
female	0.51 [0.39-0.66]	
Age (years)		
0-64 (Ref.)	1.00	<0.001
65-79	19.6 [13.6-28.4]	
80 and more	58.2 [39.5-85.8]	
Comorbidity load	1.07 [1.03-1.11]	0.001
Season		
spring (Ref.)	1.00	<0.001
summer	0.50 [0.38-0.66]	

Even the percent of hospitalized patients was different; there were 886 hospitalized cases in spring (39.3%), compared with 1784 in summer (11.9%; $P < 0.001$).

Duration of hospital treatment declined significantly as the epidemic progressed, from 22.8 ± 11.8 days in February to 7.6 ± 6.0 days in September ($P < 0.001$); the duration of hospital treatment did not show any coherent pattern in regards to the disease outcome (Supplementary Table 3). Duration of the hospital stay was the longest in the oldest age group and the shortest in the youngest age group (80+ years old: 16.6 ± 11.0 days; 64-79 years old: 13.7 ± 10.8 ; 0-64 years old: 9.4 ± 8.0 days). The comorbidity load was significantly associated with the duration of hospital stay ($P < 0.001$; $r = 0.16$).

DISCUSSION

This is the first systematic attempt to describe the dynamics and outcome of the first eight months of the SARS-CoV-2 epidemic in Croatia and compare the mortality patterns in spring and summer. While the number of newly diagnosed cases increased substantially during summer, the CFR declined. The summer case fatality was nearly halved in comparison with the spring rate, with the greatest reduction observed in the youngest age group. Additionally, the presence of any comorbidity doubled the risk of dying in spring, while the risk declined to a no detectable effect in summer. The exception was cardiovascular disease, which retained its detrimental effect. This suggests that either comorbidity could only be a conditional risk factor that operates in unfavorable conditions or that comorbidity had an additive detrimental effect in spring, which was dissolved in summer. Unfortunately, previously published papers provide little evidence as to the extent of comorbidity contribution, which ranges from substantial (23) to none (24), and any level of contribution in between.

The summer case fatality rates decline could have been caused by a number of factors, broadly classifiable into pathogen-related, host-related, or environmental (and any combination or interaction thereof). Enveloped viruses are known to have stronger seasonal occurrence (25,26), possibly due to the lipid (27) or glycosylation properties of their envelope (28,29). On the other hand, human mucosa also shows seasonal changes, probably affecting mucosal barrier and clearance mechanisms, as a response to the environmental conditions (17,30). Respiratory tract microbiome was also shown to exhibit seasonal patterns (31,32), possibly attributing to the seasonal pattern of respiratory infections. Interestingly, challenge studies of cold air inha-

lation have failed to replicate the seasonal effects (33-35), suggesting that the exact pathogenetic mechanisms of increased winter-spring burden of respiratory pathogens are very complex. Therefore, the lack of clarity and understanding of causal relationships at this stage reduce the ability to use seasonal effects mechanisms as the COVID-19 epidemic controlling measures.

Cardiovascular comorbidity seemed to have a strong modifying effect, which persisted throughout both spring and summer. Previous studies have already established this link (36-38), reporting a strong modifying effect on the COVID-19 prognosis (39-42). However, studies have reported not only that pre-existing cardiovascular comorbidity affects the prognosis, but also that SARS-CoV-2 may trigger myocardial injury, acute coronary syndrome, arrhythmias, and venous thromboembolism (39,43-45). As a result, COVID-19 patients often have the signs of myocardial injury (39,46-48), with more than half of all COVID-related deaths exhibiting signs of heart failure (49). A large study from Italy suggested that none of the comorbid conditions increased the mortality risk, but that the risk was conveyed by the impaired renal function, elevated C-reactive protein, and advanced age (50). Interestingly, in our study the detrimental effects of tumor comorbidity were dissolved in summer, conveying even a significant reduction of case fatality risk. Similar results were already reported (51), suggesting that cancer effects on the risk of dying in COVID-19 patients might even be case-, stage-, or grade-specific. Alternatively, patients with cancer might have more strictly adhered to preventative measures, which led to a lower risk of dying in summer. All these results suggest that a proper answer to the question of comorbidity contribution will have to be given in a form of a systematic review and meta-analysis, preferably using individual-level data (52).

The data presented here also suggest that clinical approach to patients with COVID-19 has likely changed. Hospitalization due to COVID-19 during summer was three times as less likely as it was during spring. While the duration of the hospital stay declined, the delay from diagnosis to hospitalization was prolonged. This means that we no longer hospitalize all the patients, and that we possibly have better treatment outcomes, which are reflected in a shorter stay. This confirms the complexity of the COVID-19 management within the health care system (53).

This study has numerous limitations. The quality of the input data had not been assessed or validated. Further, the data collection process was not harmonized or created for this

purpose, especially early on during the epidemic course, which could be a major source of risk of bias and error. Next, heterogeneity of treatment plans across different hospitals and across the epidemic duration, including diagnosis criteria, severity assessment, hospital admission, or ventilator use were changing, meaning that their direct pooling may yield spurious results. This epidemic has produced a substantial societal effect, meaning that the epidemic landscape might change quickly, as it has in the past, making all these estimates time-sensitive. The changing capacity and pattern of testing could have also affected the results, possibly by underestimating the real number of cases (54,55). The extent and direction of this is hard to estimate; low testing may underestimate the true number of positive results, while selective under-testing (and consequent under-diagnosing) of certain population groups could have substantially skewed the results; this is why the improvement of testing capacities is a critical step forward (56). Finally, the methodological framework that uses CFR and SMR is also a possible source of problems, including the analysis of the data from an ongoing epidemic, which reduced the usability of estimates toward the end of observation period.

The results of this study demonstrate the apparent changes in the outcomes during summer, which systematically suggest favorable direction and confirm previous predictions of the virus and epidemic seasonality. Therefore, this study provides an account of the mortality patterns, ultimately aiming to contribute to a better epidemic management and the prevention of health care functioning disruptions.

Acknowledgments We thank the staff of the Croatian Institute of Public Health, who maintain and curate the databases that were the primary source of data for this study.

Funding None.

Ethical approval None sought.

Author contributions OP, IKr, IK, and BK conceived the idea and planned the study. IKr, MPe, and MPa assembled the data and performed the literature search. OP performed statistical analysis. IKr, MPa, and MPe drafted the manuscript. All authors participated in the draft development, critically revised the manuscript, and approved the final version of the manuscript. All authors are accountable for all aspects of the work.

Competing interests OP and BK are scientific advisors to the Government of the Republic of Croatia for COVID-19 response, which had no influence in the decision to prepare the manuscript, its preparation or the selection of the journal where the manuscript would be submitted. OP is a member of the Editorial Board of the *Croatian Medical Journal*. To ensure that any possible conflict of interest relevant to the journal has been addressed, this article was reviewed according to best practice guidelines of international editorial organizations. All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organization for the submitted work; no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years; no other relationships or activities that could appear to have influenced the submitted work.

References

- 1 Watts CH, Vallance P, Whitty CJM. Coronavirus: global solutions to prevent a pandemic. *Nature*. 2020;578:363. Published online 02 20, 2020. [Medline:32071448](#)
- 2 Sajadi MM, Habibzadeh P, Vintzileos A, Shokouhi S, Miralles-Wilhelm F, Amoroso A. Temperature, humidity, and latitude analysis to estimate potential spread and seasonality of coronavirus disease 2019 (COVID-19). *JAMA Netw Open*. 2020;3:e2011834. [Medline:32525550](#)
- 3 Sheikh A, Sheikh Z, Dhama S, Sridhar D. What's the way out? Potential exit strategies from the COVID-19 lockdown. *J Glob Health*. 2020;10:010370. [Medline:32566161](#)
- 4 Primorac D, Matišić V, Molnar V, Bahtijarević Z, Polašek O. Pre-season football preparation in the era of COVID-19: Croatian Football Association Model. *J Glob Health*. 2020;10:010352. [Medline:32426120](#)
- 5 Rudan I. Answering the initial 20 questions on COVID-19 (January-February 2020). *J Glob Health*. 2020;10:010106. [Medline:32566149](#)
- 6 Audi A, Allbrahim M, Kaddoura M, Hijazi G, Yassine HM, Zaraket H. Seasonality of respiratory viral infections: Will COVID-19 follow suit? *Front Public Health*. 2020;8:567184. [Medline:33042956](#)
- 7 Malki Z, Atlam ES, Hassani AE, Dagnew G, Elhosseini MA, Gad I. Association between weather data and COVID-19 pandemic predicting mortality rate: Machine learning approaches. *Chaos Solitons Fractals*. 2020;138:110137. [Medline:32834583](#)
- 8 Adedokun KA, Olaninmoye AO, Mustapha JO, Kamorudeen RT. A close look at the biology of SARS-CoV-2, and the potential influence of weather conditions and seasons on COVID-19 case spread. *Infect Dis Poverty*. 2020;9:77. [Medline:32586369](#)
- 9 Mandal CC, Panwar MS. Can the summer temperatures reduce COVID-19 cases? *Public Health*. 2020;185:72-9. [Medline:32574871](#)
- 10 Jamil T, Alam I, Gojobori T, Duarte CM. No evidence for temperature-dependence of the COVID-19 epidemic. *Front Public Health*. 2020;8:436. [Medline:32984240](#)
- 11 Benedetti F, Pachetti M, Marini B, Ippodirino R, Gallo RC, Ciccozzi M, et al. Inverse correlation between average monthly high temperatures and COVID-19-related death rates in different geographical areas. *J Transl Med*. 2020;18:251. [Medline:32576227](#)
- 12 Jin K, Min J, Jin X. Re:Can the summer temperature reduce COVID-19 cases? *Public Health*. 2020;187:186. [Medline:33002712](#)
- 13 Mecenaz P, Bastos R, Vallinoto ACR, Normando D. Effects of temperature and humidity on the spread of COVID-19: A systematic review. *PLoS One*. 2020;15:e0238339. [Medline:32946453](#)
- 14 He J, Chen G, Jiang Y, Jin R, Shortridge A, Agusti S, et al. Comparative infection modeling and control of COVID-19 transmission patterns in China, South Korea, Italy and Iran. *Sci Total Environ*. 2020;747:141447. [Medline:32771775](#)
- 15 Perone G. The determinants of COVID-19 case fatality rate (CFR) in the Italian regions and provinces: An analysis of environmental, demographic, and healthcare factors. *Sci Total Environ*. 2020;755:142523. [Medline:33022464](#)
- 16 Fazzini M, Baresi C, Bisci C, Bna C, Cecili A, Giuliacci A, et al. Preliminary Analysis of relationships between COVID19 and climate, morphology, and urbanization in the Lombardy Region (Northern Italy). *Int J Environ Res Public Health*. 2020;17. [Medline:32977546](#)
- 17 Lauc G, Markotić A, Gornik I, Primorac D. Fighting COVID-19 with water. *J Glob Health*. 2020;10:010344. Published online 05 07, 2020. [Medline:32373322](#)
- 18 Lolli S, Chen YC, Wang SH, Vivone G. Impact of meteorological conditions and air pollution on COVID-19 pandemic transmission in Italy. *Sci Rep*. 2020;10:16213. Published online 10 03, 2020. [Medline:33004925](#)
- 19 Barbet JP, Chauveau M, Labbé S, Lockhart A. Breathing dry air causes acute epithelial damage and inflammation of the guinea pig trachea. *J Appl Physiol*. 1988;64:1851-7. [Medline:3391888](#)
- 20 Meo SA, Abukhalaf AA, Alomar AA, Sumaya OY, Sami W, Shafi KM, et al. Effect of heat and humidity on the incidence and mortality due to COVID-19 pandemic in European countries. *Eur Rev Med Pharmacol Sci*. 2020;24:9216-25. [Medline:32965017](#)
- 21 Carta MG, Scano A, Lindert J, Bonanno S, Rinaldi L, Fais S, et al. Association between the spread of COVID-19 and weather-climatic parameters. *Eur Rev Med Pharmacol Sci*. 2020;24:8226-31. [Medline:32767354](#)
- 22 Meo SA, Abukhalaf AA, Alomar AA, Al-Beeshi IZ, Alhwwikan A, Shafi KM, et al. Climate and COVID-19 pandemic: effect of heat and humidity on the incidence and mortality in world's top ten hottest and top ten coldest countries. *Eur Rev Med Pharmacol Sci*. 2020;24:8232-8. [Medline:32767355](#)
- 23 Gu T, Chu Q, Yu Z, Fa B, Li A, Xu L, et al. History of coronary heart disease increased the mortality rate of patients with COVID-19: a nested case-control study. *BMJ Open*. 2020;10:e038976. [Medline:32948572](#)
- 24 Brar G, Pinheiro LC, Shusterman M, Swed B, Reshetnyak E, Soroka O, et al. COVID-19 Severity and Outcomes in Patients With Cancer: A Matched Cohort Study. *J Clin Oncol*. 2020;...:JCO2001580. [Medline:32986528](#)
- 25 Price RHM, Graham C, Ramalingam S. Association between viral seasonality and meteorological factors. *Sci Rep*. 2019;9:929. [Medline:30700747](#)
- 26 Schoeman D, Fielding BC. Coronavirus envelope protein: current knowledge. *Virology*. 2019;16:69. [Medline:31133031](#)
- 27 Polozov IV, Bezrukov L, Gawrisch K, Zimmerberg J. Progressive ordering with decreasing temperature of the phospholipids of influenza virus. *Nat Chem Biol*. 2008;4:248-55. [Medline:18311130](#)
- 28 An Y, Parsons LM, Jankowska E, Melnyk D, Joshi M, Cipollo JF. N-glycosylation of seasonal influenza vaccine hemagglutinins: implication for potency testing and immune processing. *J Virol*. 2019;93. [Medline:30355697](#)

- 29 Wang YF, Chang CF, Tsai HP, Chi CY, Su IJ, Wang JR. Glycan-binding preferences and genetic evolution of human seasonal influenza A(H3N2) viruses during 1999-2007 in Taiwan. *PLoS One*. 2018;13:e0196727. [Medline:29746492](#)
- 30 Kudo E, Song E, Yockey LJ, Rakib T, Wong PW, Homer RJ, et al. Low ambient humidity impairs barrier function and innate resistance against influenza infection. *Proc Natl Acad Sci U S A*. 2019;116:10905-10. [Medline:31085641](#)
- 31 Schoos AM, Kragh M, Ahrens P, Kuhn KG, Rasmussen MA, Chawes BL, et al. Season of birth impacts the neonatal nasopharyngeal microbiota. *Children (Basel)*. 2020;7. [Medline:32403236](#)
- 32 Mika M, Mack I, Korten I, Qi W, Aebi S, Frey U, et al. Dynamics of the nasal microbiota in infancy: a prospective cohort study. *The Journal of Allergy and Clinical Immunology*. 2015;135:905-12e11. [Medline:25636948](#)
- 33 Eccles R, Wilkinson JE. Exposure to cold and acute upper respiratory tract infection. *Rhinology*. 2015;53:99-106. [Medline:26030031](#)
- 34 Eccles R. An explanation for the seasonality of acute upper respiratory tract viral infections. *Acta Otolaryngol*. 2002;122:183-91. [Medline:11936911](#)
- 35 Mourtzoukou EG, Falagas ME. Exposure to cold and respiratory tract infections. *Int J Tuberc Lung Dis*. 2007;11:938-43. [Medline:17705968](#)
- 36 Madjid M, Safavi-Naeini P, Solomon SD, Vardeny O. Potential effects of coronaviruses on the cardiovascular system: a review. *JAMA Cardiol*. 2020;5:831-40. [Medline:32219363](#)
- 37 Driggin E, Madhavan MV, Bikdeli B, Chuich T, Laracy J, Biondi-Zoccai G, et al. Cardiovascular considerations for patients, health care workers, and health systems during the COVID-19 pandemic. *J Am Coll Cardiol*. 2020;75:2352-71. [Medline:32201335](#)
- 38 Clerkin KJ, Fried JA, Raikhelkar J, Sayer G, Griffin JM, Masoumi A, et al. COVID-19 and cardiovascular disease. *Circulation*. 2020;141:1648-55. [Medline:32200663](#)
- 39 Shi S, Qin M, Shen B, Cai Y, Liu T, Yang F, et al. Association of cardiac injury with mortality in hospitalized patients with COVID-19 in Wuhan, China. *JAMA Cardiol*. 2020;5:802-10. [Medline:32211816](#)
- 40 Barton LM, Duval EJ, Stroberg E, Ghosh S, Mukhopadhyay S. COVID-19 autopsies, Oklahoma, USA. *Am J Clin Pathol*. 2020;153:725-33. [Medline:32275742](#)
- 41 Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med*. 2020;8:420-2. [Medline:32085846](#)
- 42 Vrsalovic M, Vrsalovic Presecki A. Cardiac troponins predict mortality in patients with COVID-19: A meta-analysis of adjusted risk estimates. *J Infect*. 2020;81:e99-100. [Medline:32416124](#)
- 43 Guo T, Fan Y, Chen M, Wu X, Zhang L, He T, et al. Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). *JAMA Cardiol*. 2020;5:811-8. [Medline:32219356](#)
- 44 Shi S, Qin M, Cai Y, Liu T, Shen B, Yang F, et al. Characteristics and clinical significance of myocardial injury in patients with severe coronavirus disease 2019. *Eur Heart J*. 2020;41:2070-9. [Medline:32391877](#)
- 45 Liu K, Fang YY, Deng Y, Liu W, Wang MF, Ma JP, et al. Clinical characteristics of novel coronavirus cases in tertiary hospitals in Hubei Province. *Chin Med J (Engl)*. 2020;133:1025-31. [Medline:32044814](#)
- 46 Creel-Bulos C, Hockstein M, Amin N, Melhem S, Truong A, Sharifpour M. Acute cor pulmonale in critically ill patients with Covid-19. *N Engl J Med*. 2020;382:e70. [Medline:32374956](#)
- 47 Zheng YY, Ma YT, Zhang JY, Xie X. COVID-19 and the cardiovascular system. *Nat Rev Cardiol*. 2020;17:259-60. [Medline:32139904](#)
- 48 Tersalvi G, Vicenzi M, Calabretta D, Biasco L, Pedrazzini G, Winterton D. Elevated troponin in patients with coronavirus disease 2019: possible mechanisms. *J Card Fail*. 2020;26:470-5. [Medline:32315733](#)
- 49 Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395:1054-62. [Medline:32171076](#)
- 50 Di Castelnuovo A, Bonaccio M, Costanzo S, Gialluisi A, Antinori A, Berselli N, et al. Common cardiovascular risk factors and in-hospital mortality in 3,894 patients with COVID-19: survival analysis and machine learning-based findings from the multicentre Italian CORIST Study. *Nutr Metab Cardiovasc Dis*. 2020. Published online 09 12, 2020. [Medline:32912793](#)
- 51 Kuderer NM, Choueiri TK, Shah DP, Shyr Y, Rubinstein SM, Rivera DR, et al. Clinical impact of COVID-19 on patients with cancer (CCC19): a cohort study. *Lancet*. 2020;395:1907-18. [Medline:32473681](#)
- 52 Ioannidis JP, Greenland S, Hlatky MA, Khoury MJ, Macleod MR, Moher D, et al. Increasing value and reducing waste in research design, conduct, and analysis. *Lancet*. 2014;383:166-75. [Medline:24411645](#)
- 53 Flaatten H, Van Heerden V, Jung C, Beil M, Leaver S, Rhodes A, et al. The good, the bad and the ugly: pandemic priority decisions and triage. *J Med Ethics*. 2020. Published online 06 12, 2020. [Medline:32522814](#)
- 54 Morel CM, Lindahl O, Özenci V. Lessons from COVID-19 on the role of the state and the market in providing early testing. *J Glob Health*. 2020;10:020330. [Medline:33110530](#)
- 55 Sousa-Pinto B, Fonseca JA, Oliveira B, Cruz-Correia R, Rodrigues PP, Costa-Pereira A, et al. Simulation of the effects of COVID-19 testing rates on hospitalizations. *Bull World Health Organ*. 2020;98:299. [Medline:32514192](#)
- 56 Vukičević D, Polašek O. Optimizing the diagnostic capacity for COVID-19 PCR testing for low resource and high demand settings: the development of information-dependent pooling protocol. *J Glob Health*. 2020;10:020515.