

Factors associated with posttraumatic stress disorder and depression in warsurvivors displaced in Croatia

Letica-Crepulja, Marina; Salcioglu, Ebru; Frančišković, Tanja; Basoglu, Metin

Source / Izvornik: **Croatian Medical Journal, 2011, 52, 709 - 717**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:184:008878>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-05-22**



Repository / Repozitorij:

[Repository of the University of Rijeka, Faculty of Medicine - FMRI Repository](#)



doi: 10.3325/cmj.2011.52.709

Factors associated with posttraumatic stress disorder and depression in war-survivors displaced in Croatia

Aim To examine the role of perceived stressfulness of trauma exposure and economic, social, occupational, educational, and familial adaptation after trauma in posttraumatic stress disorder (PTSD) and depression in displaced war survivors.

Methods A cross-sectional survey was conducted between March 2000 and July 2002 with a sample of 173 internally displaced persons or refugees and 167 matched controls in Croatia. Clinical measures included Structured Clinical Interview for DSM-IV and Clinician-Administered PTSD Scale.

Results Displaced war survivors reported the exposure to a mean \pm standard deviation of 13.1 ± 8.3 war stressors, including combat, torture, serious injury, death of close persons, and loss of property. Compared to controls, they reported higher rates of marked to severe impact of war on family (16.2% vs 51.6%), social (7.2% vs 43.5%), economic (12.6% vs 55.4%), occupational (1.8% vs 15.9%), and educational (2.4% vs 8.8%) adaptation. In two logistic regression analyses, the strongest predictor of PTSD and depression was high level of perceived distress during trauma exposure. PTSD but not depression was associated with economic, social, occupational, educational, and familial adaptation after trauma.

Conclusion Displaced survivors who experienced multiple war events perceived greater negative impact of war on their life domains compared to individuals who lived in a war setting but had no trauma exposure. The most important determinant of psychological outcomes was perceived stressfulness of war stressors. Although post-trauma adaptation in different life spheres had an impact, its effect was not robust and consistent across disorders. These findings suggest that it would be effective to use a trauma-focused approach in rehabilitation of war survivors.

Marina Letica-Crepulja¹,
Ebru Salcioglu^{2,3}, Tanja
Frančičković^{1,4}, Metin
Basoglu^{2,3}

¹Regional Psychotrauma Centre Rijeka and Department of Psychological medicine, Psychiatric Clinic, Clinical Hospital Centre Rijeka, Rijeka, Croatia

²Trauma Studies Unit, Institute of Psychiatry, King's College, University of London, London, UK

³Istanbul Center for Behavior Research and Therapy, Istanbul, Turkey

⁴Department for Psychiatry and Psychological Medicine, University of Rijeka School of Medicine, Rijeka, Croatia

Received: June 29, 2011

Accepted: November 23, 2011

Correspondence to:

Marina Letica-Crepulja
Regional Psychotrauma Centre
Rijeka and Department of
Psychological Medicine
Psychiatric Clinic, Clinical Hospital
Centre Rijeka
Cambierieva 17
51 000 Rijeka, Croatia
leticacrepulja@yahoo.com

Armed conflicts, wars, and associated displacement affect large numbers of people in the world (1). According to the United Nations High Commissioner of Refugees (UNHCR) (2), at the end of 2010 the number of forcibly displaced people in the world was 43.7 million. Some war survivors are displaced within the borders of their countries as internally displaced persons, while others are displaced to other countries as refugees. In addition to war-related stressor events they have been through, such displaced war survivors are believed to experience enduring contextual stressors, including socioeconomic disadvantage and poverty, changes in family structure and functioning, loss of social support, lack of access to education, overcrowded housing, hostility and racism, acculturation difficulties, marginalization and isolation, and cultural bereavement (3-7). These stressors are claimed to cause general psychological distress (8), but there are few studies that examined their contribution to mental health while controlling for the impact of other potentially important variables.

The most common mental health outcome of exposure to war-related traumatic stressors is posttraumatic stress disorder (PTSD) and depression (9). In a meta-analysis of 181 surveys comprising 81 866 refugees and other conflict-affected persons from 40 countries, the prevalence rate of PTSD across all surveys was 30.6% and that of depression was 30.8% (1). In this study, the factors that showed strong association with PTSD were cumulative exposure to traumatic events, time since conflict, and assessed level of political terror, while the factors associated with depression were cumulative exposure to traumatic events, time since trauma, torture, and residency status. An important factor that this meta-analysis did not control for (due to lack of data) was psychological processes during trauma exposure, including emotional reactions, perception of life threat, or dissociation during trauma, which have been determined to be the strongest predictors of PTSD (10-14). Indeed, in a series of studies involving war and torture survivors, distress and loss of control during exposure to traumatic stressors (9,15-19) emerged as the strongest predictors of PTSD when the effects of the number of trauma events (cumulative exposure) were statistically controlled for in a regression analysis.

Some studies identified older age (20,21), female gender (3,21), and psychiatric history and current illness (22) as predictors of traumatic stress reactions but it is not possible to reach a conclusion about these associations because these studies rarely took into account the effects of trauma exposure characteristics.

Political violence and terror during the 1991-1995 war in Croatia and the 1992-1995 war in Bosnia and Herzegovina were categorized as the highest level on the Political Terror Scale (23). Four years of conflict resulted in the displacement of almost 900 000 Croatian citizens of all ethnicities inside and outside the country. Serb secession in central and eastern parts of the country caused an internal displacement in Croatia of over 550 000 ethnic Croats. The war that erupted in Bosnia and Herzegovina in April 1992 resulted in a large population movement. Over the course of the conflict, Croatia accepted 403 000 refugees from the neighboring country (24). The total number of displaced war survivors in Croatia at the end of 1992 was more than 10% of the total Croatian population (25). In this study, we examined the factors associated with PTSD and depression in displaced war survivors in Croatia. Our aim was to determine the role of economic, social, occupational, educational, and familial adaptation after trauma on PTSD and depression while controlling for the effects of psychological processes during trauma exposure as well as demographic and personal history characteristics. Specifically, we hypothesized that 1) displaced war survivors who experienced multiple war-related potentially traumatic events would report greater negative impact on economic, social, occupational, educational, and familial adaptation compared to a control group of individuals who, despite living in a war setting, had no personal exposure to potentially traumatic events; 2) PTSD and depression would be most strongly associated with perceived stressfulness or uncontrollability of war-related traumatic stressors; and 3) economic, social, occupational, educational, and family adaptation in the aftermath of exposure to traumatic incidents would be an independent but weaker predictor of PTSD and depression.

METHODS

Sampling procedures

Our research was part of a large survey conducted between March 2000 and July 2002. Details of the study methodology were presented in the main report (9). Briefly, a cross-sectional survey was conducted with a population-based sample of 1358 war survivors who had experienced at least 1 war-related stressor (combat, torture, internal displacement, refugee experience, siege, and/or aerial bombardment) from 4 sites in former Yugoslavia (Belgrade, Serbia and Montenegro; Rijeka, Croatia; Sarajevo, Bosnia-Herzegovina; and Banja Luka, Republic of Srpska, Bosnia-Herzegovina), accessed through linkage sampling. For the

purposes of this study, from the total sample we selected 173 internally displaced persons and refugees (displaced war survivors hereafter) living in the region around Rijeka, Croatia. The internally displaced persons were mostly Croats from the Vukovar region. Inclusion criteria were age 18 through 65 years, literacy, absence of past or present psychotic illness, and written consent for participation in the study. Target sampling (26) was used to ensure adequate representation of survivor group of interest. Sampling bias in the targeted group was minimized as much as possible by using linkage sampling and snowballing (27). This strategy involved tracing and contacting survivors in the community through key informants who were asked to make a list of their friends or acquaintances with an experience of a particular index stressor, disregarding any available information on their psychological status. These survivors were then contacted and invited to participate in the study. Once the interview was completed, each survivor was asked to list all friends or acquaintances with a similar trauma experience. This "snowballing" process continued until the targeted sample size for a particular index stressor was achieved. The study design included a control group ($n = 167$) matched on a 1:1 basis with the community participants on sex, age, and education. Although most control participants were exposed to war scenes on television and/or had relatives or friends who had experienced war events, they had no direct exposure to war stressors.

Measures

Psychiatric outcome was assessed with the Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) (SCID-I/NP, version 2) (28). Assessment of current and lifetime PTSD was made with the Clinician-Administered PTSD Scale (CAPS) (29). Demographic, trauma characteristics, and direct social impact of war and displacement on economic, social, family, occupational, and educational functioning were assessed with the Semi-Structured Interview for Survivors of War (SISOW) (9), which was modified from the Semi-Structured Interview for Survivors of Torture (30) used in previous studies. The SISOW includes an Exposure to War Stressors Scale (EWSS) that elicits information on 54 war-related stressor events (eg, experience of combat, torture, witnessing atrocities, internal or external displacement, loss of close ones, etc) and perceived distress and uncontrollability associated with each event. Measurement included a Global Distress Rating (0 = not at all distressing, 4 = extremely distressing) and a Global Sense of Control Rating (0 = completely in control and 4 = not at all in control/entirely helpless) to assess dis-

placed war survivors' overall perceived distress or loss of control during exposure to such war stressors. The SISOW also provided 5 variables assessing the impact of trauma on economic, social, family, occupational, and educational life (0 = no impact at all, 4 = extremely severe impact). These variables reflected the cumulative negative impact of all traumatic events identified by the EWSS on various life spheres by comparing pre-war status with the present. In this assessment, interviewers distinguished as much as possible the impact of the traumatic stressors from the impact of PTSD on various life spheres. Specifically, the assessors were instructed to identify the trauma event responsible for the impact in a particular life sphere considering the nature of the latter (eg, physical disability, loss of job or social status, loss of educational opportunities, death in or separation from the family) and not to evaluate the negative aspect of PTSD symptoms (eg, on marital or family adaptation or work functioning).

Interviews

The interviews were conducted by 5 trained professionals in Rijeka, Croatia. The interviews were evaluated for validity of diagnoses, rapport, adherence to the assessment protocol for each instrument, and overall quality of assessment. The concordance rates (kappa values) were 0.81 for PTSD, 0.84 for depression, and ranged from 0.83 to 0.99 for SISOW items in the overall sample (9). The interviews were usually completed in 1 or 2 sessions. Written informed consent was obtained for all study procedures. Strict confidentiality was emphasized. All study participants were offered an interview fee of US \$10 and those in need of help were offered psychiatric treatment.

Data analyses

Data were analyzed with SPSS, version 14.0 (SPSS Inc., Chicago, IL, USA). Sample characteristics were examined with frequency and descriptive statistics. Displaced war survivors were compared to matched controls on study variables of interest to produce evidence in support of the validity of the economic, social, family, occupational, and educational adaptation after trauma using McNemar test for related samples and paired t tests. Pearson correlation was used to examine the intercorrelations among impact of trauma on economic, social, family, occupational, and educational adaptation variables. As most of these variables were highly intercorrelated a composite measure representing global impact of trauma on life was computed for use in prediction analyses to avoid prob-

lems of multicollinearity. Two logistic regression analyses examined the factors related to current PTSD and current major depressive episode. Logistic regression allows one to predict a discrete outcome such as disease/no disease from a set of independent variables that may be continuous, discrete, dichotomous, or a mix (31). As in multiple regression analysis, the independent variables can be correlated with one another to varying degrees and the analysis takes into account these correlations in predicting the outcome. The goal of logistic regression is to correctly predict the category of outcome for individual cases using the most parsimonious model. To accomplish this goal, a model is created that includes all predictor variables that are useful in predicting the response variable. Since logistic regression calculates the probability of success over the probability of failure, the results of the analysis are in the form of an odds ratio. A Wald test was used to test the statistical significance of each coefficient of predictor variable (β). In this study, a total of 11 independent variables were entered in the logistic regression analyses, which included age, sex (1 = male, 2 = female), marital status (0 = single, 1 = married), education (1 = primary school, 2 = secondary school, 3 = high school, 4 = university/post-graduate), income level (1 = low, 4 = very good), personal and/or family history of psychiatric illness (0 = no, 1 = yes), time since last trauma event (months), global distress (0 = not at all distressing, 4 = extremely distressing) and control during trauma (0 = completely in control and 4 = not at all in control/entirely helpless) ratings, and the composite measure of global impact of trauma on life (total score on 5 variables ranging 0 to 20).

RESULTS

Sample characteristics

The socio-demographic characteristics of the sample are given as web-extra material. There was a balanced distribution of male and female participants in the sample. The majority of the displaced war survivors were Croats ($n = 148$, 85.5%) and Catholics ($n = 136$, 78.6%). The mean \pm standard deviation time since the most distressing traumatic event was 9.0 ± 1.6 years. Participants were exposed to a mean of 13.1 ± 8.3 war-related stressors. Some of these stressors included combat experience ($n = 41$, 23.7%), prisoner of war experience ($n = 34$, 19.7%), torture ($n = 43$, 24.9%), serious injury ($n = 10$, 5.8%), death of close persons ($n = 83$, 48.0%), exposure to aerial bombardment ($n = 82$, 47.4%), and sudden loss of property ($n = 123$, 71.1%). About 70% of the participants reported severe/extremely severe distress, while 94% reported slight to extreme loss of control during exposure to such traumatic stressors. Sixty-one participants (35.5%) had lifetime PTSD, 39 (22.5%) had current PTSD, 22 (12.7%) had current depression, 20 (11.6%) had past depression, 16 (9.2%) had alcohol abuse, and 10 (5.8%) had dysthymic disorder. Rates of the other psychiatric diagnoses were lower than 3%.

The matching procedure resulted in groups similar in sex, age, education, and marital status (Table 1). There was no difference between the groups in reported rates of personal and family history of psychiatric illness. As would be expected, displaced war survivors reported significantly

TABLE 1. Comparison of displaced war survivors and matched controls on demographic, personal history, and trauma variables

	No. (%) of displaced war survivors (n = 167)	No. (%) of controls (n = 167)	McNemar test
Sex:			
men	74 (44.3)	74 (44.3)	$P = 1.00^{**}$
women	93 (55.7)	93 (55.7)	
Marital status:			
married	64 (38.3)	64 (38.3)	$\chi^2 = 0.0, P = 1.0^*$
single/widowed/divorced	103 (61.7)	103 (61.7)	
Psychiatric illness before war	2 (1.2)	5 (3.0)	$P = 0.45^{**}$
Psychiatric illness in the family	46 (27.5)	41 (24.6)	$\chi^2 = 0.25, P = 0.61^*$
Posttraumatic stress disorder	38 (22.8)	1 (0.6)	$\chi^2 = 33.23, P < 0.001^*$
Depression	20 (12.0)	4 (2.4)	$P < 0.001^{**}$
Age (years, mean \pm standard deviation)	33.2 ± 12.4	33.0 ± 12.3	$t(166) = 1.2, P = 0.22^\ddagger$
Education level (mean \pm standard deviation)	2.2 ± 0.65	2.2 ± 0.65	$t(166) = -1.6, P = 0.10^\ddagger$
Income level (mean \pm standard deviation)	1.8 ± 0.75	2.1 ± 0.74	$t(166) = -3.6, P < 0.001^\ddagger$
Number of trauma events (mean \pm standard deviation)	13.1 ± 8.3	0.6 ± 0.8	$t(172) = 19.4, P < 0.001^\ddagger$

*McNemar test.

†Binomial distribution used.

‡t test.

greater number of war stressors that they had been exposed to. The two groups also differed in terms of income level. Compared to controls, displaced war survivors had significantly higher rates of PTSD (0.6% vs 22.8%, $P < 0.001$) and depression (2.4% vs 12.0%, $P < 0.001$). Among displaced war survivors, 51.6%, 43.5%, and 55.4% reported marked to severe negative impact on their family, social, and economic adaptation, respectively (Table 2). A comparison of the survivors and controls revealed marked differences in family, social, economic, occupational, and educational adaptation after trauma (all P 's significant at 0.001 level). The latter findings supported the validity of the scale.

Factors associated with psychiatric outcome

Two logistic regression analyses (method enter) were conducted using the diagnoses of current PTSD and current depression as the dependent variables. The ratio of cases to independent variables was 15.6 to 1, which was satisfactory. In the analysis where current PTSD was used as the dependent variable (Table 3), the overall prediction model was reliable ($\chi^2 = 72.0$, $df = 11$, $P < 0.001$) and explained 52.0% of the variance (Nagelkerke R^2). PTSD significantly related to old age, male sex, high levels of perceived distress during exposure to traumatic stressors, and negative

TABLE 2. Impact of trauma on different life spheres in displaced war survivors and matched controls

Impact of war on	No. (%) of		Displaced survivors (n = 167) (mean ± standard deviation)	Controls (n = 167) (mean ± standard deviation)	
	total participants				
Family adaptation:			2.6 ± 1.3	0.4 ± 0.9	t (166) = 18.5, $P < 0.001$
no impact	91	33.6			
mild to moderate impact	40	14.8			
marked to severe impact	140	51.6			
Social adaptation:			2.5 ± 1.3	0.5 ± 0.9	t (166) = 16.9, $P < 0.001$
no impact	90	33.2			
mild to moderate impact	63	23.3			
marked to severe impact	118	43.5			
Economic adaptation:			2.8 ± 1.3	0.8 ± 1.2	t (166) = 15.7, $P < 0.001$
no impact	77	28.4			
mild to moderate impact	44	16.2			
marked to severe impact	150	55.4			
Occupational adaptation:			0.8 ± 1.4	0.1 ± 0.6	t (165) = 5.5, $P < 0.001$
no impact	206	76			
mild to moderate impact	22	8.1			
marked to severe impact	43	15.9			
Educational adaptation:			0.7 ± 1.1	0.1 ± 0.4	t (166) = 7.4, $P < 0.001$
no impact	201	74.2			
mild to moderate impact	46	17.0			
marked to severe impact	24	8.8			

TABLE 3. Factors associated with posttraumatic stress disorder in displaced war survivors persons (n = 173)

Predictors	Standard		Wald test	Odds ratio (95% confidence interval)	P
	Beta	error			
Age	0.06	0.02	5.183	1.06 (1.01-1.11)	0.02
Sex (male)	-1.51	0.53	8.192	0.22 (0.08-0.62)	0.004
Marital status	-0.16	0.60	0.071	0.85 (0.26-2.77)	0.79
Education	-0.19	0.39	0.240	0.83 (0.38-1.78)	0.62
Level of income	-0.48	0.37	1.657	0.62 (0.30-1.29)	0.20
History of psychiatric illness	1.07	1.56	0.475	2.93 (0.14-62.34)	0.49
Family history of psychiatric illness	-1.01	0.67	2.259	0.37 (0.10-1.36)	0.13
Time since trauma	-0.001	0.01	0.005	0.99 (0.98-1.02)	0.94
Distress during trauma	1.15	0.46	6.145	3.15 (1.27-7.81)	0.013
Loss of control during trauma	-0.07	0.34	0.048	0.93 (0.48-1.80)	0.83
Global impact of trauma on life	0.18	0.08	4.400	1.19 (1.01-1.41)	0.04

TABLE 4. Factors associated with depression in displaced war survivors (n = 173)

Predictors	Beta	Standard error	Wald test	Odds ratio (95% confidence interval)	P
Age	0.03	0.03	1.344	1.03 (0.98-1.09)	0.25
Sex (male)	-0.86	0.61	1.962	0.42 (0.13-1.41)	0.16
Marital status	0.19	0.64	0.093	1.22 (0.35-4.28)	0.76
Education	-0.24	0.45	0.27	0.79 (0.32-1.92)	0.60
Level of income	-0.42	0.42	0.972	0.66 (0.29-1.51)	0.32
History of psychiatric illness	-0.07	1.44	0.002	0.93 (0.06-15.62)	0.96
Family history of psychiatric illness	-1.12	0.81	1.931	0.33 (0.07-1.59)	0.17
Time since trauma	0.01	0.01	0.787	1.01 (0.99-1.04)	0.38
Distress during trauma	1.22	0.50	6.227	3.41 (1.30-8.92)	0.013
Loss of control during trauma	-0.42	0.36	1.413	0.66 (0.33-1.32)	0.24
Global impact of trauma on life	-0.004	0.09	0.002	0.99 (0.84-1.19)	0.96

cumulative impact of trauma on life spheres. When the logistic regression analysis was repeated with current major depressive episode as the dependent variable (Table 4), the overall model was again significant ($\chi^2 = 32.6$, $df = 1$, 11 , $P < 0.001$) and explained 32.2% of the variance (Nagelkerke R^2). Only high levels of distress during exposure to war stressors related to current depression. Interestingly, there was no significant association between the impact of trauma on economic, social, family, occupational, and educational adaptation and depression.

DISCUSSION

Comparisons between displaced war survivors and control participants matched on age, sex, and education revealed significant differences in terms of impact of war stressors on family, occupational, social, economic, and educational adaptation. This finding supported our first hypothesis that refugees and internally displaced persons from war settings endure socioeconomic disadvantage, problems in family functioning, lack of access to educational and occupational opportunities, loss of social support, and perhaps marginalization and isolation. It needs to be noted, however, that it is not possible to attribute this negative impact on life domains to displacement only, because our study participants had also been exposed to multiple war-related traumatic events, which may have had independent effects on adaptation.

The strongest predictor of PTSD and depression in internally displaced persons and refugees was the level of perceived distress during exposure to war-related stressors. This finding supported our second study hypothesis and is consistent with previous studies which showed that perceived stressfulness of the stressors, rather than

mere exposure to them, determines traumatic stress in war and torture survivors (9,16,19,32,33). According to the "learning theory" of traumatic stress, distress, fear, or panic brought about by an inability to escape from the stressor, avoid the occurrence of the event, or reduce its impact, leads to a sense of helplessness with respect to possible future occurrences of the event, thereby causing anxiety reactions, including PTSD symptoms (34-36). In war settings, exposure to multiple stressor events may reinforce the learned helplessness effects of the primary traumatic event and may lead to more certain helplessness and anxiety reactions (34). Indeed, in studies involving survivors of war (9) and earthquake (37) anticipatory fear of recurrence of trauma events and associated helplessness feelings and cognitions in the aftermath of trauma exposure emerged as the strongest predictor of PTSD.

The significant association between distress during trauma and major depressive episode is consistent with previous similar association in war (9), torture (18), earthquake (37,38), and tsunami survivors (39). As PTSD and depression are highly comorbid conditions (46% of survivors with PTSD also had depression in this study), it is possible that they share the same mediating factors and depression develops secondary to PTSD (38,39). Such secondary depression might be related to the helplessness- and hopelessness-generating effects of trauma exposure and subsequent traumatic stress problems. Exposure to multiple war stressors may also lead to hopelessness and depression.

Global impact of trauma on adaptation in different life domains showed independent but rather weak association with PTSD and it was not associated with depression, a finding that provided only partial support for our third study hypothesis. The mechanisms by which such second-

ary effects of trauma have an influence on PTSD symptoms are not entirely clear. On the one hand, psychological trauma could be conceptualized as involving exposure to primary traumatic events which then lead to secondary stressors, all contributing to an ongoing traumatization process (5,15,40-42). For example, loss of economic opportunities and lower income may be a direct result of the primary trauma (eg, exposure to a war-related traumatic stressor) because the latter may be associated with loss of breadwinning family members, medical problems due to torture or serious physical injury, or sudden loss of property. They may thus act as constant reminders of the primary traumatic events and aggravate PTSD symptoms. It is also possible that such secondary stressors reinforce the helplessness effects of the primary trauma. On the other hand, given that PTSD symptoms lead to serious disability in social, occupational, and family functioning, the psychological reactions in response to secondary effects of trauma on life spheres might be the consequence rather than the cause of PTSD. Although attempt was made to rule out the latter effect in assessments, the retrospective nature of the study does not preclude such confounding.

The findings need to be interpreted in the light of some methodological limitations. Although the sample size afforded sufficient power for the analyses it was still small and findings from such small samples may have limited generalizability. However, the fact that the predictors were consistent with previous research supports the validity of the findings. Furthermore, although measures were taken to minimize sampling bias, non-random selection of study participants may have resulted in a sample with limited representativeness of displaced war survivors in Croatia. Finally, the retrospective design of the study, which involved recall of events and associated reactions assessed on average nine years post-trauma may have affected the internal validity.

In conclusion, our findings suggest that psychological processes during exposure to war stressors are the most important determinants of PTSD and depression in displaced war survivors. Although adaptation in different life spheres after trauma exposure may have an impact on mental health, its effect is not robust and consistent across disorders. These findings imply that a non-trauma-focused approach in rehabilitation of displaced war survivors is not justified. Trauma-focused cognitive-behavioral treatments and exposure-based interventions have been shown to be effective in a wide range of trauma survivors (43). Given the critical role played by distress, fear, and helplessness

responses in posttraumatic stress, restoration of sense of control over fear would be effective in recovery from traumatic stress in war survivors. Indeed, treatment studies have found that a behavioral intervention designed to enhance sense of control over fear is highly effective in reducing PTSD and depression in natural disaster survivors (38,44,45) and traumatized refugees (33,34,46). More studies are needed to test the efficacy of trauma-focused interventions in refugees and internally displaced persons.

Funding This study was supported by grants from the Spunk Fund Inc (New York) and the Bromley Trust (England).

Ethical approval was received from The Research Ethics Committee of the Institute of Psychiatry, King's College, University of London and Ethics Committee of Medical School, University of Rijeka.

Declaration of authorship ES, MLC, and MB developed the study concept and design. MB and TF performed acquisition of data. ES and MLC performed the analysis and interpretation of data. MLC and ES drafted the manuscript. ES, MLC, TF, and MB made the critical revision of the manuscript for important intellectual content. ES performed the statistical analysis. MB obtained funding.

Competing interests All authors have completed the Unified Competing Interest form at www.icmje.org/coi_disclosure.pdf (available on request from the corresponding author) and declare: no support from any organization for the submitted work; no financial relationships with any organizations that might have an interest in the submitted work in the previous 3 years; no other relationships or activities that could appear to have influenced the submitted work.

References

- 1 Steel Z, Chey T, Silove D, Marnane C, Bryant RA, van Ommeren M. Association of torture and other potentially traumatic events with mental health outcomes among populations exposed to mass conflict and displacement: a systematic review and meta-analysis. *JAMA*. 2009;302:537-49. [Medline:19654388](https://pubmed.ncbi.nlm.nih.gov/19654388/) doi:10.1001/jama.2009.1132
- 2 United Nations High Commissioner for Refugees. UNHCR Global Trends 2010. Available from: <http://www.unhcr.org/4dfa11499.html>. Accessed: November 28, 2011.
- 3 Porter M, Haslam N. Predisplacement and postdisplacement factors associated with mental health of refugees and internally displaced persons: a meta-analysis. *JAMA*. 2005;294:602-12. [Medline:16077055](https://pubmed.ncbi.nlm.nih.gov/16077055/) doi:10.1001/jama.294.5.602
- 4 Bhugra D, Ayonrinde O. Depression in migrants and ethnic minorities. *Adv Psychiatr Treat*. 2004;10:13-7. doi:10.1192/apt.10.1.13
- 5 Bhugra D, Becker MA. Migration, cultural bereavement and cultural identity. *World Psychiatry*. 2005;4:18-24. [Medline:16633496](https://pubmed.ncbi.nlm.nih.gov/16633496/)
- 6 Burnett A, Peel M. Health needs of asylum seekers and refugees. *BMJ*. 2001;322:544-7. [Medline:11230074](https://pubmed.ncbi.nlm.nih.gov/11230074/) doi:10.1136/bmj.322.7285.544
- 7 Miller KE, Rasmussen A. War exposure, daily stressors, and mental health in conflict and post-conflict settings: bridging the

- divide between trauma-focused and psychosocial frameworks. *Soc Sci Med.* 2010;70:7-16. [Medline:19854552](#) [doi:10.1016/j.socscimed.2009.09.029](#)
- 8 Miller KE, Omidian P, Rasmussen A, Yaqubi A, Daudzai H. Daily stressors, war experiences, and mental health in Afghanistan. *Transcult Psychiatry.* 2008;45:611-38. [Medline:19091728](#) [doi:10.1177/1363461508100785](#)
 - 9 Basoglu M, Livanou M, Crnobaric C, Franciskovic T, Suljic E, Duric D, et al. Psychiatric and cognitive effects of war in former Yugoslavia: association of lack of redress for trauma and posttraumatic stress reactions. *JAMA.* 2005;294:580-90. [Medline:16077052](#) [doi:10.1001/jama.294.5.580](#)
 - 10 Murray J, Ehlers A, Mayou RA. Dissociation and post-traumatic stress disorder: two prospective studies of road traffic accident survivors. *Br J Psychiatry.* 2002;180:363-8. [Medline:11925361](#) [doi:10.1192/bjp.180.4.363](#)
 - 11 Lanius RA, Vermetten E, Loewenstein RJ, Brand B, Schmahl C, Bremner JD, et al. Emotion modulation in PTSD: Clinical and neurobiological evidence for a dissociative subtype. *Am J Psychiatry.* 2010;167:640-7. [Medline:20360318](#) [doi:10.1176/appi.ajp.2009.09081168](#)
 - 12 Bryant RA. Predicting posttraumatic stress disorder from acute reactions. *J Trauma Dissociation.* 2005;6:5-15. [Medline:16150665](#) [doi:10.1300/J229v06n02_02](#)
 - 13 Gil S, Caspi Y. Personality traits, coping style, and perceived threat as predictors of posttraumatic stress disorder after exposure to a terrorist attack: a prospective study. *Psychosom Med.* 2006;68:904-9. [Medline:17079704](#) [doi:10.1097/01.psy.0000242124.21796.f8](#)
 - 14 Cukor J, Wyka K, Jayasinghe N, Weathers F, Giosan C, Leck P, et al. Prevalence and predictors of posttraumatic stress symptoms in utility workers deployed to the World Trade Center following the attacks of September 11, 2001. *Depress Anxiety.* 2011;28:210-7. [Medline:21394854](#) [doi:10.1002/da.20776](#)
 - 15 Basoglu M, Paker M, Ozmen E, Tasdemir O, Sahin D. Factors related to long-term traumatic stress responses in survivors of torture in Turkey. *JAMA.* 1994;272:357-63. [Medline:8028166](#) [doi:10.1001/jama.272.5.357](#)
 - 16 Başoğlu M, Paker M, Paker O, Ozmen E, Marks I, Incesu C, et al. Psychological effects of torture: a comparison of tortured with nontortured political activists in Turkey. *Am J Psychiatry.* 1994;151:76-81. [Medline:8267139](#)
 - 17 Basoglu M, Mineka S, Paker M, Aker T, Livanou M, Gok S. Psychological preparedness for trauma as a protective factor in survivors of torture. *Psychol Med.* 1997;27:1421-33. [Medline:9403913](#) [doi:10.1017/S0033291797005679](#)
 - 18 Basoglu M, Livanou M, Crnobaric C. Torture vs other cruel, inhuman, and degrading treatment: is the distinction real or apparent? *Arch Gen Psychiatry.* 2007;64:277-85. [Medline:17339516](#) [doi:10.1001/archpsyc.64.3.277](#)
 - 19 Basoglu M. A multivariate contextual analysis of torture and cruel, inhuman, and degrading treatments: implications for an evidence-based definition of torture. *Am J Orthopsychiatry.* 2009;79:135-45. [Medline:19485630](#) [doi:10.1037/a0015681](#)
 - 20 Mollica RF, McInnes K, Sarajlic N, Lavelle J, Sarajlic I, Massagli MP. Disability associated with psychiatric comorbidity and health status in Bosnian refugees living in Croatia. *JAMA.* 1999;282:433-9. [Medline:10442658](#) [doi:10.1001/jama.282.5.433](#)
 - 21 Scholte WF, Olff M, Ventevogel P, de Vries GJ, Jansveld E, Cardozo BL, et al. Mental health symptoms following war and repression in eastern Afghanistan. *JAMA.* 2004;292:585-93. [Medline:15292084](#) [doi:10.1001/jama.292.5.585](#)
 - 22 de Jong JT, Komproe IH, Van Ommeren M, El Masri M, Araya M, Khaled N, et al. Lifetime events and posttraumatic stress disorder in 4 postconflict settings. *JAMA.* 2001;286:555-62. [Medline:11476657](#) [doi:10.1001/jama.286.5.555](#)
 - 23 Gibney M, Cornett L, Wood R. Political Terror Scale 1976-2008. Available from: <http://www.politicalerrorscale.org/>. Accessed: November 25, 2011.
 - 24 United Nations High Commissioner for Refugees. Refugee protection and humanitarian work in Croatia: An overview of UNHCR's operations in the past 18 years. Available from: <http://www.unhcr.hr/eng/>. Accessed: November 25, 2011.
 - 25 Kostović I, Henigsberg N, Judaš M. The role of Croatia in the management of the humanitarian crisis in Bosnia and Herzegovina. *National Security and the Future.* 2000;1:133-45.
 - 26 Watters JK, Biernacki P. Targeted sampling: options for the study of hidden populations. *Soc Probl.* 1989;36:416-30. [doi:10.1525/sp.1989.36.4.03a00070](#)
 - 27 Frank O, Snijders T. Estimating the size of hidden populations using snowball sampling. *J Off Stat.* 1994;10:53-67.
 - 28 First MB, Gibbon M, Spitzer RL, Williams JBW. User's guide for the Structured Clinical Interview for DSM-IV Axis I Disorders, research version 2.0. New York (NY): Biometrics Research Department, New York State Psychiatric Institute; 1996.
 - 29 Blake DD, Weathers FW, Nagy LM, Kaloupek DG, Charney DS, Keane T. Clinician-Administered PTSD Scale (CAPS): Current and Lifetime Diagnostic Version. Boston (MA): Behavioral Science Division, National Center for Posttraumatic Stress Disorder; 1990.
 - 30 Basoglu M, Paker M, Paker O, Ozmen E, Marks I, Incesu C, et al. Psychological effects of torture: a comparison of tortured with nontortured political activists in Turkey. *Am J Psychiatry.* 1994;151:76-81. [Medline:8267139](#)
 - 31 Agresti A. *Categorical Data Analysis.* New York: Wiley-Interscience; 2002.
 - 32 Basoglu M, Paker M. Severity of trauma as predictor of long-term psychological status in survivors of torture. *J Anxiety Disord.* 1995;9:339-50. [doi:10.1016/0887-6185\(95\)00014-F](#)
 - 33 Salcioglu E, Basoglu M. Control-focused behavioural treatment of female war survivors with torture and gang rape experience: four

- case studies. *Eur J Psychotraumatol*. 2011;2 Suppl 1:S192.
- 34 Basoglu M, Salcioglu E. A mental healthcare model for mass trauma survivors: Control-focused behavioral treatment of earthquake, war, and torture trauma. Cambridge (UK): Cambridge University Press; 2011.
- 35 Alloy L, Kelly K, Mineka S, Clements C. Comorbidity in anxiety and depressive disorders: a helplessness/hopelessness perspective. In: Maser JD, Cloninger CR, editors. *Comorbidity in anxiety and mood disorders*. Washington (DC): American Psychiatric Press; 1990. p. 499-543.
- 36 Mineka S, Zinbarg R. A contemporary learning theory perspective on the etiology of anxiety disorders: it's not what you thought it was. *Am Psychol*. 2006;61:10-26. [Medline:16435973](#) [doi:10.1037/0003-066X.61.1.10](#)
- 37 Basoglu M, Salcioglu E, Livanou M. Traumatic stress responses in earthquake survivors in Turkey. *J Trauma Stress*. 2002;15:269-76. [Medline:12224798](#) [doi:10.1023/A:1016241826589](#)
- 38 Salcioglu E, Basoglu M, Livanou M. Effects of live exposure on symptoms of posttraumatic stress disorder: The role of reduced behavioral avoidance in improvement. *Behav Res Ther*. 2007;45:2268-79. [Medline:17570342](#) [doi:10.1016/j.brat.2007.04.012](#)
- 39 Rosendal S, Salcioglu E, Andersen HS, Mortensen EL. Exposure characteristics and peri-trauma emotional reactions during the 2004 tsunami in Southeast Asia—what predicts posttraumatic stress and depressive symptoms? *Compr Psychiatry*. 2011;52:630-7. [Medline:21349509](#) [doi:10.1016/j.comppsy.2010.12.004](#)
- 40 Steel Z, Chey T, Silove D, Marnane C, Bryant RA, van Ommeren M. Association of torture and other potentially traumatic events with mental health outcomes among populations exposed to mass conflict and displacement: a systematic review and meta-analysis. *JAMA*. 2009;302:537-49. [Medline:19654388](#) [doi:10.1001/jama.2009.1132](#)
- 41 Hollifield M, Warner TD, Lian N, Krakow B, Jenkins JH, Kesler J, et al. Measuring trauma and health status in refugees: a critical review. *JAMA*. 2002;288:611-21. [Medline:12150673](#) [doi:10.1001/jama.288.5.611](#)
- 42 Porter M, Haslam N. Forced displacement in Yugoslavia: a meta-analysis of psychological consequences and their moderators. *J Trauma Stress*. 2001;14:817-34. [Medline:11776427](#) [doi:10.1023/A:1013054524810](#)
- 43 Bradley R, Greene J, Russ E, Dutra L, Westen D. A multidimensional meta-analysis of psychotherapy for PTSD. *Am J Psychiatry*. 2005;162:214-27. [Medline:15677582](#) [doi:10.1176/appi.ajp.162.2.214](#)
- 44 Basoglu M, Livanou M, Salcioglu, Kalender D. A brief behavioral treatment of chronic posttraumatic stress disorder in earthquake survivors: Results from an open clinical trial. *Psychol Med*. 2003;33:647-54. [Medline:12785466](#) [doi:10.1017/S0033291703007360](#)
- 45 Basoglu M, Salcioglu E, Livanou M, Kalender D, Acar G. Single-session behavioral treatment of earthquake-related posttraumatic stress disorder: A randomized waiting list controlled trial. *J Trauma Stress*. 2005;18:1-11. [Medline:16281190](#) [doi:10.1002/jts.20011](#)
- 46 Basoglu M, Ekblad S, Baarnhielm S, Livanou M. Cognitive-behavioral treatment of tortured asylum seekers: A case study. *J Anxiety Disord*. 2004;18:357-69. [Medline:15125982](#) [doi:10.1016/S0887-6185\(02\)00248-7](#)