

Stroke prevention in atrial fibrillation and 'real world' adherence to guidelines in the Balkan Region: The BALKAN-AF Survey

(BALKAN-AF Investigators) Potpara, Tatjana S.; ...; Brusich, Sandro; Lip, Gregory Y. H.

Source / Izvornik: **Scientific Reports, 2016, 6, 1 - 10**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.1038/srep20432>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:184:477266>

Rights / Prava: [Attribution 4.0 International](#)/[Imenovanje 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2025-01-15**



Repository / Repozitorij:

[Repository of the University of Rijeka, Faculty of Medicine - FMRI Repository](#)



SCIENTIFIC REPORTS



OPEN

Stroke prevention in atrial fibrillation and 'real world' adherence to guidelines in the Balkan Region: The BALKAN-AF Survey

Received: 15 September 2015

Accepted: 04 January 2016

Published: 12 February 2016

Tatjana S. Potpara^{1,2}, Gheorghe-Andrei Dan³, Elina Trendafilova⁴, Artan Goda⁵, Zumreta Kusljagic⁶, Sime Manola⁷, Ljilja Music⁸, Rodica Musetescu⁹, Elisabeta Badila¹⁰, Gorana Mitic¹¹, Vilma Papparisto⁵, Elena S. Dimitrova⁴, Marija M. Polovina^{1,2}, Stanislav L. Petranov¹², Hortensia Djergo⁵, Daniela Loncar⁶, Amira Bijedic⁶, Sandro Brusich¹³, Gregory Y. H. Lip^{1,14} & The BALKAN-AF Investigators[#]

Data on the management of atrial fibrillation (AF) in the Balkan Region are limited. The Serbian AF Association (SAFA) prospectively investigated contemporary 'real-world' AF management in clinical practice in Albania, Bosnia&Herzegovina, Bulgaria, Croatia, Montenegro, Romania and Serbia through a 14-week (December 2014-February 2015) prospective, multicentre survey of consecutive AF patients. We report the results pertinent to stroke prevention strategies. Of 2712 enrolled patients, 2663 (98.2%) with complete data were included in this analysis (mean age 69.1 ± 10.9 years, female 44.6%). Overall, 1960 patients (73.6%) received oral anticoagulants (OAC) and 762 (28.6%) received antiplatelet drugs. Of patients given OAC, 17.2% received non-vitamin K antagonist oral anticoagulants (NOACs). CHA_2DS_2-VASc score was not significantly associated with OAC use. Of the 'truly low-risk' patients ($CHA_2DS_2-VASc = 0$ [males], or 1 [females]) 56.5% received OAC. Time in Therapeutic Range (TTR) was available in only 18.7% of patients (mean TTR: $49.5\% \pm 22.3\%$). Age ≥ 80 years, prior myocardial infarction and paroxysmal AF were independent predictors of OAC non-use. Our survey shows a relatively high overall use of OAC in AF patients, but with low quality of vitamin K antagonist therapy and insufficient adherence to AF guidelines. Additional efforts are needed to improve AF-related thromboprophylaxis in clinical practice in the Balkan Region.

In parallel with increasing global burden of atrial fibrillation (AF), accumulating high-quality evidence from randomized clinical trials on AF management inform frequent updates of AF guidelines¹. However, guideline implementation into daily clinical practice might be incomplete for many reasons and monitoring of routine practice through ongoing large, well-conducted long-term registries^{2,3} helps to understand and attenuate barriers for evidence-based management of AF in 'real-world' setting.

¹School of Medicine, Belgrade University, Belgrade, Serbia. ²Cardiology Clinic, Clinical Centre of Serbia, Belgrade, Serbia. ³Medicine University "Carol Davila", Colentina University Hospital, Bucharest, Romania. ⁴National Heart Hospital, Sofia, Bulgaria. ⁵Clinic of Cardiology, University Hospital centre Mother Theresa, Tirana, Albania. ⁶Cardiology Department, Clinic for internal diseases, Tuzla, Bosnia & Herzegovina. ⁷Clinical Hospital Centre Sestre Milosrdnice, Zagreb, Croatia. ⁸University Clinical Centre Podgorica, Podgorica, Montenegro. ⁹Cardiology Centre, County Emergency Hospital, Craiova, Romania. ¹⁰Medicine University "Carol Davila", Emergency Clinical Hospital, Internal Medicine Department, Bucharest, Romania. ¹¹Clinical Centre Vojvodina, Novi Sad, Serbia. ¹²Multiprofile Hospital for Active Treatment, Bourgas, Bulgaria. ¹³Clinical Hospital Centre Rijeka, Rijeka, Croatia. ¹⁴University of Birmingham Centre for Cardiovascular Sciences, City Hospital, Birmingham B18 7QH, UK. [#]A comprehensive list of authors and affiliations appear at the end of the paper. Correspondence and requests for materials should be addressed to T.P. (email: tanjapotpara@gmail.com or tatjana.potpara@mfub.bg.ac.rs)

Recent reports from contemporary European AF registries have provided important insights into AF management in clinical practice^{2–5}, including the observation of certain regional differences in management across Europe⁶. Most countries from the Balkan Region (comprising an area of >50 million inhabitants) were not participating in prior registries, and in contrast to other European regions, ‘real-world’ data on the management of AF in Balkan countries are limited.

In this study, we investigated contemporary real-world patterns of AF management in the Balkan Region through a prospective 14-week survey of consecutive AF patients in clinical practice, and we report the results pertinent to stroke prevention.

Methods

Study design and patient selection. A detailed report on the Balkan-AF study protocol has been published⁷. A 14-week prospective, multicentre ‘snapshot’ Balkan-AF survey of consecutive patients with electrocardiographically documented AF, who were seen by cardiologists or internal medicine specialists (in centres where a cardiologist was not available), was conducted from December 2014 to February 2015 in Albania, Bosnia & Herzegovina, Bulgaria, Croatia, Montenegro, Romania and Serbia (a total of ~40 million inhabitants). The survey was designed and conducted by the Serbian Atrial Fibrillation Association (SAFA), which is a non-profit multidisciplinary association of expert physicians involved in AF management and AF research.

The survey was announced to the National Cardiology Societies and relevant Working Groups or associations in Albania, Bosnia & Herzegovina, Bulgaria, Croatia, Former Yugoslav Republic Macedonia, Montenegro, Romania, Slovenia and Serbia. In the participating countries Balkan-AF survey was approved by the national and/or local Institutional Review Board, or the need for approval was waived according to the regulations in the respective country. In concordance with the local policy, a signed patient informed consent was obtained from each patient before enrolment. The study protocol conforms to the ethical guidelines of the 1975 Declaration of Helsinki as reflected in a priori approval by the institution’s human research committee.

Each country participated with university and non-university hospitals and outpatient health centres in- and outside the capital cities. Patients younger than 18 years and patients with prosthetic mechanical heart valves or significant valve disease requiring surgical repair were not included.

Data collection. Data were collected via a web-based electronic case report form (CRF) with a range of pre-specified plausibility checks for the entries. The CRF was formulated to obtain the information on patients’ characteristics including demographics, cardiovascular risk factors, medical history, AF-related data regarding symptoms, prior history of AF, AF clinical type, prior use of antithrombotic medication, antiarrhythmic drugs or other therapies, health care setting (i.e., university/non-university health centre, in- or outside the capital city, in-hospital or outpatient, internal medicine specialist/cardiologist, main reason for current visit/hospitalization, emergency or non-emergency setting, length of hospitalization, etc.) and patient’s presentation, AF management at enrolling visit or hospitalization (i.e., medication, cardioversion, AF ablation) and further management strategy post discharge, and diagnostic procedures performed due to AF during enrolling visit/hospitalization or within the last 12 months (the latter was not applicable to patients with first-diagnosed AF). A detailed list of cardiovascular risk factors, diseases and risk scores definitions used in the Balkan-AF survey is provided in the Supplementary Appendix 1.

Systematic monitoring of centres was not performed due to the relatively short duration of the survey. The national coordinators and all investigators are the guarantors of the consecutiveness of enrolment, accuracy and completeness of data. The CRF, patient files, and medical records (paper or database) serve as source documents.

Statistical analysis. Following a test of statistical normality, continuous variables were presented as mean with standard deviation (SD), or with a skewed distribution as median with interquartile range (IQR, 25th–75th quartile). Categorical variables were reported as counts with percentages. The Student t-test was used for comparison of continuous variables with normal distribution, and Mann-Whitney test for continuous variables with skewed distribution. Differences in categorical variables were tested by Chi-square test.

Univariate and multivariable logistic regression analyses were used to investigate the associations of variables shown in Table 1 (that is, demographic data, patient clinical characteristics and AF characteristics) and health care setting with the use of oral anticoagulants (OAC) and other antithrombotic therapies (that is, antiplatelet drugs), as prescribed at discharge from enrolling visit or hospitalization. Variables statistically significant on univariate analysis were entered into the multivariable model to identify independent predictors of OAC use. All analyses were adjusted for country code, to account for differences in the health care systems among the participating countries.

Because the main reason for enrolling visit or hospitalization could have been either AF or some other condition, we have performed two sensitivity analyses. First, we excluded patients seen for other reasons (in whom the use of OAC might have been influenced by other condition) and performed the analysis of OAC use as in the main cohort. Second, we excluded patients presenting with an acute coronary syndrome and then performed the OAC use analyses in the rest of the main cohort.

All results are reported as Odds Ratio (OR) with 95% Confidence Interval (CI). All statistical analyses were performed using SPSS 20.0 software package (SPSS Inc., Chicago, Illinois). A two-sided P value of <0.05 was considered statistically significant.

Results

A total of 2712 patients were enrolled in 49 centres from seven Balkan countries; 27 centres (55.1%) were university hospitals enrolling 2161 patients (86.6%). Eighteen centres (36.7%) were situated in the capital cities and

	Total	Albania	B&H	Bulgaria	Croatia	Montenegro	Romania	Serbia
	n = 2663	n = 313	n = 265	n = 443	n = 159	n = 102	n = 699	n = 682
<i>Demographic</i>								
Age, years (mean ± SD)	69.1 ± 10.9	68.2 ± 10.2	69.2 ± 10.6	70.2 ± 10.7	69.6 ± 11.1	65.0 ± 10.8	70.9 ± 10.8	67.4 ± 11.0
Age ≥ 65–74 years (%)	878 (33.0)	114 (36.4)	99 (37.4)	139 (31.4)	46 (28.9)	39 (38.2)	215 (30.8)	226 (33.1)
Age ≥ 75 years (%)	942 (35.4)	94 (30.0)	87 (32.8)	176 (39.7)	62 (39.0)	20 (19.6)	301 (43.1)	202 (29.6)
Age ≥ 80 years (%)	418 (17.7)	39 (12.5)	40 (15.1)	76 (17.2)	31 (19.5)	6 (5.9)	143 (20.5)	83 (12.2)
Female sex (%)	1188 (44.6)	151 (48.2)	115 (43.4)	189 (42.7)	67 (42.1)	35 (34.3)	325 (46.5)	306 (44.9)
Cigarette smoking ever (%)	776 (29.1)	111 (35.5)	66 (24.9)	102 (23.0)	48 (30.2)	38 (37.3)	154 (22.0)	257 (37.7)
Cigarette smoking current (%)	339 (12.7)	74 (23.6)	32 (12.1)	48 (10.8)	29 (18.2)	22 (21.6)	58 (8.3)	76 (11.1)
Alcohol abuse (%)	110 (4.1)	30 (9.6)	9 (3.4)	35 (7.9)	4 (2.5)	0 (0.0)	20 (2.9)	12 (1.8)
Body mass index (mean ± SD)	27.7 ± 4.4	28.0 ± 3.9	26.8 ± 4.3	27.9 ± 3.8	27.8 ± 3.4	27.3 ± 3.7	28.5 ± 5.1	27.2 ± 4.3
Systolic BP, mmHg (mean ± SD)	134.6 ± 22.0	131.5 ± 25.0	133.3 ± 24.6	135.2 ± 19.0	137.5 ± 19.3	137.7 ± 19.6	136.2 ± 24.0	133.4 ± 19.8
Diastolic BP, mmHg (mean ± SD)	81.0 ± 12.2	81.4 ± 13.7	81.5 ± 12.7	81.6 ± 11.1	83.6 ± 10.0	86.8 ± 10.7	78.8 ± 13.1	81.0 ± 11.4
<i>Characteristics of AF</i>								
First diagnosed AF (%)	626 (23.5)	107 (34.4)	92 (34.7)	109 (24.6)	21 (13.2)	22 (21.6)	146 (20.9)	129 (18.9)
<i>Known history of AF</i>								
Paroxysmal (%)	554 (27.2)	43 (21.0)	23 (13.3)	77 (23.1)	45 (32.6)	31 (38.8)	106 (19.2)	229 (41.4)
Persistent (%)	319 (15.7)	13 (6.3)	12 (6.9)	72 (21.6)	46 (33.3)	13 (16.2)	77 (13.9)	86 (15.6)
Long-standing persistent (%)	64 (3.1)	7 (3.4)	12 (6.9)	7 (2.1)	6 (4.3)	1 (1.2)	9 (1.6)	22 (4.0)
Permanent (%)	1081 (53.1)	142 (69.3)	126 (72.8)	178 (53.2)	41 (29.7)	34 (42.5)	360 (65.1)	200 (36.2)
<i>Clinical parameters</i>								
Arterial hypertension (%)	2108 (79.2)	218 (69.6)	210 (79.2)	398 (89.8)	127 (79.9)	75 (73.5)	518 (74.1)	562 (82.4)
Heart failure ever (%)	1157 (43.5)	137 (43.8)	129 (48.9)	230 (51.9)	52 (32.7)	5 (4.9)	465 (66.6)	163 (23.9)
Signs of heart failure at present (%)	1104 (41.5)	113 (36.1)	129 (48.9)	223 (50.3)	34 (21.4)	3 (2.9)	454 (65.0)	124 (18.9)
Coronary artery disease (%)	816 (30.7)	99 (31.6)	106 (40.0)	147 (33.2)	35 (22.0)	16 (15.7)	256 (36.7)	157 (23.1)
Prior PCI/stenting (%)	224 (8.4)	35 (11.2)	6 (2.3)	62 (14.0)	9 (5.7)	8 (7.8)	50 (7.2)	54 (7.9)
Prior CABG (%)	97 (3.6)	12 (3.8)	13 (4.9)	17 (3.8)	3 (1.9)	1 (1.0)	15 (2.1)	36 (5.3)
Prior myocardial infarction (%)	365 (13.7)	50 (16.0)	53 (20.0)	40 (9.0)	13 (8.2)	8 (7.8)	115 (16.6)	86 (12.6)
Stable coronary artery disease (%)	592 (22.0)	64 (20.4)	100 (37.7)	85 (19.2)	26 (16.4)	8 (7.8)	206 (29.5)	103 (15.1)
Valvular disease (%)	933 (35.0)	69 (22.0)	60 (22.6)	159 (35.9)	39 (24.5)	3 (2.9)	414 (59.2)	189 (27.7)
Mitral valve disease (%)	844 (31.7)	53 (16.9)	54 (20.4)	131 (29.6)	34 (21.4)	1 (1.0)	388 (55.5)	183 (26.8)
Mitral valve regurgitation (%)	818 (30.7)	53 (16.9)	51 (19.2)	127 (28.7)	33 (20.8)	1 (1.0)	381 (54.5)	172 (25.2)
Aortic valve disease (%)	299 (11.2)	28 (8.9)	23 (8.7)	66 (14.9)	8 (5.0)	2 (2.0)	131 (18.7)	41 (6.0)
Dilated cardiomyopathy (%)	216 (8.1)	21 (6.7)	14 (5.3)	8 (1.8)	18 (11.3)	1 (1.0)	100 (14.3)	54 (7.9)
Hypertrophic cardiomyopathy (%)	52 (2.0)	6 (1.9)	12 (4.5)	2 (0.5)	4 (2.5)	0 (0.0)	20 (2.9)	8 (1.2)
Restrictive cardiomyopathy (%)	4 (0.2)	0 (0.0)	0 (0.0)	0 (0.0)	1 (0.6)	0 (0.0)	3 (0.4)	0 (0.0)
Congenital heart disease (%)	7 (0.3)	0 (0.0)	0 (0.0)	1 (0.2)	0 (0.0)	0 (0.0)	2 (0.3)	4 (0.6)
Other cardiac disease (%)	206 (7.7)	11 (3.5)	8 (3.0)	9 (2.0)	4 (2.5)	0 (0.0)	92 (13.2)	82 (12.0)
Peripheral arterial disease (%)	122 (4.6)	13 (4.2)	7 (2.7)	19 (4.3)	5 (3.1)	2 (2.0)	40 (5.7)	36 (5.3)
Diabetes mellitus (%)	666 (25.0)	99 (31.6)	76 (28.7)	109 (24.6)	30 (18.9)	19 (18.6)	178 (25.5)	155 (22.7)
Chronic kidney disease (%)	411 (15.5)	36 (11.5)	33 (12.5)	78 (17.6)	21 (13.3)	1 (1.0)	164 (23.5)	78 (11.5)
Chronic hepatic disease (%)	96 (3.6)	7 (2.2)	9 (3.4)	13 (2.9)	2 (1.3)	0 (0.0)	54 (7.7)	11 (1.6)
Prior stroke (%)	280 (10.5)	29 (9.3)	40 (15.1)	47 (10.6)	13 (8.2)	7 (6.9)	65 (9.3)	79 (11.6)
Prior TIA (%)	83 (3.1)	16 (5.1)	18 (6.8)	14 (3.2)	5 (3.1)	0 (0.0)	14 (2.0)	16 (2.3)
Prior bleeding (%)	135 (5.0)	17 (5.4)	18 (6.8)	19 (4.3)	3 (1.9)	0 (0.0)	40 (5.7)	35 (5.0)
CHA ₂ DS ₂ -VASc score	3.48 ± 1.78	3.37 ± 1.79	3.66 ± 1.78	3.71 ± 1.76	3.21 ± 1.89	2.54 ± 1.61	3.81 ± 1.64	3.20 ± 1.76
HASBLED score	1.97 ± 1.23	1.87 ± 1.28	1.91 ± 1.18	1.92 ± 1.12	1.77 ± 1.17	1.87 ± 1.22	2.25 ± 1.26	1.86 ± 1.23

Table 1. Patient demographic data, clinical characteristics and AF characteristics. B&H: Bosnia & Herzegovina; SD: standard deviation; BP: blood pressure; AF: atrial fibrillation; PCI: percutaneous coronary intervention; CABG: coronary artery bypass grafting; TIA: transient ischemic attack.

enrolled 1241 patients (45.8%). A total of 2147 patients (79.2%) were enrolled by cardiologists, and 717 patients (26.8%) were seen in outpatient setting. Full data on antithrombotic therapy prescribed at current visit/hospitalization were available in 2663 patients (98.2%) and those patients were included in this analysis.

Demographic data (mean age 69.1 ± 10.9 years, range 18–96; female 44.6%), clinical characteristics of the study population and AF characteristics are shown in Table 1.




Figure 1. Stroke and bleeding risk. ‘Truly low-risk’: CHA₂DS₂-VASc = 0 in males, or CHA₂DS₂-VASc = 1 in females; B&H: Bosnia & Herzegovina.

	Total	Albania	B&H	Bulgaria	Croatia	Montenegro	Romania	Serbia
	n = 2663	n = 313	n = 265	n = 443	n = 159	n = 102	n = 699	n = 682
No antithrombotic therapy (%)	264 (9.9)	23 (7.3)	28 (10.6)	46 (10.4)	15 (9.4)	11 (10.8)	67 (9.6)	74 (10.9)
Oral anticoagulant therapy-overall (%)	1960 (73.6)	229 (73.2)	133 (50.2)	319 (72.0)	134 (84.3)	72 (70.6)	534 (76.4)	539 (79.0)
VKAs (%)	1662 (60.9)	205 (65.5)	111 (41.9)	191 (43.1)	110 (69.2)	68 (66.7)	482 (69.0)	455 (66.7)
NOACs (%)	338 (12.7)	24 (7.7)	22 (8.3)	128 (28.9)	24 (15.1)	4 (3.9)	52 (7.4)	84 (12.3)
Oral anticoagulant therapy alone (%)	1637 (61.5)	162 (51.8)	124 (46.8)	286 (64.6)	121 (76.1)	50 (49.0)	446 (63.8)	448 (65.7)
Antiplatelet therapy (%)	762 (28.6)	128 (40.9)	113 (42.6)	111 (25.1)	23 (14.5)	41 (40.2)	186 (26.6)	160 (23.5)
Single antiplatelet drug only (%)	320 (12.0)	39 (12.5)	88 (33.2)	54 (12.2)	9 (5.7)	14 (13.7)	61 (8.7)	55 (8.1)
DAPT only (%)	119 (4.5)	22 (7.0)	16 (6.0)	24 (5.4)	1 (0.6)	5 (4.9)	37 (5.3)	14 (2.1)
Dual therapy ^a	240 (9.0)	51 (16.3)	9 (3.4)	17 (3.8)	9 (5.7)	14 (13.7)	74 (10.6)	66 (9.7)
Triple therapy ^b	83 (3.1)	16 (5.1)	0 (0.0)	16 (3.6)	4 (2.5)	8 (7.8)	14 (2.0)	25 (3.7)

Table 2. Country-specific distribution of OAC and antiplatelet therapies. B&H: Bosnia & Herzegovina; OAC: oral anticoagulant; VKA: vitamin-K antagonist; NOAC: non-vitamin K antagonist; DAPT: dual antiplatelet therapy. ^aDual therapy: OAC plus single antiplatelet agent. ^bTriple therapy: OAC plus dual antiplatelet therapy.

Stroke and bleeding risk profile. Mean CHA₂DS₂-VASc score was 3.48 ± 1.77 (range 0–9, median 3.0, IQR 2.0–5.0), and a score of ≥2 was present in 2290 patients (86.0%). The mean CHADS₂ score was 2.15 ± 1.29 (≥2 in 65.6% of patients), and mean HASBLED score was 1.97 ± 1.23 (range 0–6). Country-specific stroke and bleeding risk distribution is shown in Table 1, Fig. 1 and Supplemental Table 1.

Antithrombotic therapies. Country-specific distribution of antithrombotic therapies is shown in Table 2. Overall, 264 patients (9.9%) were not given any antithrombotic therapy, 1960 patients (73.6%) were prescribed OAC, and 762 patients (28.6%) received an antiplatelet drug. OAC as only antithrombotic drug was given to 1637 patients (61.5%), whilst an antiplatelet drug only was given to 320 patients (12.0%, and in 91.2% of patients that was aspirin).

NOACs (i.e., dabigatran, rivaroxaban or apixaban) were given to 338 patients (12.7% of the whole study population or 17.2% of patients receiving OAC).

One patient previously underwent a left atrial appendage closure device implantation.




Figure 2. The use of antithrombotic therapies by CHA₂DS₂-VASc and HASBLED risk strata. APLT: antiplatelet therapy; DAPT: dual antiplatelet therapy; OAC: oral anticoagulant.

Adherence to guidelines. The proportions of OAC and other antithrombotic therapies by CHA₂DS₂-VASc and HASBLED score strata are shown in Fig. 2.

In the ‘truly’ low-risk group (that is, CHA₂DS₂-VASc score of 0 in males, or 1 in females) only 44 patients (33.6%) were not given any antithrombotic therapy, whilst 74 patients (56.5%) received OAC and 18 patients (13.7%) received an antiplatelet therapy, alone or in combination with OAC (3.8%). The use of OAC in ‘truly’ low-risk patients could be attributed to planned cardioversion or AF catheter ablation in only 21 patients (16.0%).

Of 2290 patients with CHA₂DS₂-VASc ≥ 2 , 194 (8.5%) received no antithrombotic therapy, 1401 (61.2%) were given OAC only and 393 patients (17.2%) received an antiplatelet drug alone. A combination of OAC and an antiplatelet drug was given to 302 patients (13.2%).

There was no significant association between the CHA₂DS₂-VASc score and OAC prescription on multivariate analysis (Table 3). The non-relationship to CHADS₂ score was similar (OR 1.06; 95% CI, 0.98–1.13, $p = 0.119$). The use of OAC increased with increasing HASBLED score, but the difference was significant only on univariate analysis ($p = 0.042$), Table 3.

The use of antiplatelet drugs was significantly associated with CHA₂DS₂-VASc score only on univariate analysis (Table 3).

Determinants of OAC use. Independent predictors of the use of antithrombotic therapies are shown in Table 3, including significant univariate associations with the use of OAC or antiplatelet drug only (full univariate analyses list is shown in Supplemental Table 2).

OAC monotherapy. On univariate analysis, most of the CHA₂DS₂-VASc score components were not significantly associated with OAC use, including prior stroke (OR 1.02; 0.76–1.37; $p = 0.878$). Patients with hypertension were more likely to use OAC, whilst older age (≥ 80 years) and coronary artery disease (CAD) were inversely associated with OAC use (Supplemental Table 2).

Increasing body mass index (BMI), mitral valve disease, dilated cardiomyopathy and thyroid disease were associated with increased use of OAC, whilst patients with chronic kidney disease (CKD) on dialysis and patients with chronic obstructive pulmonary disease (COPD) were less likely to use OAC (Supplemental Table 2).

The proportions of OAC use according to AF clinical type are shown in Fig. 3. Known history of AF was associated with increased use of OAC, whilst patients with paroxysmal AF were less likely to receive OAC. Patients treated in the health centres situated in the capital city or in university centres and patients managed by a cardiologist were more likely to use OAC compared to other patients (Supplemental Table 2).

Independent predictors of OAC use were hypertension, mitral valve disease, dilated cardiomyopathy, known history of AF and treatment in the capital city health centres, whilst age ≥ 80 years, prior MI and paroxysmal AF were independent predictors of OAC non-use (Table 3).

Antiplatelet drug (aspirin) monotherapy. Independent predictors of aspirin monotherapy were age ≥ 80 years, CAD, aortic valve disease, COPD and paroxysmal AF, whilst BMI, known history of AF and treatment in the capital city health centre or treatment by a cardiologist were negatively associated with aspirin use (Table 3).

Antithrombotic therapy	Univariate analysis (significant variables only)			Multivariate analysis			Antithrombotic therapy	Multivariate analysis		
	OR	95%CI	P	OR	95%CI	P		DAPT	OR	95%CI
OAC only										
HASBLED (cont. variable)	1.08	1.01–1.16	0.042				PCI	4.47	2.69–7.43	<0.001
Age ≥80 years	0.62	0.50–0.78	<0.001	0.54	0.37–0.79	0.002	Coronary artery disease (any)	9.67	5.81–16.10	<0.001
Hypertension	1.67	1.36–2.05	<0.001	1.85	1.30–2.63	0.001	COPD	1.95	1.14–3.33	0.015
Prior MI	0.47	0.36–0.61	<0.001	0.58	0.37–0.91	0.018	Known history of AF	0.45	0.28–0.72	0.001
Prior PCI	0.32	0.22–0.47	<0.001				Paroxysmal AF	2.31	1.45–3.69	<0.001
Coronary artery disease (any)	0.49	0.41–0.60	<0.001							
Stable coronary artery disease	0.55	0.46–0.67	<0.001							
Mitral valve disease	1.40	1.15–1.70	0.001	1.56	1.07–2.28	0.021				
Dilated cardiomyopathy	2.06	1.40–3.04	<0.001	1.72	1.10–2.68	0.018	Dual or triple therapy	OR	95%CI	P
CKD on dialysis	0.16	0.04–0.60	0.007				Age ≥80 years	0.58	0.39–0.88	0.009
COPD	0.73	0.57–0.94	0.016				PCI	3.69	2.46–5.46	<0.001
Thyroid disease	1.54	1.13–2.11	0.007				Coronary artery disease (any)	2.78	1.94–4.00	<0.001
Known history of AF	2.53	2.07–3.08	<0.001	1.51	1.04–2.20	0.032	Hospital-based centre	2.99	1.51–5.91	0.002
Paroxysmal AF	0.38	0.32–0.46	<0.001	0.44	0.32–0.62	<0.001	HASBLED (cont. variable)	1.24	1.11–1.38	<0.001
Body mass index	1.06	1.04–1.08	<0.001	1.04	1.01–1.08	0.031				
Centre in the capital city	2.02	1.68–2.43	<0.001	2.14	1.50–3.05	<0.001				
University centre	2.09	1.62–2.69	<0.001							
Cardiologist	1.55	1.26–1.92	<0.001							
Antiplatelet drug only	OR	95%CI	P	OR	95%CI	P	OAC (alone or in combination)	OR	95%CI	P
CHA ₂ DS ₂ -VAsC (cont. variable)	1.08	1.01–1.16	0.021				HASBLED ≥3	1.28	1.02–1.61	0.036
Age ≥80 years	1.88	1.42–2.49	<0.001	1.99	1.46–2.73	<0.001	Hypertension	1.76	1.40–2.22	<0.001
Coronary artery disease (any)	1.63	1.26–2.10	<0.001	1.35	1.03–1.77	0.033	Age ≥80 years	0.52	0.39–0.67	<0.001
Stable coronary artery disease	1.59	1.23–2.07	<0.001				Coronary artery disease (any)	0.74	0.60–0.92	0.007
Aortic valve disease	2.52	1.64–3.86	<0.001	1.52	1.06–2.17	0.022	Mitral valve disease	1.29	1.03–1.62	0.030
Other cardiac disease	0.50	0.28–0.90	0.021				Dilated cardiomyopathy	1.76	1.15–2.67	0.009
Prior TIA	1.78	1.01–3.13	0.047				Thyroid disease	1.57	1.10–2.25	0.013
COPD	1.56	1.13–2.16	0.007	1.55	1.09–2.19	0.014	Known history of AF	1.50	1.16–1.93	0.002
Body mass index	0.96	0.93–0.99	0.003	0.97	0.94–0.99	0.033	Paroxysmal AF	0.32	0.24–0.42	<0.001
Known history of AF	0.57	0.43–0.74	<0.001	0.68	0.51–0.91	0.010	Body mass index	1.06	1.03–1.08	<0.001
Paroxysmal AF	1.91	1.50–2.48	<0.001	2.42	1.84–3.20	<0.001	Centre in the capital city	1.98	1.58–2.49	<0.001
Centre in the capital city	0.45	0.34–0.59	<0.001	0.40	0.30–0.54	<0.001	Treatment by a cardiologist	1.37	1.06–1.78	0.016
Treatment by a cardiologist	0.68	0.51–0.90	0.007	0.67	0.50–0.91	0.010				

Table 3. Determinants of the use of antithrombotic therapies for stroke prevention in AF patients (see also Supplemental Table 2–5). OAC: oral anticoagulant; OR: Odds Ratio; CI: Confidence Interval; DAPT: dual antiplatelet drug therapy; AF: atrial fibrillation; MI: Myocardial infarction; PCI: percutaneous coronary intervention; CKD: chronic kidney disease; COPD: chronic obstructive pulmonary disease; TIA: transient ischemic attack.




Figure 3. The use of oral anticoagulation by AF clinical type. APLT: antiplatelet therapy; OAC: oral anticoagulant.

On univariate analysis, increasing CHA₂DS₂-VASc score and prior transient ischemic attack (TIA) were also associated with increased use of aspirin only (Table 3).

Other antithrombotic therapies. Independent predictors of dual antiplatelet drug therapy (DAPT) use were PCI, any CAD, COPD and paroxysmal AF, whilst patients with known history of AF were less likely to receive DAPT (Table 3). Independent predictors of the use of OAC combined with one or two antiplatelet drugs were CAD, PCI and treatment in a hospital-based centre, whilst patients aged ≥ 80 years were less likely to be given such therapy. Increasing HASBLED score was positively associated with the use of combined therapy (Table 3).

Indices of VKA anticoagulation quality. An International Normalized Ratio (INR) obtained within previous 3 weeks was available in 946 (79.0%) of 1198 patients who were previously taking a VKA for at least 6 months or longer. The most recent INR value ranged from 1 to 10 (mean 2.42, SD 1.0, median 2.28), and was within the target range of 2.0 to 3.0 in 522 patients (55.2%) whilst in 281 (29.5%) and 143 patients (15.1%) the INR was below and above the target range, respectively.

The Time in Therapeutic Range (TTR) from the previous 3 months was available in only 224 patients (18.7%). Mean TTR was $49.5\% \pm 22.3\%$ (median 50.0%, range 10–100%) and only 66 patients (29.5%) had a TTR of $\geq 65\%$. Labile INR was reported in 439 patients (36.6%) in whom TTR was not available.

Sensitivity analyses. AF was the main reason for enrolling visit or hospitalization in 1329 patients (49.9%). Compared to the main cohort, these patients were younger (mean age 66.9 ± 11.3 years), with lower CHA₂DS₂-VASc (mean 2.95 ± 1.74) and lower HASBLED score (1.72 ± 1.19), all $p < 0.01$. OAC was given to 1000 patients (75.2%). Univariate and multivariable determinants of OAC use in patients with AF as the main reason for enrolling visit or hospitalization are shown in Supplemental Table 4. Similar to the main cohort, there was no significant relationship between the CHA₂DS₂-VASc score and the use of OAC in the multivariable analysis. The use of OAC was driven by the presence of hypertension, younger age (< 75 years), increasing BMI, non-cardiac comorbidities (thyroid disease), centre localization (capital city) and university centre type, whilst patients with COPD, malignancy and paroxysmal AF were less likely to be prescribed OAC (Supplemental Table 4).

An acute coronary syndrome was the main reason for enrolling hospitalization in 206 patients (7.7%) and they were excluded from this analysis. In the remaining cohort of 2457 patients (mean age 68.9 ± 11.0 years, mean CHA₂DS₂-VASc 3.42 ± 1.77 , mean HASBLED 1.97 ± 1.23) OAC was prescribed to 1843 patients (75.0%), and the use of OAC was driven by broadly similar determinants as in the main cohort (Supplemental Table 5).

Discussion

This snapshot survey provides, for the first time, a contemporary insight into routine clinical practice in AF management from a large region of Europe where data on AF management are generally scarce, especially since countries participating in this survey were largely under-represented in recent European AF surveys^{2,4,5}. This survey therefore complements the European AF data and reduces a gap in the European ‘map’ of contemporary real-world management of AF.

Our findings suggest that the overall use of OAC for stroke prevention in the Balkan region is relatively high (~74%), but poorly associated with individual patient stroke risk as recommended by the European Society of Cardiology guidelines¹. Whilst mostly VKA were used, the quality of anticoagulation was poor, with less than a third of patients having a TTR of $\geq 65\%$. Overall, the use of NOACs was slightly higher than recently reported in a ‘real-world’ European survey³, and the use of antiplatelet drugs was comparably high as in other parts of Europe. We also observed significant differences in the use of antithrombotic therapies according to the physician speciality and health centre location. Our results have important practical implications and may help in recognizing the ‘action points’ needed to improve the management of AF patients at risk of stroke in daily clinical practice in the Balkan region.

The Balkan-AF cohort stroke risk profile was broadly similar to recent data from the EURObservational Pilot AF Registry³, with some minor differences (e.g., hypertension, diabetes mellitus and prior stroke were slightly more prevalent, whilst CAD, HF and valvular disease were slightly less frequent in the Balkan-AF cohort). However, bleeding risk was higher, and the difference might possibly be driven by labile INRs in many patients.

The overall use of OAC in Balkan-AF cohort was close to that in recent European reports^{3,4}, but was not significantly associated with CHA₂DS₂-VASc score, even when categorised to < 2 vs. ≥ 2 . Despite clear evidence of low stroke risk in male AF patients with a CHA₂DS₂-VASc score of 0 and those with a score of 1 due to female sex^{8–12}, in whom no antithrombotic therapy is recommended¹, as many as 56.5% of such patients in the Balkan-AF cohort received OAC despite only a minority being scheduled for cardioversion or AF ablation.

Some deviations from evidence-based stroke prevention strategies were evident in the Balkan-AF cohort. For example, elderly patients were more likely to receive aspirin, despite clear evidence of net benefit with OAC in elderly^{13,14}. Also, patients with stable CAD were more likely to receive aspirin or DAPT, or OAC plus antiplatelet drugs instead of OAC monotherapy, although evidence showed that combining OAC with antiplatelet drugs in AF patients with stable vascular disease resulted only in increased risk of major bleeding with no additional reduction of thromboembolism^{15,16}. Such patterns of OAC use were also noted in other European countries⁶.

Independent predictors of increased OAC use in the Balkan-AF survey were hypertension, dilated cardiomyopathy and mild-to-moderate mitral valve disease (essentially mitral regurgitation). In contrast to mitral stenosis, data on the risk of stroke in AF patients with mitral regurgitation are controversial¹⁷ and decisions on OAC should be driven by the presence of well documented stroke risk factors.

The ‘chronicity’ of AF strongly influenced OAC use in the Balkan-AF cohort, with a 56% lower probability for OAC in paroxysmal AF and a 49% greater probability of OAC therapy in patients with history of AF. A large

body of evidence suggests that stroke risk is comparable with paroxysmal or permanent AF^{18,19}. Recently, this has been challenged by a meta-analysis of two non-anticoagulated AF cohorts with systematically adjudicated adverse events which showed AF type to be a strong independent predictor of stroke²⁰, but the annual stroke rate with paroxysmal AF was still sufficiently high (2.1%) to warrant OAC therapy¹. Hence, the decision on OAC should be guided by the presence of conventional stroke risk factors.

The use of aspirin alone was high across all CHA₂DS₂-VAsC score strata and increased from 10% to 17.2% with increasing score. Despite sufficient evidence of only modest efficacy and similar safety compared to OAC^{13,14}, aspirin is still used as monotherapy in around 15% of AF patients in Europe². Independent predictors of aspirin use in the Balkan-AF survey (i.e., advanced age, paroxysmal AF, COPD, mild-to-moderate aortic valve disease) likely reflect a mixture of misperceptions of aspirin better safety (e.g., in elderly) or of lower stroke risk (e.g., paroxysmal AF, COPD, etc.) and an unjustified favouring of aspirin over OAC (e.g., in patients with stable CAD or aortic valve disease).

However, the HASBLED score was not a significant determinant of OAC use in our cohort (indeed, the use of OAC increased with increasing HASBLED), suggesting that either the role of HASBLED was correctly interpreted (that is, the score was used to flag up modifiable bleeding risk factors, and not to preclude OAC use) or perhaps the score was ignored. The latter seems to be the case particularly with the use of combined OAC plus antiplatelet drug therapy, which increased with increasing HASBLED score. Alternatively, sicker patients might have needed such therapy more often.

Around 80% of AF patients in our survey were managed by a cardiologist, and these patients were less likely to receive aspirin. Patients managed in health centres in the capital cities less often received aspirin and more often were given OAC, the latter likely resulting from the clustering of tertiary health centres in the capital cities in most of participating countries. Of note, similar influence of clinical background and specialty of treating physicians on antithrombotic treatment strategies in AF patients has been also described in other European countries²¹. However, the availability of a cardiologist with proper level of expertise for stroke risk management in AF patients in real-world clinical practice may significantly differ among countries.

The signal of poor quality of anticoagulation with VKA in the Balkan-AF cohort (as reflected by a low proportion of patients with a TTR of $\geq 65\%$ and high proportion of patients with labile INRs) is particularly worrisome and calls for urgent action, as suboptimal TTR (<65–70%) is associated with an excess of both stroke and bleeding (and mortality)^{15,22,23}. Indeed, a TTR of $>70\%$ is recommended in guidelines and position documents, when VKAs are used¹⁵. Also, missing TTR in $>80\%$ of patients in the Balkan-AF cohort might indicate that the calculation of TTR is not commonly used in routine clinical practice in Balkan countries. Given the low quality of therapy with VKA compared to other European countries²⁴, increasing use of NOACs (17% of all patients taking OAC) is encouraging, as it could facilitate adequate stroke prevention in Balkan countries. An analysis of factors influencing the choice of NOAC over VKA in Balkan countries is underway.

This study is limited by its observational snapshot registry design, but we made every effort to include consecutive patients. Although we tried to capture a sample representative of real-world clinical practice by recruiting a range of different types of centres in each country (i.e., university and non-university hospitals and outpatient centres in- and outside the capital cities), there still may be a selection bias due to variable health care setting in the participating countries.

The proportion of cardiologists versus internal medicine specialists participating in the Balkan-AF survey may not fully reflect daily practice in the participating countries, since we might have not adequately covered the rural areas. Still, participating centres situated outside capital cities enrolled about 55% of patients, and in smaller countries many AF patients are often referred to the tertiary centres at least for initial evaluation.

Since TTR was available in a small proportion of patients treated with OAC, our results may under- or overestimate the quality of VKA treatment in our cohort, and we cannot conclude whether or not VKA therapy is systematically monitored in clinical practice.

Conclusion

This survey provide important insights into contemporary routine practices for stroke prevention in AF patients in Balkan countries, thus reducing a gap in the European 'map' of contemporary real world management of AF. Our results show a broadly similar patient stroke risk profile in the Balkan Region and similar OAC use as in other European countries. However, routine practices for stroke prevention in AF patients in Balkan countries are less influenced by the presence of conventional well documented stroke risk factors in AF patients, and the quality of VKA therapy is low. Thus, additional efforts are needed to increase the adherence to AF guidelines and improve the management of AF-related risk of stroke in routine clinical practice in the Balkan Region.

References

1. Camm, A. J. *et al.* 2012 focused update of the ESC Guidelines for the management of atrial fibrillation: an update of the 2010 ESC Guidelines for the management of atrial fibrillation. Developed with the special contribution of the European Heart Rhythm Association. *Eur Heart J* **33**, 2719–47 (2012).
2. Lip, G. Y. *et al.* A prospective survey in European Society of Cardiology member countries of atrial fibrillation management: baseline results of EURObservational Research Programme Atrial Fibrillation (EORP-AF) Pilot General Registry. *Europace* **16**, 308–19 (2014).
3. Lip, G. Y. *et al.* Prognosis and treatment of atrial fibrillation patients by European cardiologists: one year follow-up of the EURObservational Research Programme-Atrial Fibrillation General Registry Pilot Phase (EORP-AF Pilot registry). *Eur Heart J* **35**, 3365–76 (2014).
4. Kirchhof, P. *et al.* Management of atrial fibrillation in seven European countries after the publication of the 2010 ESC Guidelines on atrial fibrillation: primary results of the PREvention of thromboembolic events–European Registry in Atrial Fibrillation (PREFER in AF). *Europace* **16**, 6–14 (2014).
5. Arbelo, E. *et al.* The atrial fibrillation ablation pilot study: a European Survey on Methodology and results of catheter ablation for atrial fibrillation conducted by the European Heart Rhythm Association. *Eur Heart J* **35**, 1466–78 (2014).

6. Lip, G. Y. *et al.* Regional differences in presentation and treatment of patients with atrial fibrillation in Europe: a report from the EURObservational Research Programme Atrial Fibrillation (EORP-AF) Pilot General Registry. *Europace* **17**, 194–206 (2015).
7. Potpara, T. S., Balkan, A. F. I. & Lip, G. Y. Patterns in atrial fibrillation management and 'real-world' adherence to guidelines in the Balkan Region: an overview of the Balkan-atrial fibrillation survey. *Eur Heart J* **36**, 1943–4 (2015).
8. Olesen, J. B., Torp-Pedersen, C., Hansen, M. L. & Lip, G. Y. The value of the CHA2DS2-VASc score for refining stroke risk stratification in patients with atrial fibrillation with a CHADS2 score 0–1: a nationwide cohort study. *Thromb Haemost* **107**, 1172–9 (2012).
9. Olesen, J. B. *et al.* Risks of thromboembolism and bleeding with thromboprophylaxis in patients with atrial fibrillation: A net clinical benefit analysis using a 'real world' nationwide cohort study. *Thrombosis and Haemostasis* **106**, 739–749 (2011).
10. Friberg, L., Rosenqvist, M. & Lip, G. Y. Net clinical benefit of warfarin in patients with atrial fibrillation: a report from the Swedish atrial fibrillation cohort study. *Circulation* **125**, 2298–307 (2012).
11. Potpara, T. S. *et al.* A 12-year follow-up study of patients with newly diagnosed lone atrial fibrillation: implications of arrhythmia progression on prognosis: the Belgrade Atrial Fibrillation study. *Chest* **141**, 339–47 (2012).
12. Lip, G. Y. Stroke and bleeding risk assessment in atrial fibrillation: when, how, and why? *Eur Heart J* **34**, 1041–9 (2013).
13. Mant, J. *et al.* Warfarin versus aspirin for stroke prevention in an elderly community population with atrial fibrillation (the Birmingham Atrial Fibrillation Treatment of the Aged Study, BAFTA): a randomised controlled trial. *Lancet* **370**, 493–503 (2007).
14. Connolly, S. J. *et al.* Apixaban in patients with atrial fibrillation. *N Engl J Med* **364**, 806–17 (2011).
15. De Caterina, R. *et al.* Vitamin K antagonists in heart disease: current status and perspectives (Section III). Position paper of the ESC Working Group on Thrombosis–Task Force on Anticoagulants in Heart Disease. *Thromb Haemost* **110**, 1087–107 (2013).
16. Bernard, A. *et al.* Anticoagulation in patients with atrial fibrillation undergoing coronary stent implantation. *Thromb Haemost* **110**, 560–8 (2013).
17. De Caterina, R. & Camm, A. J. What is 'valvular' atrial fibrillation? A reappraisal. *Eur Heart J* **35**, 3328–35 (2014).
18. Hart, R. G. *et al.* Stroke with intermittent atrial fibrillation: incidence and predictors during aspirin therapy. Stroke Prevention in Atrial Fibrillation Investigators. *J Am Coll Cardiol* **35**, 183–7 (2000).
19. Friberg, L., Hammar, N. & Rosenqvist, M. Stroke in paroxysmal atrial fibrillation: report from the Stockholm Cohort of Atrial Fibrillation. *Eur Heart J* **31**, 967–75 (2010).
20. Vanassche, T. *et al.* Risk of ischaemic stroke according to pattern of atrial fibrillation: analysis of 6563 aspirin-treated patients in ACTIVE-A and AVERROES. *Eur Heart J* **36**, 281–7a (2015).
21. Kirchhof, P. *et al.* Impact of the type of centre on management of AF patients: surprising evidence for differences in antithrombotic therapy decisions. *Thromb Haemost* **105**, 1010–23 (2011).
22. Gallagher, A. M., Setakis, E., Plumb, J. M., Clemens, A. & van Staa, T. P. Risks of stroke and mortality associated with suboptimal anticoagulation in atrial fibrillation patients. *Thromb Haemost* **106**, 968–77 (2011).
23. Gallego, P. *et al.* Cessation of oral anticoagulation in relation to mortality and the risk of thrombotic events in patients with atrial fibrillation. *Thromb Haemost* **110**, 1189–98 (2013).
24. Le Heuzey, J. Y. *et al.* Differences among western European countries in anticoagulation management of atrial fibrillation. Data from the PREFER IN AF registry. *Thromb Haemost* **111**, 833–41 (2014).

Acknowledgements

We thank all Balkan-AF investigators and Ms Zlatiborka Mijatovic for their hard work and contribution.

Author Contributions

T.S.P. study design, data collection, data analysis, writing of the manuscript, MMP: eCRF development, data collection, manuscript review, G.A.D., E.T., A.G., Z.K., S.M. and L.J.M. coordinating the study, data collection, manuscript reviewing, R.M., E.B., G.M., V.P., E.S.D., S.L.P., H.D.J., D.L., A.B. & S.B. data collection, manuscript reviewing, GYHL: study design, data analysis and interpretation, writing of the manuscript. The BALKAN-AF Investigators: data collection.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: TSP: speaker fees from Pfizer and Bayer; MMP: no conflicts of interest to declare; GAD: speaker fees from Boehringer Ingelheim; ET: speaker fees from Bayer, Pfizer, Boehringer Ingelheim, Astra Zeneca, Servier, Merck Serono, Gedeon Richter, Actavis, Berlin Chemie, AG, ZK, SM: no conflicts of interest to declare; LJM: speaker fees from Bayer and Boehringer Ingelheim; RM: no conflicts of interest to declare; EB: speaker fees from Astra Zeneca, Berlin Chemie, Pfizer, Antibiotice Iasi, Gedeon Richter; GM: speaker fees from Pfizer and Boehringer Ingelheim; VP, ESD, SLP, HDj, DL, AB: no conflicts of interest to declare; SB: speaker fees from Bayer, Pfizer, Boehringer Ingelheim; GYHL: consultant fees from Bayer, Merck, Sanofi, BMS/Pfizer, Daiichi-Sankyo, Biotronic, Medtronic, Portola and Boehringer Ingelheim and speaker fees from Bayer, BMS/Pfizer, Boehringer Ingelheim, Daiichi-Sankyo, Medtronic.

How to cite this article: Potpara, T. S. *et al.* Stroke prevention in atrial fibrillation and 'real world' adherence to guidelines in the Balkan Region: The BALKAN-AF Survey. *Sci. Rep.* **6**, 20432; doi: 10.1038/srep20432 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

Consortia

The BALKAN-AF Investigators

Tatjana S. Potpara¹⁵, Marija Polovina¹⁵, Srdjan Milanov¹⁶, Marija Pavlovic¹⁶, Marijana Petrovic¹⁶, Stefan Simovic¹⁶, Gorana Mitic¹⁷, Marko Milanov¹⁸, Jelena Savic¹⁹, Sanja Gnip¹⁹, Pavica Radovic¹⁹, Snezana Markovic²⁰, Ivana Koncarevic²¹, Jelena Gavrilovic²¹, Tijana Acimovic²¹, Dijana Djikic²², Semir Malic²², Jusuf Hodzic²², Milovan Stojanovic²³, Marina Deljanin Ilic²³, Milan Zlatar²⁴, Dragan Matic²⁴, Snezana Lazic²⁵, Vladan Peric²⁵, Sanja Markovic²⁵, Snezana Kovacevic²⁶, Aleksandra Arandjelovic²⁷, Milika Asanin²⁸, Marija Zdravkovic²⁹, ROMANIA: Gheorghe-Andrei Dan³⁰, Anca Breha³⁰, Anca Rodica Dan³⁰, Rodica Musetescu³¹, Mircea Ioachim Popescu³², Elisabeta Badila³³, Catalina Arsenescu Georgescu³⁴, Sorina Pop³⁵, Raluca Popescu³⁶, Simina Neamtu³⁷, Floriana Oancea³⁸, BULGARIA: Elina Trendafilova³⁹, Elena Dimitrova³⁹, Evgenii Goshev³⁹, Anna Velichkova³⁹, Stanislav Petranov⁴⁰, Delyana Kamenova⁴¹, Penka Kamenova⁴², Svetoslava Elefterova⁴³, Valentin Shterev⁴⁴, Maria Zekova⁴⁵, Stela Diukiandzhieva⁴⁶, Boiko Dimitrov⁴⁷, Tihomir Sotirov⁴⁸, Valentina Simeonova⁴⁹, Dimitrina Drianovska⁴⁹, Liliya Ivanova Vasileva Boiadzhieva⁵⁰, Darina Buchukova⁵¹, ALBANIA: Artan Goda⁵², Vilma Papparisto⁵², Hortensia Gjergo⁵², Alma Mijo⁵², Ervina Shirka⁵², Viktor Gjini⁵³, Uliks Ekmekci⁵⁴, Ina Refatllari⁵⁴, BOSNIA & HERZEGOVINA: Zumreta Kusljugic⁵⁵, Daniela Loncar⁵⁶, Denis Mrsic⁵⁶, Hazim Tulumovic⁵⁶, Belma Pojskic⁵⁷, Alma Sijamija⁵⁸, Amira Bijedic⁵⁹, Indira Karamujic⁵⁹, Irma Bijedic⁶⁰, Sanela Halilovic⁶¹, Sekib Sokolovic⁶², CROATIA: Sime Manola⁶³, Ivan Zeljkovic⁶³, Nikola Pavlovic⁶³, Vjekoslav Radeljic⁶³, Sandro Brusich⁶⁴, Ante Anic⁶⁴, Melita Jeric⁶⁵, Petar Pekic⁶⁶, Kresimir Milas⁶⁷, MONTENEGRO: Ljilja Music⁶⁸, Nebojsa Bulatovic⁶⁸, Ana Nenezic⁶⁹, Dijana Asanovic⁶⁹

¹⁵Cardiology Clinic, University Clinical Center of Serbia, School of Medicine, Belgrade University, Belgrade.

¹⁶Cardiology Clinic, University Clinical Center of Kragujevac, Kragujevac. ¹⁷Hematology Clinic, University Clinical Center of Vojvodina, University of Novi Sad, Medical Faculty, Novi Sad. ¹⁸University Clinical Center Zvezdara, Cardiology Department, Belgrade. ¹⁹Hematology Clinic, University Clinical Center of Vojvodina, Novi Sad. ²⁰General Hospital, Pirot. ²¹University Clinical Center Bezanijaska kosa, Cardiology Department, Belgrade. ²²General Hospital Gracanica, Gracanica. ²³Clinic for Cardiovascular Diseases, Institute Niska Banja, Niska Banja. ²⁴Emergency Center, Coronary Care Unit, University Clinical Center of Serbia, Belgrade. ²⁵Internal Medicine Clinic, University Clinical Center of Pristina, Pristina. ²⁶General Hospital Sabac, Sabac. ²⁷University Clinical Center Zvezdara, Cardiology Department, School of medicine, Belgrade University, Belgrade. ²⁸Emergency Center, Coronary Care Unit, Clinical Center of Serbia, School of medicine, Belgrade University, Belgrade. ²⁹University Clinical Center Bezanijaska kosa, Cardiology Department, School of medicine, Belgrade University, Belgrade. ³⁰Medicine University "Carol Davila", Colentina University Hospital, Bucharest. ³¹Cardiology Centre – County Emergency Hospital, Craiova. ³²Cardiology department, Emergency Hospital, Oradea. ³³Emergency Clinical Hospital, Internal Medicine Department, Bucharest. ³⁴Institute for Cardiovascular Diseases 'Prof Dr George I.M. Georgescu', Iasi. ³⁵General Practice in Cluj-Napoca, Cluj-Napoca. ³⁶Colentina University Hospital, Cardiology Department, Bucharest. ³⁷General Practice in Timisoara, Timisoara. ³⁸Cardiology Department, County Emergency Hospital, Sibiu. ³⁹National Heart Hospital, Coronary Care Unit, Sofia. ⁴⁰Health Center, Bugras, Bugras. ⁴¹Health Center Vidin, Vidin. ⁴²District Hospital/MHAT, Ruse. ⁴³University Hospital, Varna. ⁴⁴Health Center, Varna. ⁴⁵University Hospital, Plevna. ⁴⁶Health Center Gorna Oryahovitza, Gorna Oryahovitza. ⁴⁷Community Hospital, Montana. ⁴⁸Community Hospital, Haskovo. ⁴⁹Health Center, Sofia. ⁵⁰MBAL Dr Stefan Cherkezov AD/ Second Internal Department, Veliko Trnovo. ⁵¹University of Medicine, Sofia. ⁵²Clinic of Cardiology, University Hospital Center Mother Theresa, Tirana. ⁵³Regional Hospital Fier. ⁵⁴Clinic of Internal Medicine, University Hospital Center Mother Theresa, Tirana. ⁵⁵Clinic of Internal Medicine, Cardiology Department, University Clinical Center Tuzla, Medical Faculty, Tuzla. ⁵⁶Clinic of Internal Medicine, Department of Intensive Care and Therapy, University Clinical Center Tuzla, Tuzla. ⁵⁷General Hospital, Zenica. ⁵⁸General Hospital, Travnik. ⁵⁹Clinic of Internal Medicine, Cardiology Department, University Clinical Center Tuzla, Tuzla. ⁶⁰Clinic of Internal Medicine, University Clinical Center Tuzla, Tuzla. ⁶¹Clinic of Heart and Rheumatologic Diseases, University Clinical Center Sarajevo, Sarajevo. ⁶²Clinical Center "Sestre Milosrdnice", Zagreb. ⁶³Clinical Center Rijeka, Rijeka. ⁶⁴General Hospital, Zadar. ⁶⁵General Hospital Varazdin, Varazdin. ⁶⁶Clinical Hospital "Sveti Duh", Zagreb. ⁶⁷General Hospital, Pula. ⁶⁸Cardiology Clinic, University Clinical Center of Montenegro, University of Podgorica, Medical Faculty, Podgorica. ⁶⁹Cardiology Clinic, University Clinical Center of Montenegro, Podgorica.