

Influence of a dispergent on the cell membrane potential

Žauhar, Gordana; Lekić, Andrica; Šolić, Franjo

Source / Izvornik: **Periodicum biologorum, 1995, 97, 343 - 346**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:184:359761>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2024-12-25**



Repository / Repozitorij:

[Repository of the University of Rijeka, Faculty of Medicine - FMRI Repository](#)





Original scientific paper

Influence of a dispergent on the cell membrane potential

GORDANA ŽAUHAR, ANDRICA LEKIĆ and FRANJO ŠOLIĆ

Received October 11, 1995.

INTRODUCTION

In recent decades the effects of different chemicals, heavy metals and complex wastes of effluents on freshwater organisms of aquatic ecosystems have been described (8, 9). The injury of several organisms structures, and disturbances of physiological processes, particularly photosynthesis in freshwater algae were described specifically (7, 6).

The membrane potential of cell is very sensitive to the physical and chemical effect of the surroundings (5, 1).

The purpose of this work was to investigate the influence of the dispergent, as the possible pollutant of aquatic ecosystems on the bipotential of algal cells.

MATERIAL AND METHODS

In the experiments described herein, the dispergent HEMISOL, manufactured in Modriča, was used. Hemisol belongs to the medium toxicity group of dispergents.

Experiments were performed on freshwater algae *Nitella* belonging to the *Characeae* family. The algae were cultured at the room temperature in glass containers filled with tap water. All experiments were done on cells constituting the second internodus from the top. The length of cells varied from 10 to 15 mm, and the diameter from 0.30 mm to 0.45 mm. After isolation, the cells were kept for one hour in a medium containing equal parts of 0.1 mM KCl and 1.0 mM NaCl with the aim to obtain the balance of ions (basal solution).

The membrane potentials were measured by standard microelectrode technique, using glass microcapillaries filled with 3 M KCl solution. A microelectrode was inserted into the vacuola of the cells, and the reference electrode was immersed into the surrounding solution. The potentials measurements were carried out in a grounded metallic box to pre-

vent the influence of electromagnetic fields from outside. All measurements were done at the room temperature and the constant white light intensity. The apparatus used has been already described (3).

The membrane potentials of the cells were measured continuously, first in the basal solution, and then in basal solution contaminated with various concentrations of a dispergent. The bathing solution was contaminated with increasing doses of the dispergent ranging from 0.001‰ to 100‰. The dispergent was added each time when the stable membrane potential was reached. The membrane potentials of contaminated cells were measured after contamination until cell death. For investigations in post treatment period the cells were kept in basal solution at the daily illumination and room temperature.

The results were evaluated by using the methods of testing the differences of arithmetical means and proportions for small independent samples, and analysis of variance (4).

RESULTS

The changes of resting potentials during contamination with increasing doses of a dispergent are illustrated in Figure 1. The values of potentials changes are indicated by ΔM and expressed in percentages of initial resting potentials of non-contaminated basal solution. Each point represents the mean value of 9-6 measurements. Error bars represent 95% confidence interval. It is evident that the dispergent caused decrease of resting potentials, i.e. the depolarisation of the cells membrane. The value of depolarisation increased with the dispergent concentration.

During the experiments the polarisation in some cells at the concentrations of 0.1‰ and 1.0‰ of the dispergent occurred. In Table 1 the observed number of such cells are indicated by N_p , whereas in the bars the total number of treated cells by N .




FIGURE 1. The changes of resting potentials as a function of dispersant concentration.

The potentials changes seen in both of the experiments are compared in the last column of Table 1. Significant differences were found for dispersant concentration of 0.1‰.

The resting potential is an indicator of cell activity; when the cell dies, the biopotential disappears. After the contamination the membrane potentials were measured at intervals of two days until the cell death. Three different dispersant concentrations were used. Cells survivals are represented in Table 2. Each value presents the means of 7-6 observations (\pm mean error). The differences of cells survival was tested by the analysis of variance. The difference of survivals between contaminated and non-contami-

TABLE 1

The changes of resting potentials between successive and directly dispersant contaminations

concentration per thousand	successive contamination		direct contamination		difference (significancy)
	N_p (N)	Δ PM %	N_{HP} (N)	Δ PM %	
0.1	4 (8)	-26.4 ± 6.2	4 (7)	1.3 ± 1.8	27.7 ± 6.5 ($P < 0.05$)
1	2 (7)	-37.9 ± 8.2	0 (6)	-36.6 ± 9.1	1.3 ± 12.2 ($P > 0.05$)

To investigate how each of these concentrations influences the potential changes, the basal solution was contaminated directly with 0.1‰ or 1.0‰ of dispersant concentrations. The results of both experiments are shown in Table 1. The number of hyperpolarised cells is indicated by N_{HP} . The hyperpolarisation was found only at the dispersant concentration of 0.1‰.

TABLE 2

Survival of the cells at different concentrations of dispersant.

concentration per thousand	experimental cells			control cells
	0.1	1	100	0
N	7	6	6	6
survival of the cells / days	11 ± 2.5	15.6 ± 3.5	12.5 ± 1.7	25.8 ± 3.2

nated cells is statistically significant at the dispersant concentration of 0.1‰ and 100‰.

In treated (experimental) and control cells, the oscillations of membrane potentials was observed in certain sets of cells. In Table 3 the number of these cells is expressed in percentage. The oscillations in experimental cells are lower than in control cells, where differences were not statistically significant.

TABLE 3

Oscillations of membrane potential in contaminated and not contaminated cells.

	experimental cells	control cells
N	22	17
Osc %	27.3	58.8
Δp %	31.5 ± 15.9	
significancy	$0.05 < P < 0.055$	

DISCUSSION

Comparing the potentials changes in cells contaminated with the increasing doses of dispersant concentrations from 0.001‰ to 0.1‰ and potential changes in cells contaminated with a single dose of a dispersant, we found that contamination with the dispersant at the concentration of about 0.1‰ resulted in hyperpolarisation. Namely, the potentials changes in successively contaminated cells suggested that the membrane permeability was effected by previous contaminations.

The dispersant decreased the life time of cells. This conclusion is not valid for concentration of 1 symbol 1‰ because of a high variability of cells survival. To achieve a reasonable conclusion, the number of cells in this sample should be higher.

The oscillations of membrane potentials in a certain number of the cells are present continuously. It is believed that these are generated spontaneously, but their origin is not clear yet (2).

REFERENCES

- MARIĆ G, ŠOLIĆ F, MILOTIĆ B, ŽAUHAR G 1990 Membrane potential of *Nitella* as an indicator of chemical contamination of water. *Period Biol* 92: 453-454
- MARKEVICH N I, SELKOV E E 1990 Theoretical analysis of the oscillations in the membrane potential in *Characean algae*. *Studia biophysica* 138: 147-156
- MILOTIĆ B, ŠOLIĆ F 1982 The effect of electromagnetic visual light irradiation on biopotentials of plant cells. *Period Biol* 84: 163-165
- PETZ B 1981 Uvod u analizu varijance. In: Rubelli-Ligutić L (ed) Osnovne statističke metode za nematematičare. SNL, Zagreb, p 258
- PYATYGIN S S, OPRITOV V A 1990 Vlyanie temperatury na generaciju potencialov deistviya vzbudimymi klietkami vysshego rastenya. *Biofizika* 35: 444-449
- RÖSICK E, STADTLANDER K, AHLERS J 1985 Environmental chemicals and biomembranes. *Chemosphere* 14: 529-544
- STRATTON G W 1989 Effect of the Solvent Acetone on Membrane Integrity in the Green Alga *Chlorella pyrenoidosa*. *Bull Environ Contam Toxicol* 42: 754-760
- WALSH G E, BAHNER L H, HORNING W B 1980 Toxicity of textile mill effluents to freshwater and estuarine algae, crustaceans and fishes. *Environ Pollut (Ser A)* 21: 169-179
- WALSH G E, DUKE K M, FOSTER R B 1982 Algae and crustaceans as indicator of bioactivity of industrial wastes. *Water Res* 16: 879-883

ABSTRACT

Influence of a dispersant on the cell membrane potential

Background and purpose: Many investigations of the influence of temperature, ultrasound, electromagnetic waves, heavy metals, and ionic solutions on the cell membrane potentials have confirmed that the membrane potential is very sensitive to physical and chemical effect of the surroundings. The purpose of this work was to investigate the influence of a dispersant, as the possible pollutant of aquatic ecosystems, upon the membrane potential of algal cells.

Material and methods: Experiments were performed on freshwater algae *Nitella* belonging to the Characeae family. The potentials were measured using the microelectrode technique. A microelectrode was inserted into the cell vacuole and the reference electrode was immersed into the basal solution containing equal parts of 0.1 mM KCl and 1.0 mM NaCl. This solution was contaminated with various concentrations of a dispersant (from 0.0001% to 10%).

Results: The observed changes of resting membrane potentials showed that the dispersant had caused the depolarisation of the membrane, proportionally with concentration if the dispersant was successively increased. If the basal solution was contaminated directly with dispersant concentration of 0.01%, the hyperpolarization of the membrane occurred. Control cells lived longer than experimental ones that were contaminated with 0.01% and 10% concentrations of the dispersant. Control cells had more oscillations than experimental cells, but the observed difference is not statistically significant.

Conclusion: The observed changes of resting membrane potentials showed that dispersant caused the depolarisation of the membrane, proportionally with its concentration.

Dept. of Physics, Faculty of Medicine, Braće Branchetta 20, 51000 Rijeka, Croatia

Correspondence: Gordana Žauhar, Dept. of Physics, Faculty of Medicine, Braće Branchetta 20, 51000 Rijeka, Croatia

Key words: membrane potential, dispersant