

Is encephalopathic brain genetically more prone to dementia?

Šoša, Ivan; Strenja-Linić, Ines

Source / Izvornik: **Open Medicine, 2013, 8, 730 - 731**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.2478/s11536-013-0229-7>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:184:340548>

Rights / Prava: [Attribution-NonCommercial-NoDerivatives 4.0 International/Imenovanje-Nekomercijalno-Bez prerada 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2024-07-13**



Repository / Repozitorij:

[Repository of the University of Rijeka, Faculty of Medicine - FMRI Repository](#)



Is encephalopathic brain genetically more prone to dementia?

Letter to the Editor

Ivan Sosa*¹, Ines Strenja-Linic²

1 Department of Forensic Medicine and Criminalistics: University of Rijeka School of Medicine, B. Branchetta 20, 51 000 Rijeka

2 Rijeka University Hospital; Neurology Clinic; Neurovascular Diseases and Expansive Processes Department

Received 20 January 2013; Accepted 22 May 2013

Disclaimer and acknowledgements statement

All the authors declare that no support for the submitted work has been received. The authors acknowledge any person's engagement that is not acknowledged in the list of authors.

To the Editor,

Encouraging progress in understanding the genetics of Alzheimer's disease (AD) merely highlights the work that remains. AD and vascular dementia are two commonest dementias. Etiology of AD involves genetic factors in 25–40% [1]. Thus far, three fully penetrant genes' mutations have been identified in patients younger than 60 years.

First, this was assigned to the β amyloid precursor protein gene (*APP*) on chromosome 21 and after that to the autosomal dominant gene defects of presenilin-1 (*PS1*) gene on chromosome 14. Presenilin-2 (*PS2*) gene on chromosome 1 was the last connected to early-onset AD. Each of these mutations causes abnormal proteins to be formed. In 2012 for example, Kamboh identifies nine other genes/loci (*CR1*, *BIN1*, *CLU*, *PICALM*, *MS4A4/MS4A6E*, *CD2AP*, *CD33*, *EPHA1*

and *ABCA7*) for late-onset Alzheimer's disease (LOAD) using genome-wide association studies (GWASs) [2].

With identification of more of the genes involved in development of the disorder, and a greater understanding of their action, AD research holds the hope of reducing or potentially eliminating the burden of this overwhelming disease [1]. The single-gene mutations directly responsible for early-onset Alzheimer's disease do not seem to be involved in late-onset Alzheimer's.

A specific gene causing the late-onset form of the disease has not been identified yet. However, one genetic risk factor does appear to increase a person's risk of developing the disease. This is related to the apolipoprotein E (*APOE*) gene found on chromosome 19. *APOE* encodes for a protein that helps carry cholesterol and other types of fat in the bloodstream. *APOE* comes in several different forms, or alleles. Three forms – *APOE* ϵ 2, *APOE* ϵ 3, and *APOE* ϵ 4 – occur most frequently. *APOE* ϵ 4 is present in about 25 to 30 % of the population and in about 40 % of all people with late-onset AD. People who develop AD are more likely to have an *APOE* ϵ 4 allele than people who do not develop the disease.

Testing for the *E-4* allele cannot predict who will get the disease. Two national panels have raised their voice against the use of *apoE-4* as a predictive test for AD. This could be justified even more, if one bears in mind that a great majority of people who carry *E4* will not develop AD, respecting many people who do not.

* E-mail: egle.dalinkeviciene@gmail.com

Just alike, chronic traumatic encephalopathy (CTE) [3], usually seen in athletes (i.e. TBI-related encephalopathy), affects competitors of different backgrounds, origin and colour of skin and people who inherit one or two APOE ϵ 4 alleles tend to develop the disease at an earlier age than those who do not have any APOE ϵ 4 alleles yet more likely if athletes (see McKee *et al.*) [4].

Already damaged nervous system is ideal ground for expression of mutations as suggested by Mattson *et al.* There is always a question, would some dementive disorder come obvious without contact sport history [5]

Ivan Sosa, MD
< Department of Forensic
Medicine and Criminalistics >

Ines Strenja Linic, PhD
< Neurology Clinic >

References

- [1] Hollingworth P., Harold D., Jones L., Owen M.J., Williams J. Alzheimer's disease genetics: current knowledge and future challenges. *Int. J. Geriatr. Psychiatry.*, 2011; 26:793-802
- [2] Kamboh M.I., Demirci F.Y., Wang X., Minster R.L., *et al.* Genome-wide association study of Alzheimer's disease. *Transl Psychiatry.* 2012; 2:e117. doi: 10.1038/tp.2012.45
- [3] Ledic D., Sosa I., Linic I.S., Cvijanovic O., Kovacevic M., Desnica A., Banicek I.. Vomiting as a reliable sign of concussion. *Med Hypotheses.*, 2012;78(1):23-25
- [4] McKee A., Cantu R.C., Nowinski C.J., Hedley-Whyte E.T., Gavett B.E., Budson A.E., Santini V.E., Lee H.S., Kubilus C.A., Stern R.A. Chronic traumatic encephalopathy in athletes: progressive tauopathy after repetitive head injury. *J. Neuropathol. Exp. Neurol.*, 2009; 68: 709-735
- [5] Rabadi M.H., Jordan B.D. The cumulative effect of repetitive concussion in sports. *Clin. J. Sport. Med.*, 2001; 11:194-198