

Antimicrobial activity of the volatile phase of essential oils and their constituents on *Legionella pneumophila*

Bićanić, Lucia; Mežnarić, Silvestar; Gobin, Ivana

Source / Izvornik: **Sanitarno inženirstvo International Journal of Sanitary Engineering Research, 2020, 14, 54 - 61**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.2478/ijser-2020-0005>

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:184:286845>

Rights / Prava: [Attribution-NonCommercial-NoDerivatives 4.0 International/Imenovanje-Nekomercijalno-Bez prerada 4.0 međunarodna](#)

Download date / Datum preuzimanja: **2024-07-25**



Repository / Repozitorij:

[Repository of the University of Rijeka, Faculty of Medicine - FMRI Repository](#)



Antimicrobial activity of the volatile phase of essential oils and their constituents on *Legionella pneumophila*

Lucia BIČANIĆ¹, Silvestar MEŽNARIĆ¹, Ivana GOBIN²

ABSTRACT

Pathogenic bacteria of the genus *Legionella* cause atypical pneumonia known as Legionnaires' disease and flu – like disease known as Pontiac fever. As pathogens of the respiratory system, these bacteria represent a public health problem and there is a need for examine new alternative ways to inactivate them. These bacteria live naturally in water and are transmitted by infectious aerosols. To purify the air, essential oils that show antimicrobial properties are widely used. The anti-*Legionella* activity of five exotic essential oils and five Mediterranean essential oils characteristic for coastal Croatia was examined. Model organism used in experiments was *L. pneumophila* (strain 130b). This experiment was conducting with modified version of sealed plate method using a BCYE medium. The exotic essential oil with highest anti-*Legionella* activity was Niaouli essential oil, and the best anti-*Legionella* activity among Mediterranean essential oils showed Immortelle essential oil. Anti- *Legionella* activity of four main chemical compounds was examined and compound that show significant highest anti-*Legionella* activity was α – pinene. Volatile components of essential oils have a great potential as anti-*Legionella* agents and further research are needed.

Key words: essential oils; volatile components; *Legionella*; antibacterial effect

POVZETEK

Patogene bakterije iz rodu *Legionella* povzročajo atipično pljučnico, poznano pod imenom Legionarska bolezen in Pontiaška vročica. Tako kot drugi povzročitelji boleznih respiratornega sistema tudi te bakterije predstavljajo javnozdravstveni problem. Iz tega sledi nenehna potreba po iskanju novih alternativnih načinov preprečevanja in obvladovanja njihovega pojava. Bakterije iz rodu *Legionella* lahko najdemo v naravnih in umetnih vodnih sistemih, prenašajo pa se preko onesnaženih aerosolov. Za čiščenje zraka se pogosto uporabljajo eterična olja, ki delujejo protimikrobno. Z uporabo BCYE medija smo analizirali protimikrobno delovanje na bakterijo *Legionella pneumophila* (sev 130b). Pri tem je bilo testiranih pet eksotičnih eteričnih olj in pet sredozemskih eteričnih olj, značilnih za obalo Hrvaške. Med eksotičnimi

Original scientific article

Received: 29. 02. 2020

Accepted: 30. 12. 2020

Published: 31. 12. 2020

¹ Student at Faculty of Medicine, University of Rijeka, Braće Branchetta 20, 51000 Rijeka, Croatia

² Department of Microbiology and Parasitology, Faculty of Medicine, University of Rijeka, Braće Branchetta 20, 51000 Rijeka, Croatia

* Corresponding author
Ivana Gobin, PhD, Associate professor
Department of Microbiology and Parasitology, Faculty of Medicine, University of Rijeka, Braće Branchetta 20, 51000 Rijeka, Croatia
E-mail: ivana.gobin@uniri.hr

© 2020 Lucia Bičanić, Silvester Mežnarić, Ivana Gobin. This is an open access article licenced under the Creative Commons Attribution NonCommercial-NoDerivs license as currently displayed on <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

eteričnimi olji smo največjo protimikrobno aktivnost zabeležili pri olju »Niaouli«. Med sredozemskimi eteričnimi olji pa je bilo to olje »Immortelle«. Med glavnimi kemičnimi snovmi ima največji antibakterijski učinek spojina α – pinen. Velik potencial protimikrobnega delovanja hlapnih snovi eteričnih olj na bakterijo *Legionella pneumophila* kaže potrebo po nadaljnjih raziskavah.

Ključne besede: eterična olja; hlapne snovi; *Legionella*; antibakterijski učinek

INTRODUCTION

The genus *Legionella* contain a group of pathogenic bacteria that very often cause respiratory diseases collectively called legionellosis. Species from the genus *Legionella* belong to the family *Legionellaceae*, and the main cause of most legionellosis is *Legionella pneumophila*. Legionellosis has two forms: Pontiac fever and Legionnaires' disease. Legionnaires' disease or atypical pneumonia is a very common and severe systemic disease. Another mild legionellosis that has flu-like symptoms is known as Pontiac fever [1, 2]. The largest, natural habitat of *L. pneumophila* is freshwater environment and very often can be present in various artificial and natural water systems such as cooling towers, air conditioners, humidifiers and dehumidifiers, boilers, swimming pools, fountains and other water systems [3]. The most common cases of infection with this bacterium are associated with travel and staying in different hotels [4]. For this reason, it is important to prevent the presence of this bacteria in places where people are gathering, so that epidemics do not occur. One of such important places are hotels and spa centers where water analysis must be performed regularly, and measures must be taken to prevent legionellosis.

Essential oils are composed of volatile molecules and are produced from aromatic or medicinal plants by a process of secondary metabolism. The largest share, in essential oils, are aromatic hydrocarbons and terpenes, which give them their characteristic pleasant and intense odor. It is very important to note that when isolating oil from the same plant and the same process, do not always results with oil that have the same composition, it will largely depend on various factors such as growth and cultivation conditions and the time of harvesting the plant itself. EOs are usually obtained as a result of hydrodistillation, steam distillation, dry distillation, or the mechanical cold pressing of plants [5]. Essential oils have been used for thousands of years, not only as ingredients of perfumes or for the aromatization of food, but also in folk medicine, because of their many different biological properties, including antimicrobial properties.

The aim of this study was to determine the antimicrobial activity of volatile phase of different Mediterranean and exotic essential oils and their active volatile components.

The largest, natural habitat of *L. pneumophila* is freshwater environment and very often can be present in various artificial and natural water systems such as cooling towers, air conditioners, humidifiers and dehumidifiers, boilers, swimming pools, fountains and other water systems.

Essential oils are composed of volatile molecules and are produced from aromatic or medicinal plants by a process of secondary metabolism.

Essential oils have been used for thousands of years, not only as ingredients of perfumes or for the aromatization of food, but also in folk medicine, because of their many different biological properties, including antimicrobial properties.

MATERIALS AND METHODS

Model organism

Model organism used in experiments was *L. pneumophila* serogroup 1, strain 130b or ATCC BAA-74 (clinical isolate). This bacterial strain was obtained from the collection of the Department of Microbiology and Parasitology, University of Rijeka. The bacterium was kept in 10% glycerol broth, stored in a freezer at -80 °C. After cultivation for 3–5 days on buffered charcoal yeast extract (BCYE) agar (Oxoid, Altrincham, UK) at 35 ± 2 °C in an aerobic atmosphere, the bacterium was used in experiments.

Essential oils and their basic chemical components

The essential oils were obtained from the company Dea Flores d.o.o. (Rijeka, Croatia) and a stock solution of essential oil was prepared using DMSO (dimethylsulfoxide) (Kemika, Zagreb, Croatia) in a concentration of 100 mg/mL. The tested oils were divided into two groups. The first group consists of exotic essential oils: Spikenard (*Nardostachys jatamansi*), Niaouli (*Melaleuca quinquenervia*), Hyssop (*Hyssopus officinalis*), Palmarosa (*Cymbopogon martinii*) and Ravensara (*Ravensara aromatica*). The second group consists of essential oils from coastal region of Croatia: Juniper berry (*Juniperus communis*), Immortelle (*Helichrysum italicum*), Sage (*Salvia officinalis* L.), Lavander (*Lavandula x hybrida*) and Rosemary (*Rosmarinus officinalis*). The antibacterial activity of bioactive components of the essential oils α -pinene, β -pinene, γ -terpinene and eugenol (Sigma-Aldrich, MO, USA) were also tested. The concentration of the stock suspension was 200 mg/mL.

Preparation of bacterial inoculum

The number of bacteria in bacterial suspension was determined spectrophotometrically. The optical density of bacterial suspension was set to OD₆₀₀ value 1,0 which indicates 10⁹ CFU/mL. The bacterial concentrations of approximately 1 × 10⁸ CFU/mL were used in the experiments. The number of bacteria in the inoculum was confirmed by cultivation of ten-fold dilutions on BCYE medium.

Determination of the antimicrobial activity of volatile components by sealed plate method

Modified version of sealed plate method has been used [6]. Briefly, using a sterile swab, bacteria suspension (10⁸ CFU/mL) were spread evenly onto pre-warmed 37 °C BCYE agar plates. A 5 mm diameter cellulose disc soaked with 5 μ L of the test essential oil or tested components was then placed in the center of each lid of the Petri dish. To set up the assay the bottom of plate containing the bacteria was inverted over the plate carrying volatile compound and was sealed against each other tightly using petri-seal to prevent the essential oil from evaporating outside the Petri dish. The BCYE agar were then placed in a thermostat at 35 ± 2 °C for a 72-hour incubation. Plates without volatiles served as control. After a 72-hour incubation, the




Figure 1.

The Petri dish with cellulose disc soaked with essential oil (left) and zone of inhibition made by volatile components of essential oil (right).

results were obtained by measuring the diameter of the zone of inhibition below the disc caused by the inhibitory action of the volatile components of the essential oils on *Legionella*. For each oil, the diameter of the inhibition zone was measured, and the result was expressed in millimeters. A pure DMSO control was included with each test to ensure that microbial growth was not inhibited by DMSO itself.

Statistical analysis

All experiments were performed three times in duplicate. The results are expressed as the mean \pm standard deviation. Results were statistically analyzed with Tibco Statistic 13.5.0. at a significance level of $p < 0.05$. The normality of the data distribution was tested by the Kolmogorov-Smirnov test. The distribution was normal and parametric T-test was used.

RESULTS AND DISCUSSION

The antimicrobial properties of the volatile components of the exotic essential oils Spikenard, Niaouli, Hyssop, Palmarosa and Ravensara were investigated, and the results are shown in Figure 2. All tested essential oils showed anti-*Legionella* activity. The best result was obtained with the Niaouli essential oil with inhibition zone of 33.3 mm, while the lowest result of *Legionella* inhibition was measured with Spikenard essential oil (16.83 mm). In the case of essential oils from the coastal region of Croatia (Figure 3, Table 1), the inhibition zone was statistically higher from those of exotic essential oils. The strongest antimicrobial activity of volatile components was shown by the immortelle (*H. italicum*) essential oil with an average zone of inhibition of 55.5 mm. Sage (*S. officinalis* L.) and lavender (*L. hybrida* L.) essential oils showed a weaker inhibitory activity with a zone of inhibition of 25.3 mm and 24.0 mm, respectively (Figure 2, Table 1).

The chemical components of the essential oils were also tested separately, and it was examined whether dilution affects the antimicrobial activity of the volatile phase (Figure 4, Table 1). The antimicrobial effect of two concentrations, 100 mg/mL and 200 mg/mL was investigated. The largest zone of inhibition was caused by α -pinene, whose zone of inhibition was almost the entire surface of the Petri dish (concentration 200 mg/mL). γ -terpinene showed no antibacterial effect,

All tested essential oils showed anti-*Legionella* activity. The best result was obtained with the Niaouli essential oil with inhibition zone of 33.3 mm, while the lowest result of *Legionella* inhibition was measured with Spikenard essential oil (16.83 mm).

Figure 2.

Antimicrobial activity of volatile components of an exotic essential oil at concentrations of 200 mg/mL. The results are expressed as a mean value \pm SD of a minimum of three repeated experiments.




Figure 3.

Antimicrobial activity of volatile essential oils of coastal region of Croatia at concentrations of 200 mg/mL. The results are expressed as a mean value \pm SD of a minimum of three repeated experiments.





Figure 4.

Antimicrobial activity of volatile active components at concentrations of 200 mg/mL and 100 mg/ml. The results are expressed as a mean value \pm SD of a minimum of three repeated experiments. Lowercase letters on top shows statistical significance ($p < 0.05$).



and we did not detect a zone of inhibition. Eugenol and β -pinene showed a statistically weaker effect than α -pinene, with a higher concentration of the test compound (concentration 200 mg/mL) showing a better antibacterial effect.

Since essential oils are complex mixtures of many phytochemicals, it is very difficult to determine which of them is responsible for the antimicrobial activity.

Since essential oils are complex mixtures of many phytochemicals, it is very difficult to determine which of them is responsible for the antimicrobial activity. Furthermore, it is also important to know ratios of individual components present in essential oil as well as their synergistic or antagonistic activity. Numerous studies have shown the antimicrobial properties of various essential oils on bacteria, however in most studies essential oils have been added to the medium in which the test was performed. The higher susceptibility of Gram-positive bacteria when

Table 1. Antimicrobial activity of volatile phase of essential oils and active components

Essential oils	Zone of inhibition (mm)±SD
Exotic essential oils	
Nard	16,83±4,19
Niauli	33,27±3,47
Isop	30,27±4,29
Palmerosa	25,27±3,06
Revensara	25,27±1,84
Mediterranean essential oils	
Juniper	47,00±7,35
Immortelle	55,83±4,90
Sage	23,00±2,16
Lavander	25,33±1,25
Rosemary	42,33±2,05
Volatile components	
α-pinene (200 mg/mL)	70,50±3,19
β-pinene (200 mg/mL)	25,83±4,37
γ-terpinene (200 mg/mL)	0,00
eugenol (200 mg/mL)	43,77±0,61
α-pinene (100 mg/mL)	61,60±0,83
β-pinene (100 mg/mL)	18,00±1,63
γ-terpinene (100 mg/mL)	0,00
eugenol (100 mg/mL)	29,67±2,05

compared with Gram-negative strains was found, and a large variability among essential oils in the antibacterial potential has also been observed [7-10].

The main advantage of essential oils is that they do not enhance antibiotic resistance with the long-term use what is the case for synthetic antibiotics, and they showed synergy in antimicrobial activity with conventional antibiotics [11-13]. For this reason, majority of studies carried out on extracts and pure components isolated from essential oils analyzing their antibacterial activities.

Volatile Organic Compounds (VOCs) emitted by plants or essential oils are largely lipophilic products with molecular masses under 300 Da. The vast majority are isoprenoids, including hemiterpenes (C₅H₈) such as isoprene, monoterpenes (C₁₀H₁₆), irregular acyclic homoterpenes (C₁₁H₁₈ or C₁₆H₂₆), and sesquiterpenes (C₁₅H₂₄) [14].

Research into the antimicrobial properties of volatile components of essential oils is rare. Laird and Phillips showed that the use of essential oil vapors in food relates to their antimicrobial activity against food pathogens and food spoilage microorganisms also [15]. Advantage of application of essential oil vapor phase, is that the components are dispersed and tend not to affect the organoleptic properties of the food like liquid essential oil. A study by Nedorostova *et al.* (2009) showed some antibacterial activity of 27 different essential oils vapors against five foodborne pathogens (*Escherichia coli*, *Listeria monocytogenes*, *Salmonella enterica ssp. enteritidis*, *Staphylococcus aureus* and *Pseudomonas aeruginosa*) using the disc volatilization method [16].

The main advantage of essential oils is that they do not enhance antibiotic resistance with the long-term use what is the case for synthetic antibiotics, and they showed synergy in antimicrobial activity with conventional antibiotics.

Advantage of application of essential oil vapor phase, is that the components are dispersed and tend not to affect the organoleptic properties of the food like liquid essential oil.

Our results showed that α -pinene showed a stronger effect on *Legionella* growth inhibitions.

All tested essential oils have the great potential to be used as a volatile antibacterial agent to control legionellosis and could be used as air purifiers in various rooms in hotels and spas.

Exotic essential oil with highest anti-*Legionella* activity was Niaouli essential oil, and the best anti-*Legionella* activity among Mediterranean essential oils have Immortelle essential oil.

The antimicrobial activity of essential oils on *Legionella* has been poorly investigated. Chang *et al.* examined the antimicrobial activity of essential oils extracted from *Cinnamomum osmophloeum* leaves and *Cryptomeria japonica* and its major constituent, cinnamaldehyde, possess strong anti-*Legionella* activities [17]. We have not found data on the antimicrobial activity of volatile phase of essential oils on *Legionella*. Since α -pinene shows high anti-*Legionella* activity one can assume that essential oil with high proportion of this component will give such activity. Ramanoelina *et al.* demonstrate that essential oil of Niaouli (*M. quinquenervia*) have high α -pinene [18] content and it can be cause of high anti-*Legionella* activity as is shown in figure 2. Research conducted by Han *et al.* shows that Immortelle essential oil also have high percentage of α -pinene [19]. Our research has shown a strong antimicrobial activity of volatile components of essential oils that predominantly have a α -pinene in their composition. Alpha-pinene is the most widely encountered terpenoid in nature and is highly repellent to insects. There are two structural isomers of pinene found in nature: α -pinene and β -pinene [20]. The antimicrobial activities of the isomers of pinene was shown against different bacterial and fungal species [20, 21].

Our results showed that α -pinene showed a stronger effect on *Legionella* growth inhibitions. Reason why γ -terpinene did not show any anti-*Legionella* activity could be that this compound on its own has no antimicrobial activity. All tested essential oils have the great potential to be used as a volatile antibacterial agent to control legionellosis and could be used as air purifiers in various rooms in hotels and spas. Furthermore, the potentially synergistic effect of volatile components of different combinations of essential oils remains to be investigated.

CONCLUSION

Wide use of essential oils demands a scientific background so antimicrobial activity of essential oil is examined all over the world. Exotic essential oil with highest anti-*Legionella* activity was Niaouli essential oil, and the best anti-*Legionella* activity among Mediterranean essential oils have Immortelle essential oil. Chemical compound that shows significant highest anti-*Legionella* activity was α -pinene and essential oils with strongest anti-*Legionella* activity are timely the ones with highest proportion of this compound. Essential oils show great benefits as antimicrobial agents and different approach to their use for this purpose is to be examined.

Acknowledgments

The described research was funded by the University of Rijeka grant uniri-biomed-18-171.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

REFERENCES

- [1] Fraiser Dw. *et al.* Legionnaires disease description of epidemic pneumonia. *NEJM*.1997;22:1189-97.
- [2] Benitez AJ., Winchell JM. Clinical application of a multiplex Real-Time PCR assay for simultaneous detection of Legionella species. Legionella pneumophila, and Legionella pneumophila serogroup 1. *JCM*. 2013; 348-51.
- [3] Bartam J. *et al.* Legionella and prevention of Legionellosis. WHO. 2007.
- [4] Kerbel W. *et al.* Recognition, Evaluation, and control of Legionella in building water system. Falls Church, VA, USA: industrial Hygiene Association. 2015.
- [5] Winska K. *et al.* Essential oils as antimicrobial agents – Myth or real alternative? *Molecules*. 2019;24(11):2130.
- [6] Lopez P. *et al.* Solid and vapor-phase antimicrobial activities of six essential oils: susceptibiliti of selectrd foodborne bacterial and fungal strains. *J. Agric. Food Chem*. 2005;53:6939-46.
- [7] Murbach Tele Andrade BF. *et al.* Antimicrobial activity of essential oils. *JEOR*. 2013;26:34-40.
- [8] Malenica Staver M. *et al.* In vitro antiproliferative and antimicrobial activityof the essential oil from the flowers and leavesof helichrysum italicum (roth) g. Don growing incentral dalmatia (Croatia). *JEOBP*. 2018;21:77-91.
- [9] Inouye S., Takizawa T., Yamagucchi H. Antibacterial activity of essential oils and their major constituents against respiratory tract pathogens by gaseous contact. *J. Antimicrob. Chemother*. 2017;47(5):565-73.
- [10] Shojee-Aliabdi S., Marzieh Hosseini S., Mirmoghtadaie L. Antimicrobial activity of essential oil. *Essential oils in food procesing: Chemistry, Safety and Applications*. 2017;191-229.
- [11] Aleksic V. *et al.* Synergistic effect of Myrtus communis L. essential oils and conventional antibiotics against multi-drug resistant Acinetobacter baumannii wound isolates. *Phytomedicine*. 2014;12(21):1666-74.
- [12] Knezevic P. *et al.* Antimicrobial activity of Eucalyptus camaldulensis essential oils and their interactions with conventional antimicrobial agents against multi-drug resistant Acinetobacter baumannii. *J Ethnopharmacol*. 2016;178:125-36.
- [13] Duarte A. *et al.* Synergistic activity of coriander oil and conventional antibiotics against Acinetobacter baumannii. *Phytomedicine*. 2012: 19:236-8.
- [14] Antonelli M. *et al.* Forest volatile organic compounds and their effects on human health: a state-of-the-art review. *Int. J. Environ. Res. Public Health*. 2020;17(18):6506.
- [15] Laird K., Phillips C. Vapour phase: a potential future use for essential oils asantimicrobials? *Lett. Appl. Microbiol*. 2012;3(54):169-74.
- [16] Nedorstova L. *et al.* Antimicrobial properties of selected essential oils in vapuor phase against foodborne bacteria. *Food control*. 2009;20:157-60.
- [17] Chang CW. *et al.* Antibacterial activities of plant essential oils against Legionella pneumophila. *Wather Res*. 2008;42:278-86.
- [18] Ramanoelina PA., Bianchini JP., Andriantsiferana M. Chemical Composition of Niaouli Essential Oils from Madagascar. *J. Essent. Oil Res*. 1992; 657-8.
- [19] Han X. *et al.* Chemical composition analysis and in vitro biological activities of ten essential oils in human skin cells. *Biochim. Open*. 2017;5:1-7.
- [20] Rivas da Silva AC. *et al.* Biological activities of a-pinene and β-pinene enantiomers. *Molecules*. 2012;17:6305-16.
- [21] de Sousa EL. *et al.* Antibacterial activity and time-kill kinetics of positive enantiomer of α-pinene against strains of staphylococcus aureus and escherichia coli. *Curr Top Med Chem*. 2018; 18(11)917-24.