

Biljni lektini u prevenciji i liječenju raka

Giacometti, Jasminka

Source / Izvornik: **Medicina Fluminensis : Medicina Fluminensis, 2015, 51, 211 - 229**

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

Permanent link / Trajna poveznica: <https://urn.nsk.hr/urn:nbn:hr:184:052288>

Rights / Prava: [In copyright](#)/[Zaštićeno autorskim pravom.](#)

Download date / Datum preuzimanja: **2025-02-27**



Repository / Repozitorij:

[Repository of the University of Rijeka, Faculty of Medicine - FMRI Repository](#)



Plant lectins in cancer prevention and treatment

Biljni lektini u prevenciji i liječenju raka

Jasminka Giacometti

Abstract. Plant lectins are specific carbohydrate-binding proteins that are widely distributed in various plant species. They participate in many physiological processes and are capable of modulating the immune response. Recently, greater attention has been drawn to their remarkable anticancer properties. Lectins are associated with cell adhesion, cell proliferation and induction of apoptosis. However, some of the questions related to the molecular mechanism / metabolic pathways and biological effects of lectins are still open. New challenges in the research of lectins are related to their application in nanotechnology and development of glycoproteomics. In addition, glycoproteomics is a powerful tool in the characterization of lectins and will be indispensable for development of lectin based drugs in the near future. This review provides a brief outline of the up-to-date advances in the field of plant lectins, focusing on their complex mechanisms implicated in apoptosis and autophagy. The current applications in cancer treatment are also described.

Key words: anticancer agents; lectins; plant lectins

Sažetak. Biljni lektini su specifični glikoproteini koji su široko rasprostranjeni u različitim biljnim vrstama. Sudjeluju u mnogim fiziološkim procesima gdje moduliraju imunološki odgovor. Velik interes za ovu skupinu proteina javio se zbog njihove sposobnosti da utječu na protutumorsku aktivnost putem stanične adhezije, stanične proliferacije i indukcije apoptoze. No još su uvijek nerazjašnjena neka pitanja koja se odnose na molekularni mehanizam / metaboličke puteve i biološke učinke lektina. Novi izazovi u istraživanju lektina odnose se na njihovu primjenu u nanotehnologiji te razvoju glikoproteomike. Glikoproteomika je jedan od moćnih alata u karakterizaciji lektina, a u bliskoj budućnosti i neizostavni alat u razvoju lijekova koji se temelje na lektinima. Ovaj pregledni rad ukratko opisuje izvore, strukturu i primjenu biljnih lektina, povezujući njihove složene mehanizme djelovanja u apoptozu i autofagiju. Opisana je primjena lektina u terapiji raka.

Ključne riječi: biljni lektini; lektini; protutumorski spojevi

Department of Biotechnology,
University of Rijeka, Rijeka

Primljeno: 24. 2. 2015.

Prihvaćeno: 11. 3. 2015.

Corresponding author:

Prof. dr. sc. Jasminka Giacometti
University of Rijeka,
Department of Biotechnology,
R. Matejčić 2, HR-51 000 Rijeka, Hrvatska
e-mail: jgiacometti@biotech.uniri.hr

<http://hrcak.srce.hr/medicina>

ABBREVIATIONS

ATG families – autophagic family proteins;
Bcl-2 – apoptosis regulator protein;
BNIP3 – BCL2/adenovirus E1B 19 kDa protein -interacting protein 3;
ERK – extracellular signal-regulated kinase;
Gal – galactose;
GalNAc – N-acetylgalactosamine;
GlcNAc – N-acetylglucosamine;
HILIC – hydrophilic interaction liquid chromatography;
LC-MS/MS – liquid chromatography-tandem mass spectrometry;
MALDI TOF – time-of-flight mass spectrometer with matrix-assisted laser desorption/ionization;
MALDI TOF/TOF – tandem time-of-flight mass spectrometer systems with MALDI;
Man – mannose;
MRM MS – Multiple Reactions Monitoring Mass Spectrometry;
PI3K/Akt pathway – intracellular signaling pathway;
PNGase – peptide-N4-(N-acetyl-beta-glucosaminyl)asparagine amidase;
PNGase F – amidase that cleaves between the innermost GlcNAc and asparagine residues of high mannose, hybrid, and complex oligosaccharides from N-linked glycoproteins;
Ras/Raf – signal transduction pathway;
Sia – sialic acid.


INTRODUCTION

Lectins are a complex and heterogeneous group of non-enzymatically carbohydrate-binding proteins that specifically recognize and bind reversibly to specific mono- and oligosaccharides on cell surfaces, the extracellular matrix, and secreted glycoproteins. More than a hundred of these molecules have been isolated from plants, viruses, bacteria, invertebrates and vertebrates, including mammals. They bind carbohydrates and possess the capability to agglutinate cells or precipitate polysaccharides and glycoconjugates. Lectins are a component of traditional herbs such as dietary and medicinal plants.

Various plants contain different plant lectins related to their molecular specificity. Plant lectins can be classified into three groups based on: i) their overall mature structure, ii) different families, according to some common features, and iii) diversity of carbohydrate-binding specificities. Differences in lectin structure and carbohydrate specificity are related to their different functions¹. Depending on carbohydrate specificity, major lectins are divided into mannose binding lectins, galactose/*N*-acetylgalactosamine binding lectins, *N*-acetylglucosamine binding lectins, *N*-acetylneuraminic acid binding lectins and fucose binding lectins as shown in Figure 1. However,

Table 1. Overview of the plant carbohydrate-binding motifs according to Ref. 1.

Lectin domain	Carbohydrate specificity	Examples
<i>Agaricus bisporus</i> agglutinin domain	T-antigen	ABA, MarpoABA
Amaranthins	T-antigen	Amaranthin, HFR2
Class V chitinase homologs	Blood group B, high-man N-glycans	RobpsCRP
Cyanovirin domain	High-man N-glycans	CV-N
<i>Euonymus europaeus</i> agglutinin domain	Blood group B, high-man N-glycans	EEA
<i>Galanthus nivalis</i> agglutinin domain	Man, oligomannosides, high-man N-glycans, complex, N-glycans	GNA, ASA II, ASAL, ACA, LOA
Hevein domain	Chitin, high-man, Man, N-glycans	Hevein, UDA, WGA, HFR3
Jacalins	Gal, T-antigen, Man, N-glycans	Jacalin, Heltuba, HFR1
Legume lectin domain	Man/Glc, Gal/GalNAc, (GlcNAc) _n , Fuc, Sial2, 3Gal/GalNAc, complex N-glycans	PHA, ConA, Gleheda, PSA, GSII
LysM domain	Chitin-oligosaccharides	LysM, CEBiP
<i>Nicotiana tabacum</i> agglutinin domain	GlcNAc-oligomers, high-man N-glycans	NICTABA, PP2
Ricin-B domain	Gal/GalNAc, Sial2-6Gal/GalNAc	Ricin, SNA-I



genome/transcriptome analyses revealed that plant lectins can be classified into twelve distinct families of evolutionary and structurally related lectin domains which are presented in Table 1.

Many of the characterized plant lectins interact with monosaccharides, but their affinity for simple sugars is lower in comparison with more complex carbohydrate structures such as *N*-glycans². The *Leguminosae* is the best-characterized family of plant lectins. This family includes lectins such as Concanavalin A (ConA), soybean agglutinin (SBA), and lentil lectin. Two other smaller families of plants whose lectins have been characterized are the *Gramineae* (cereals, such as wheat germ) and *Solanaceae* (potatoes and tomatoes).

One example of their mature structure is ConA, as a tetrameric protein which binds specifically α -D-mannosyl, and α -D-glucosyl residues. Another example is peanut (*Arachis hypogaea*) agglutinin, which is homotetrameric non-glycosylated protein (without RIP activity) and shows specificity for the tumor-associated T-antigenic disaccharide Gal β 1, 3GalNAc. Ricin is classified as both lectin and type II ribosome inactivating protein (RIP) and consists of two parts; an A chain (with *N*-glycosidase activity/RIP activity) and a B chain (hemagglutinating/lectin activity) with the B chain capable of binding different carbohydrates, such as β -D-glucose and β -D-galactose. The list of origin of lectins and their abbreviations is presented in Table 2.

Due to their ability to distinguish carbohydrates in human blood cells, different specific lectins can be used in blood typing to differentiate between blood types⁴.

One of the earliest findings was related to the biological role of some lectins as cell surface sugars

- Plant lectins seem to have great potential as anti-cancer therapeutic agents.
- Lectins can cause cancer cell agglutination and/or aggregation and blocking of further migration.
- Some plant lectins are capable of modulating the immune response in different ways.
- Toxic plant lectins can be used as a supportive therapy to improve health-related quality of life (HRQoL).
- Glycoproteomics is one of the tools in characterizing lectins and the development of lectin based drugs.

and its mitogenic stimulation on the surface of the lymphocytes. This property makes their glycosylation useful tools in cancer research, especially for the isolation and characterization of polysaccharides and glycoconjugates as diagnostic tools for the investigation of early cell-membrane alterations and carbohydrate changes that accompany neoplastic processes, and in immunological studies⁵.

In the past, numerous lectins were isolated from plants as well as from microorganisms and animals, however, their structure and function as recognition molecules in cell-molecule and cell-cell interactions in various biological systems, have been established during the past two decades.

Despite the disparity in physicochemical and biochemical characteristics, lectins from different sources exhibit common biological activity. They are involved in the strategies of different scientific and practical fields such as agricultural, agro-economy, food production and food science, life

Table 2. List of origin of some lectins

Abbreviations	Name of Lectin	Origin (eng.)	Origin (lat.)
AAL	<i>Aleuria Aurantia</i> Lectin	Orange Peel Fungus	<i>Aleuria aurantia</i>
ABA	<i>Agaricus bisporus</i> agglutinin	Edible mushroom, white button mushroom	<i>Agaricus bisporus</i>
Abrin A	Abrin A	Crab's Eye	<i>Abrus precatorius</i>
ACA	<i>Amaranthus caudatus</i> agglutinin	Amaranth	<i>Amaranthus caudatus</i>
AGG	Gamma-globulin		
AML	<i>Astragalus membranaceus</i> lectin	Huáng qí	<i>Astragalus membranaceus</i>
AMML	<i>Astragalus membranaceus</i> var. Mongholicus lectin	Milk vetch / Huang qi	<i>Astragalus mongholicus</i>
ASAL	<i>Allium sativum</i> leaf agglutinin	Garlic	<i>Allium sativum</i>
ASA I	<i>Allium sativum</i> bulb agglutinin I	Garlic	<i>Allium sativum</i>
ASA II	<i>Allium sativum</i> bulb agglutinin II	Garlic	<i>Allium sativum</i>
CEBiP	Chitin elicitor binding protein	Rice	<i>Oryza sativa</i>
CMA	Chelidonium majus agglutinin	Greater celandine	<i>Chelidonium majus</i>
CML	<i>Cratylia mollis</i> lectin	Cratylia	<i>Cratylia mollis</i>
Con A	Concanavalin A	Jack bean	<i>Canavalia ensiformis</i>
CV-N	Cyanovirin-N	Cyanobacterium Nostoc ellipsosporum	<i>Nostoc ellipsosporum</i>
EEA	<i>Euonymus europaeus</i> agglutinin	European spindle	<i>Euonymus europaeus</i>
Gleheda	<i>Glechoma hederacea</i> agglutinin	Ground-ivy	<i>Glechoma hederacea</i>
GNA	<i>Galanthus nivalis</i> agglutinin	Snowdrop	<i>Galanthus nivalis</i>
GS-II	<i>Griffonia simplicifolia</i> agglutinin II	Griffonia seed	<i>Griffonia simplicifolia</i>
GSA-IA4	<i>Griffonia simplicifolia</i> agglutinin	Griffonia seed	<i>Griffonia simplicifolia</i> , <i>Griffonia (Bandeiraea) simplicifolia</i>
Heltuba	<i>Helianthus tuberosus</i> agglutinin	Jerusalem artichoke	<i>Helianthus tuberosus</i>
HFR1	Hessian fly responsive 1	Hessian fly	<i>Mayetiola destructor</i>
HFR2	Hessian fly responsive 2	Hessian fly	<i>Mayetiola destructor</i>
HFR3	Hessian fly responsive 3	Hessian fly	<i>Mayetiola destructor</i>
HEV1	Hevein	Pará rubber tree	<i>Hevea brasiliensis</i>
JAC	Jack fruit lectin, <i>Artocarpus integrifolia</i> agglutinin, Jacalin	Jack fruit	<i>Artocarpus integrifolia</i>
Jacalin	Jacalin	Jackfruit	<i>Artocarpus integrifolia</i>
Lentil	Lentil lectin	Lentil	<i>Lens culinaris</i>
LCA	<i>Lens culinaris</i> agglutinin	Lentil	<i>Lens culinaris</i>
LOA	<i>Listera ovata</i> agglutinin	Twayblade	<i>Listera ovata</i>
LysM	LysM domain		
MAL	<i>Maackia amurensis</i> agglutinin	Amur maackia	<i>Maackia amurensis</i>
MarpoABA	<i>Agaricus bisporus</i> agglutinin homolog	Edible mushroom, white button mushroom	<i>Agaricus bisporus</i>
ML-I	Mistletoe lectin I	Mistletoe	<i>Viscum album</i> L.
ML-II	Mistletoe lectin II	Mistletoe	<i>Viscum album</i> L.
ML-III	Mistletoe lectin III	Mistletoe	<i>Viscum album</i> L.
MLL	Mulberry leaf lectin	White mulberry	<i>Morus alba</i>
Nictaba	<i>Nicotiana tabacum</i> agglutinin	Tobacco	<i>Nicotiana tabacum</i>
NPA	<i>Narcissus pseudonarcissus</i> agglutinin	Common daffodil	<i>Narcissus pseudonarcissus</i>
PCL	<i>Pleurotus citrinopileatus</i> lectin	Citrinopileatus	<i>Pleurotus citrinopileatus</i>
PHA	Phytohemagglutinin, <i>Phaseolus vulgaris</i> leucoagglutinin	Common bean, kidney bean	<i>Phaseolus vulgaris</i> L.
PNA	Peanut agglutinin	Peanut	<i>Arachis hypogaea</i>
POL	<i>Pleurotus ostreatus</i> lectin	Mushroom	<i>Agaricus bisporus</i>
PP2	<i>Nicotiana tabacum</i> agglutinin domain	Tobacco	<i>Nicotiana tabacum</i>

Abbreviations	Name of Lectin	Origin (eng.)	Origin (lat.)
PSA	<i>Pisum sativum</i> agglutinin	Pea	<i>Pisum sativum</i>
RBA	Rice bran agglutinin	Rice bran	<i>Oryza sativa</i> L.
Ricin (RCA)	<i>Ricinus communis</i> , Anti-B4-blocked ricin	Castor bean	<i>Ricinus communis</i>
Ricin A	<i>Ricinus communis</i> , Anti-B4-blocked ricin	Castor bean	<i>Ricinus communis</i>
RLL	<i>Russula lepida</i> lectin	<i>Russula rosea</i>	<i>Russula lepida</i>
rML	Recombinant mistletoe lectin	Mistletoe	<i>Viscum album</i> L.
RobpsCRA	Chitinase-related agglutinin homolog	black locust	<i>Robinia pseudoacacia</i>
Saracin	<i>Saraca indica</i> lectin	Ashoka	<i>Saraca indica</i>
SBA	Soybean agglutinin	soybean	<i>Glycine max</i>
SBL	Soybean lectin	soybean	<i>Glycine max</i>
SNA	<i>Sambucus nigra</i> agglutinin	Elderberry, European black elderberry	<i>Sambucus nigra</i>
STL	<i>Solanum tuberosum</i> lectin	Potato	<i>Solanum tuberosum</i>
TML	<i>Tricholoma mongolicum</i> lectin	Paimo mushroom	<i>Tricholoma mongolicum</i>
UDA	<i>Urtica dioica</i> agglutinin	Stinging nettle	<i>Urtica dioica</i>
VAA	<i>Viscum album</i> agglutinin	mistletoe	<i>Viscum album</i>
VCA	<i>Viscum album coloratum</i> agglutinin	Korean mistletoe	<i>Viscum album</i> L. var. <i>coloratum</i>
VFA	<i>Vicia faba</i> agglutinin, Broad bean lectin	Broad bean, fava bean	<i>Vicia faba</i>
WGA	Wheat germ agglutinin	Wheat	<i>Triticum aestivum</i> L.

science, health science, pharmaceuticals, etc. Many plant lectins showed anticancer properties *in vivo*, and *in vitro*, thus they have a potential for use as a therapeutic agent in the malignant neoplastic disease treatment. This review discussed today's main research interest of plant lectins as the next generation of anticancer drugs.

PRODUCTION, PURIFICATION AND PROTEOMIC APPLICATION OF LECTINS

Two main ways, enabling the production of lectins, a) isolation from their natural sources by chromatographic procedures, or b) production by recombinant DNA technology as shown in Table 3. The yields of animal lectins are usually low in comparison to the yield of plant lectins such as legume lectins⁶.

Isolation of lectins integrates different purification techniques, such as precipitation (using acids, organic solvents and salts) and chromatographic methods such as affinity chromatography (AC), ion-exchange chromatography (IEX), hydrophobic interaction chromatography (HIC) and gel permeation (GF)⁷. Lectin affinity chromatography (LAC), often uses the immobilized lectins such as ConA in separation and isolation of glycopeptides

that express *N*-linked structures and high-mannose glycans⁷⁻¹¹.

Recombinant DNA technology has been used for cloning and characterizing newly pure and sequence-defined lectins. Recombinant lectins are often produced in *Escherichia coli* and by post-translational modified recombinant lectins are produced in eukaryotic organisms. These recombinant lectins may have different applications such as *i*) in cancer diagnostics and/or therapy, *ii*) antimicrobial, antiviral and anti-insect agents, or *iii*) in microarray for glycome profiling¹². Although recombinant lectins can be synthesized, due to high costs and low yield, more acceptable production is isolated from plant sources. This especially applies to the isolation of high-yield *Phaseolus* cultivars cvs. French bean 35 (Table 3).

Current methods to quantify lectin levels in foods and other matrix analyses are based on immunosorbent assay (ELISA), which mainly relies on specific monoclonal antibody or pre-labelled well-known lectins, or toxicity tests. Wang et al.¹⁴ suggested a new strategy to detect the specific carbohydrate binding capability of lectins based on enzyme-linked adsorbent assay applying different monosaccharide-polyacrylamide conjunc-

Table 3. Yields of animal and plant lectins (from different *Phaseolus* cultivars) obtained by chromatographic isolation and plant lectins produced by recombinant DNA techniques modified according to Ref. 13.

Plant lectins from <i>Phaseolus</i> cultivar (yield mg/100 g seed)				
Source	Chromatographic purification	Yield	Sugar specificity	Reference
Anasazi bean	Affi-gel blue gel, Mono S and Superdex 200	13	N.F.	82
Dark red kidney bean	DEAE-cellulose and Affi-gel blue gel	107	N.F.	83
Escumite bean	AC	163	N-acetyllactosamine-type glycans	84
Extralong autumn purple bean	Blue-Sepharose, Q-Sepharose, Mono Q and Superdex 75	35	Galactose	85
French bean 12	SP-Sepharose, Affi-gel blue, Q-Sepharose, and Superdex 200	4.8	N.F.	86
French bean 35	Blue-Sepharose, Q-Sepharose and Superdex 75	1100	N.F.	87
Red kidney bean	Affi-gel blue gel and CM-Sepharose	27.5	Lactoferrin, ovalbumin, thyroglobulin	88
Animal lectins				
<i>Acropora millepora</i> (coral) plasma fluid	Mannose AC	0.7 mg/100 ml plasma	N.F.	89
<i>Aristichthys nobilis</i> (bighead carp) gills	DEAE-Sepharose, Sephacryl S-200 and Superdex 200	9.4 mg/100 g	N.F.	90
<i>Bubalus bubalis</i> (Buffalo) heart tissue	Ammonium sulfate precipitation and Sephadex G50	0.97 mg/100 g	N.F.	91
<i>Capra hircus</i> (goat) heart tissue	Ammonium sulfate precipitation and Sephadex G50, Lactosyl-Sepharose 4B AC	1.09 mg/ 100 g	Galactose	92
<i>Holothuria scabra</i> (sea cucumber) coelomic fluid	Ultrafiltration and Phenyl-Sepharose	1.6 mg/100 ml	N.F.	93
<i>Macoma birmanica</i> (marine bivalve) foot muscles	Ammonium sulfate precipitation and N-acetylglucosamine Sepharose 4B	4.5 mg/100 g	N.F.	94
<i>Nemopilema nomurai</i> (jellyfish)	SP-Sepharose and BSM- Toyopearl	0.35 µg/100 g	N.F.	95
Plant lectins produced by recombinant DNA techniques				
Natural source of lectin	Yield (mg/L culture medium)	Genetically modification in cells		Reference
<i>Allium sativum</i> (garlic) leaf	5	cDNA was cloned into NdeI and BamHI restricted plasmid pET19b and expressed in <i>E. coli</i> strain BL21 (DE3) cells		96
<i>Artocarpus incise</i> (breadfruit)	16	cDNA was cloned into the pET-25b(+) and expressed in <i>E. coli</i> .		97
<i>Artocarpus incise</i> (breadfruit)	18–20	cDNA was cloned into EcoRI/XbaI restricted plasmid pUC57 and expressed in <i>E. Coli</i>		98
<i>Glycine max</i> (Soybean)	0.1	cDNA was cloned NcoI/NdeI/BamHI restricted plasmid PET-3d and expressed in <i>E. coli</i> strain BL21(DE3)pLysS		99
<i>Nicotiana tabacum</i> (tobacco) leaves	6	cDNA was cloned EcoRI/NotI restricted plasmid and expressed in <i>E. coli</i> strain top10F		100
<i>Oryza sativa</i> (rice) roots	14.6	cDNA was cloned into NdeI/BamHI restricted pET 3D plasmid and expressed in <i>E. coli</i> strain BL21 (DE3) cells		101
<i>Pisum sativum</i> (pea)	2–5	cDNA was cloned into HindIII/PstI/BamHI restricted plasmid and expressed in <i>E. coli</i> strain W3110		102
<i>Polyporus squamosus</i> fruiting bodies	4–7	cDNA was cloned into NdeI/BamHI restricted plasmid and expressed in <i>E. coli</i> strain Nova Blue (DE3)		103

N.F. – not found; AC – affinity chromatography




Figure 2. Glycoproteomics based on affinity chromatography at the protein level with immobilized lectin columns modified according to Ref. 7.

gates as capturing agents for screening lectins in biological samples.

In summary, plant lectins have application in LAC, blotting, affinity electrophoresis, immune-electrophoresis as well as in microarrays, as in evanescent-field fluorescence-assisted lectin microarray¹⁵.

Proteomic strategies to quantitative analysis of plant lectins include the use of chromatographic or electrophoretic strategies combined with mass spectrometry (LC-MS/MS, MALDI-TOF MS or MALDI-TOF/TOF MS). The workflow often involves a combination of LAC, tryptic digestion, ion-pairing HILIC, and precursor ion-driven data-dependent MS/MS analysis with a script to facilitate the identification and characterization of occupied *N*-linked glycosylation sites^{16,17} (Figure 2).

Proteomic approach was used in investigation of quantitative differences in aberrant glycosylation of target glycoproteins between noncancerous group and patient group with carcinoma such as adenocarcinoma lung cancer (ADLC)¹⁸, liver cancer¹⁹ developed by cooperatively using comparative lectin-capturing, targeted mass spectrometry (MRM MS), and antibody/lectin sandwich ELISA. This different proteomic approach can be useful for identifying and verifying biomarker candidate involved in aberrant protein glycosylation⁷.

SOME IMPORTANT ANIMAL AND HUMAN LECTINS

Twelve structural families of lectins are known to exist in mammals where carbohydrates bind to another structure such as protein–protein, protein–lipid or protein–nucleic acid. Although they have other functions, their main function is generally related to the recognition molecules within the immune system, as direct first-line defense against pathogens, cell trafficking, immune regulation and prevention of autoimmunity²⁰.

C-type lectins with C-type lectin domain-containing proteins (CTLDs) are characteristic of mammals. Their seven subgroups are based on the order of the various protein domains in each protein³. Changes in the amino acid residues that interact with the carbohydrate alter the carbohydrate-binding specificity of the lectins. A calcium ion bridges the protein and the sugar through direct interactions with sugar hydroxyl groups as shows Figure 3. These proteins function as adhesion and signaling receptors in many immune functions such as inflammation and immunity to tumors and virally infected cells. A large class found in animals includes collectins, selectins, endocytic receptors, and proteoglycans that can play an important role in cellular functions^{21,22}.




Figure 3. Structure of a C-Type, carbohydrate-binding domain from an animal lectin according to Ref. 3.

All selectins are single-chain transmembrane glycoproteins that share similar properties to C-type lectins due to a related amino terminus and calcium-dependent binding on immune-system cells to the sites of injury in the inflammatory response²³. The L, E, and P forms of selectins bind specifically to carbohydrates on lymph-node vessels, the endothelium, or activated blood platelets, respectively. P- and E-selectins are highly expressed on the luminal plasma membrane of

vascular endothelial cells at sites of inflammation, therefore, can be smart targets for the delivery of anti-inflammatory drugs²⁴.

New therapeutic agents that control inflammation may emerge from a detailed understanding of how selectins bind and distinguish different carbohydrates²⁵. P-selectin glycoprotein ligand 1 (PSGL-1) is the only transmembrane glycoprotein characterized at the molecular, cellular and functional levels, and which is comprised of extracellular, transmembrane, and cytoplasmic domains²⁶ (see Figure 4). PSGL-1 is one of the promising selectin inhibitor, which has entered clinical trials^{26,27}.

The ability of viruses to infect specific cell types is partially certain by the ability of these viruses to bind particular structures or receptors on the surfaces of cells. In some cases, these receptors are carbohydrates.

Viral infections often coincide with platelet activation. Increased levels of E-selectin on the endothelial cell surface were found in Dengue virus-infected patients²⁸. In these patients, increased E-selection encouraged enhanced activation of adhesion as well as enhanced activation of the coagulation cascade. The viral protein from influenza virus recognizes sialic acid residues found in cell-surface glycoproteins (hemagglutinin)²⁹. P-selectin is an important adhesion molecule in regulating T cell responses which may be important for T cell memory and immunity to influenza virus³⁰.

Soluble P-selectin levels (sP-selectin) in plasma were higher in hepatitis C patients with low




Figure 4. Scheme of selectins and their common ligand PSGL-1 modified according to Ref. 26.

platelet counts. This result indicates that hepatitis C virus infection (HCV) might be directly responsible for a condition of *in vivo* platelet activation in patients with chronic C hepatitis³¹. Also, low platelet and sP-selectin levels are related to the degree of liver disease and thrombosis in patients with cirrhosis³².

APPLICATIONS OF LECTINS

Plant lectins: possible application in diagnostics and therapy

Due to the large range of natural sources as well as high specificity, lectins are important tools in cell biology and immunology. This part is focused on the anticancer activity of selected plant lectins *in vitro*, *in vivo* and in human case studies. Lectins can penetrate into cells, causing cytotoxicity, apoptosis, cancer cell agglutination and/or aggregation, and inhibition of tumor growth. Several studies demonstrated a strong correlation between specific lectin-binding patterns and their biological effects in various tumors.

Agglutination is primarily done by binding to the glycoprotein receptors on cell membranes, resulting in blocking further migration. Thus, they can affect cancer cells, by modulating the status of the immune system by altering the production of various interleukins, certain protein kinases, and proteins themselves by binding to ribosomes. Plant lectins affect both apoptosis and autophagy by modulating representative signalling pathways involved in Bcl-2 family, caspase family, p53, PI3K/Akt, ERK, BNIP3, Ras-Raf and ATG families³³.

In vitro studies

Although carbohydrates are associated with cell growth and viability, glycosylation also has an integral role in many processes leading to cell death. Glycans, simple or complexed with glycan-binding proteins, can transfer intracellular signals or control extracellular processes and so promote initiation, implementation and resolution of cell death programs.

In vitro studies have shown that plant lectins elicit apoptosis in different cancer cell lines. Plant lectins can modify the cell cycle by inducing non-apoptotic G1-phase accumulation mechanisms

and G2/M phase cell cycle arrest and apoptosis^{34,35}.

This can be explained by the binding between lectin-tumor cells which depend on lectins with different sugar-binding specificities such as galactosyl-specificity of the mushroom *Pleurotus ostreatus* lectin (POL). The specificity was increased by substitution at the C-2 position of the galactosyl residue with a fucosyl or acetylamino group³⁶. Different carbohydrate-binding specificities were studied by Wang et al.³⁷ on human hepatoma (H3B), human choriocarcinoma, mouse melanoma, and rat osteosarcoma cell lines. In comparison to other cells, POL inhibited more sarcoma S-180 cells³⁷.

The fresh oyster mushroom *Pleurotus ostreatus* produced the most significant cytotoxicity on human androgen-independent cancer PC-3 cells among the mushroom species tested³⁸. Three proteoglycan fractions from *P. ostreatus* mycelia were tested for *in vitro* and *in vivo* immunomodulatory and anticancer effects on Sarcoma-180-bearing mouse model. Reduced number of S-180 tumor cells and cell cycle analysis showed that most of the cells were found to be stopped in pre-G0/G1 phase of the cell cycle. Three tested proteoglycan fractions elevated mouse natural killer (NK) cell cytotoxicity and stimulated macrophages to produce nitric oxide³⁹. The mechanism of this anticancer effect may be explained by the improvement of the host immune system.

In comparison to some plant lectins, dietary lectins may not be toxic. Overview of selected plant lectins which is important in cell biology and immunology is shown in Table 4.

Vicia faba agglutinin (VFA) is a dietary lectin, with D-glucose and D-mannose sugar specificity, which is present in broad beans. VFA can alter the proliferation of colon cells by aggregation, stimulation of the morphological differentiation and reduction of the malignant phenotype of human colon cancer cells by acting to direct binding to N-glycosylated epithelial cell adhesion molecule (epCAM) or through a pathway involving epCAM⁴⁰.

Wheat germ agglutinin (WGA) is N-acetylglucosamine binding lectin. Its inhibitory effect is linked to a small decrease in α -amylase secretion

Table 4. Inhibitory effects of plant lectins on malignant cells *in vitro* modified according to Ref. 34.

Lectins	Tumor cells	Type of effect	References
AAL, WGA, MAL, TML, STL	H3B human hepatoma, Jar human choriocarcinoma, and ROS rat osteosarcoma	C/TI	37
LCA, NPA	H3B human hepatoma		
AML	K562 leukemia cell line	C/TI, apoptosis activation of the caspase cascade	104
AMML	Human cervical carcinoma cell line (HeLa)	Apoptosis, Cell cycle arrest at S phase	105
ABA, WGA	LS174T, SW1222, and HT29 human colon cancer	C/TI, CA/A	40
VFA, PNA	SW1222 and HT29 human colon cancer		
VFA	LS174T human colon cancer	C/TI, stimulation of morphological differentiation, reduction of malignant phenotype, CA/A	
Abrin A	Jurkat and CCRF-HSB-2 adult T-cell lymphoblastic leukemia cells	C/TI, CA/A	47
	Molt-4 and HPB-ALL adult T-cell lymphoblastic leukemia cells		
	RPMI 8402 and BALL-1 adult T-cell lymphoblastic leukemia cells	C/TI	
	CCRF-CEM adult T-cell lymphoblastic leukemia	C/TI, CA/A	
	BALM-1 Acute B-cell lymphoblastic leukemia	C/TI, CA/A	
	NALM6 Acute B-cell lymphoblastic leukemia		
ACA, JAC	HT29 human colon cancer	C/TI	106
Anti-CD64 Ricin A	Acute human myeloid leukemia	C/TI, apoptosis	107
CD22-rec Ricin-a	Daudi and Ramos B-cell lines (Burkitt lymphomas)	C/TI	108
CD22-rec Ricin-a	Chronic B-cell lymphocytic leukemia (B-CLL)		
CD22-rec Ricin-a	Acute B-cell lymphoblastic leukemia (B-ALL)	C/TI	
CMA, ConA, LCA, UDA, WGA	Merkel cell skin carcinomas	DC/A/BindCellMemR	109
Con A, GSA-IA4	Hs729 (HTB-153) human rhabdomyosarcoma	C/TI, DC/A/BindCellMemR	110
Con A, GSA-IA4, WGA	SK-UT-1 and SK-LMS-1 human leiomyosarcoma	C/TI, DC/A/BindCellMemR	
PHA	Hs729 (HTB-153) human rhabdomyosarcoma and SK-UT-1 and SK-LMS-1 human leiomyosarcoma	C/TI, DC/A/BindCellMemR	
PNA	SK-UT-1 (HTB-114) human leiomyosarcoma	C/TI, DC/A/BindCellMemR	
	SK-LMS-1 (HTB-88) human leiomyosarcoma	DC/A/BindCellMemR	
WGA	Hs729 (HTB-153) human rhabdomyosarcoma	DC/A/BindCellMemR	
ConA, GSA-IA4, WGA	SK-MEL-28, HT-144, and C32 human melanoma	C/TI	111
PHA	SK-MEL-28, HT-144 and C32 human melanoma	C/TI	
JAC, WGA	Adenomatous polyps and colorectal neoplasms	DC/A/BindCellMemR	112
ML-I, ML-II, ML-III	Molt-4 human lymphocyte	C/TI	52
ML-I	Molt-4 human lymphocyte	Ribosome binding/inhibition of protein synthesis, DC/A/BindCellMemR, internalization of lectin, apoptosis	53
ML-II	U937 human monoblastic leukemia	Apoptosis, activation of extracellular signal-regulated kinases, activation of p38 mitogen-activated protein kinase, alteration of cellular signaling pathways	54
	U937 human myeloleukemic	Apoptosis, activation of the caspase cascade	55
	Jurkat T, RAW 264.7, HL-60, DLD-1, primary acute myelocytic leukemic	Apoptosis	56

Lectins	Tumor cells	Type of effect	References
ML-I	Malignant melanoma	C/TI	58
ML-II, ML-III		C/TI	
MLL	MCF-7 human breast cancer cells	C/TI, stimulation of morphological differentiation, reduction of malignant phenotype, DNA fragmentation, activation of the caspase cascade, increase percentage of cells in sub G0/G1 phase	113
	HCT-15 human colon cancer cells		
PHA	SP2 myeloma, Lox-2 Ab-producing hybridoma	CA/A	114
	B-DLCL human large B-cell lymphoma	DC/A/BindCellMemR	115
PNA	Human melanoma cells	DC/A/BindCellMemR	116
RBA	Human monoblastic leukemia U937	C/TI, apoptosis, chromatin condensation/ nuclear fragmentation/DNA release, externalization of membrane phosphatidylserine, DNA ladder formation, G2/M phase cell cycle arrest	117
Ricin	BEL7404 hepatoma	Apoptosis, upregulation of Bak	118
Ricin A	Human A431 epidermoid	Ribosome binding/inhibition of protein synthesis	46
TGF- α -rec Ricin A	A431 human epidermoid cancer	C/TI	
TGF- α -rec Ricin A	H226Br brain metastatic <i>var.</i> human NSCLC squamous cells	C/TI	
Saracin	Human T-lymphocytes	Apoptosis, induction of IL-2 secretion	119
SNA	Surgically removed human colon cancer cells	DC/A/BindCellMemR	120
VAA	Murine melanoma and HeLa human cervical cancer	C/TI	121
	A549 human lung carcinoma	Non-apoptotic G1-phase accumulation mechanisms	122
VCA	SK-Hep-1 (p53+), Hep 3B (p53-) hepatic cancer	Apoptosis, down-regulation of Bcl-2/upregulation of Bax, down-regulation of telomerase activity	123
	Human breast cancer	Increased TNF- α , IL-6, IFN- γ and/or IL-4 secretion, Th1- shift in the Th1/Th2 balance	124
WGA	Human pancreatic carcinoma	DC/A/BindCellMemR, internalization of lectin, apoptosis, chromatin condensation/ nuclear fragmentation/DNA release	125
	AR42J rat pancreatic cell line	C/TI	41

Note: *italic* indicate weak effects.

Abbreviations: Cytotoxicity/tumor inhibition – C/TI; cell agglutination/aggregation – CA/A; Direct contact/adhesion/binding to cell membrane or receptors – DC/A/BindCellMemR

in rat pancreatic tumor cell line AR42J⁴¹. The alterations of the carbohydrate structures of cellular glycoconjugates may be related to goblet cell differentiation in normal, benign and malignant human colorectal tissues⁴². WGA proved highly toxic to human pancreatic carcinoma cells *in vitro*, primarily to sialic acid residues, with lectin internalization. Cytotoxic effect was found in leukemia cells, several human breast cell lines, skin, and liver cancer cell lines *in vitro* as shows Table 4.

Ribosome-inactivating proteins (RIPs) irreversibly inhibit protein synthesis through the removal of

one or more adenine residue from ribosomal RNA (rRNA). RIPs type I (approximately 30 kDa) consist of an enzymatically active A-chain, while RIPs type II (approximately 56-65 kDa) consists of chain A and chain B which is specific for galactose-like sugars⁴³. Type I is less cytotoxic because this structure does not contain chain B. They are important in several clinical medicine and biomedical research, especially in immunological research and as individual or combined human immunodeficiency virus (HIV) drug therapy as well as anticancer therapy⁴³. The RIPs type II such as ebulin I, foetidissimin

II, mistletoe, nigrin b, riproximin have shown anti-cancer activity *in vitro* and *in vivo*. Ricin, abrin-a, *Sambucus nigra* agglutinin (SNA) and related plant lectins belong to RIPS^{44,45}.

Recently, these toxins have been investigated in experimental models which elucidate the intracellular trafficking of endocytosed proteins⁴⁶. Transforming growth factor (TGF- α) specifically binds and stimulates phosphorylation of the EGF receptor (EGFR) and activates protein kinase activity during cell signaling. TGF- α is highly expressed in human cancer cells. Synthesis of ricin A conjugate and TGF- α trigger cell proliferation⁴⁶, and available levels of EGFR influence the cytotoxic effect on human cancer cells, indicating the involvement of receptor-mediated endocytosis of the conjugate⁴⁶.

Lectin-binding specificity as a rule of recognition carbohydrates allows phenotypic and functional characterization of membrane-associated glycoproteins expressed on cancer cells. In comparison to normal lymphocytes, toxin abrin-a showed greater cytoagglutination against human cultured cell lines derived from acute lymphoblastic leukemia and adult T-cell leukemia⁴⁷.

Lectins (ML-I, ML-II, and ML-III) are the main constituents of mistletoe (MLs) which are responsible for its anticancer and immunomodulatory effects. Nowadays, most researchers focus on investigation of mistletoe lectins, particularly mistletoe lectin I (ML-I). Cytotoxic A-chain inhibits the elongation step of protein biosynthesis by catalyzing the hydrolysis of the *N*-glycosidic bond on ribosomes, resulting in apoptosis or necrosis cell death⁴⁸. Chain B is immunomodulatory, enhancing the secretion of cytokines and activates NK cell⁴⁹ which are involved in anticancer activity. This has been observed in 20 mammary carcinoma patients who received ML-I by subcutaneous injections⁵⁰. Recently, the high resemblance between the 3D structure of mistletoe lectin and the shiga toxin from *Shigella dysenteriae* was found, which represents the bacterial origin of this protein⁵¹. Furthermore, it was suggested that a combination of mistletoe lectin with other forms of recognition receptor ligand substances enhances the immune stimulatory effect.

Several *in vitro* experiments have investigated the anticancer effect of mistletoe extract and its

active compounds in breast cancer and predominantly reported their anticancer and cytotoxic effects in cancer cell lines, as presented in Table 5. Mistletoe extracts exhibit substantial cytotoxic effects *in vitro* and none of the studies reported growth stimulation and proliferation of tumor cell lines⁵²⁻⁵⁷.

Viscum album is the European type of mistletoe. Numerous preparations of this plant from different host trees like apple, pine, oak, and others have been used. Also, mistletoe extract preparations are commercially available, including *Israel*, *Cefaleksin*, *Lektinol*, *Eurixor*, *Iscador*, *Helixor*, *Isucucin*, and *Abnobaviscum*⁵⁸. The antineoplastic activity of *Viscum album* agglutinin-1 (VAA-1) alone or in combination with other chemotherapeutic drugs, including doxorubicin, cisplatin, and taxol, was evaluated in the human lung carcinoma cell line A549. Stronger synergistic effects were noticed using VAA-1 for all drugs tested. Moreover, VAA-1 was able to induce nonapoptotic G1-phase accumulation mechanisms⁵⁹. Recombinant mistletoe lectin alone or in combination with ionizing radiation also showed down regulation of the proliferative activity and cell killing of transforming murine tumor cells⁶⁰.

In vivo studies

The cytotoxic and anticancer activity of plant lectins tested on different animal models has been demonstrated in most of the investigations. They are administrated as oral, intramuscular, intrapleural, intraperitoneal, and intratumoral on relevant sites.

Mistletoe lectins are the most studied lectins in preclinical studies and clinical trials. Until to date, the PubMed database alone lists more than 1280 citations for "mistletoe," of which 113 are clinical studies. Preclinical and clinical studies demonstrated immune response, cytotoxicity, proapoptotic effects, antiangiogenesis, and DNA stabilization⁶¹⁻⁶⁵.

Drees et al.⁶⁶ reported the reduction of cell proliferation in MAXF 449 cell line, sc/Nude mice using *Abnobaviscum M*. Beuth et al.⁶⁷ demonstrated the dose-dependent anticancer activity of *Helixor* using a BALB/c-mouse/BT474 ductal breast carcinoma model. In their *in vivo* experiment, standardized mistletoe extracts harvested from de-

Table 5. Inhibitory effects of plant lectins on malignant cells *in vivo* modified according to Ref. 34.

Lectins	Animal model	Type of effects	References
Anti-CD64-Ricin-A	Acute human myeloid leukemia in NOD/SCID mice	C/TI	107
Con A	B16 melanoma cells in mice	C/TI	71
rML	Human ovarian cancer in SCID mice	C/TI	126
	Chemically induced urinary bladder cancer in mice	C/TI, lower expression of IL-10	127
	Nitrosurea-induced urinary bladder cancer in rat	C/TI, DC/A/BindCellMemR, lower expression of IL-10	128
	Murine melanoma in mice	C/TI, inhibition of metastasis	121
PHA	Non-Hodgkin's lymphoma in mice	C/TI, competition for polyamines	129
	NMRI mice injected with Krebs II lymphosarcoma	C/TI	130
	MCF7 and T47D metastatic human breast cancer lines, SCID mice	DC/A/BindCellMemR	131
	HBL100, BT20, and HS578T human breast cancer lines, SCID mice	<i>DC/A/BindCellMemR</i>	
	HT29 highly metastatic colon cancer, SCID mice	DC/A/BindCellMemR	
	CACO2 colon cancer, SCID mice	DC/A/BindCellMemR	
	HT29 highly metastatic colon cancer, SCID mice	DC/A/BindCellMemR	
	VAA Urinary bladder carcinoma MB49 in mice	C/TI, reduction of malignant phenotype, <i>inhibition of metastasis</i>	132
	VCA C57BL6 mice with B16-BL6 melanoma cells	C/TI, apoptosis, inhibition of angiogenesis, inhibition of metastasis	133
WGA	Colon carcinoma in F-344 rats	C/TI	134
	Human colorectal cancer	Inhibition of metastasis better prognosis/longer survival times	135
MCL	Nasopharyngeal carcinoma (NPC), Nude mice	C/TI, 45 % remission of NPC xenograft tumors	136
SBL	Dalton' lymphoma bearing mice (DL)	Inhibition of tumor proliferation	137
AGG	Nude mice bearing HepG2 xenografts	Reduced tumor growth, increased TUNEL expression, decreased CD-31 and Ki-67 expression	138

Note: *italic* indicate weak effects.

fined host trees ME-A and ME-M (fir tree *Abies*, ME-A, Helixor[®]A, and apple tree *Malus*, ME-M, Helixor[®]M) were applied to the breast carcinoma model intratumorally. *In vivo* investigations of the ability of mistletoe extract to improve tumor survival induce apoptosis and necrosis and inhibit cancer cell proliferation in animal models that have yielded inconsistent results. Seifert et al.⁶⁸ investigated both the cytotoxic effect and the mechanism of action of two standardized aqueous MEs (ME-A and ME-P obtained from the pine

tree) in an *in vivo* the severe combined immunodeficiency model (SCID) mice of B-precursor acute lymphoblastic leukemia (pre-B ALL) cell line (NALM-6). Both MEs significantly improved survival (up to 55.4 days) at all tested concentrations in contrast to controls (34.6 days) without side effects. However, some research showed decreases in the rate of cell proliferation and improvement in tumor survival.

Mushrooms have become popular sources of natural anticancer, antiviral, antibacterial, antiox-

idative, and immunomodulatory agents. *Pleurotus citrinopileatus* lectin (PCL)⁶⁹ and *Russula lepida* lectin (RLL)⁷⁰ exerted potent anticancer activity in mice bearing sarcoma 180, and caused inhibition of tumor growth when administered intraperitoneally.

It has also been shown that the chemical modification of polyethylene glycol-modified concanavalin A (PEG-Con A) enhanced the anticancer cytotoxicity of peripheral lymphocytes against melanoma B16 cells in mice⁷¹. The encapsulation of *Cratylia mollis* lectin (CML) with liposomes lowered its tissue toxicity in the liver and kidney, and improved its anticancer activity in Swiss mice inoculated with sarcoma 180 cell line⁷². Reduction in the tumor size and inhibition of growth represents basic outcomes of used lectin therapy. The second type of lectins application is combined with conventional anticancer therapy. Toxic lectins can be often used as a supportive therapy to improve health-related quality of life (HRQoL). It has been shown that the use of some types of complementary and alternative medicine (CAM) in breast cancer patients has rapidly increased⁷³. In recent years there has been an increase of research studies on mistletoe therapy, including studies of its co-administration alongside chemotherapy to reduce adverse effects and to improve quality of life in breast cancer, ovarian cancer, and lung cancer patients. These clinical trials have not found significant clinical efficacy in terms of tumor control and survival time for patients. However, studies have shown a positive outcome given the HRQoL⁶¹⁻⁶⁵.

To design effective cancer vaccines, the best tumor antigens should be combined with the most effective immunogen to achieve better clinical results. Plant lectins can be also applied as immunoadjuvants to enhance antigen-specific tumor activity. Ricin toxin (RTB) was used as immunoadjuvants fused with HPV-16 E7 to prepare an effective vaccine, which could inhibit tumor growth in the lung. The immunization with E7-RTB protein without adjuvant can generate anticancer effects in mice challenged with TC-1 cells. This research confirms the clinical application of therapeutic vaccines with lectins as immunoadjuvants is directed to design effective cancer vaccines⁷⁴.

PERSPECTIVE

Plant lectins have been investigated for a long time. However, new research challenges are present. Some of them will be indicated below.

Applications of glycosylated nanomaterials in nanotechnology have gained significant attention in recent years due to their unique structural properties and compatibility in biological systems⁷⁵⁻⁷⁷. Strategies for building various types of glyco-nanoparticles (glyco-NPs) and functionalized carbon nanotubes (CNTs)⁷⁸ and highlights their potential in targeted drug delivery and molecular imaging as well as their uses in bioassays and biosensors. Glyco-NPs contain a nano-sized metallic core that exhibits carbohydrate ligands on the surface in three dimensions polyvalent displays similar to the glycocalyx structures of cell membranes. The most recent examples of glyco-NPs are as vaccine candidates and probes for assaying enzymes with bond-forming activities. CNTs have attracted great attention in biomedical applications due to their molecular size and unique properties. Introduction of biofunctionalities by integration of carbohydrate with CNTs provide new tools for glycobiological studies⁷⁸.

Carbohydrates are crucial for a wide variety of cellular processes ranging from cell-cell communication to immunity, and they are altered in disease states such as cancer and inflammation.

Development of glycan analysis towards high-throughput analytics are new challenges in the fields of glycomics and glycoproteomics. These include advances in applying separation, mass spectrometry, and microarray methods to the fields of glycomics and glycoproteomics. These new bioanalytical techniques influenced the progress in understanding the importance of glycosylation in biology and disease⁷⁹⁻⁸¹.

ACKNOWLEDGEMENTS

This work has been supported by the University of Rijeka under the project number 13.11.1.2.02. The author thanks to Professor Arijana Krišković for the revision of the English and Professor Aleina Buretić-Tomljanović for useful suggestions.

Conflicts of interest statement: The authors report no conflicts of interest.

REFERENCES

1. Van Damme JM, Lannoo N, Peumans WJ. Plant lectins. *Adv Bot Res* 2008;48:107-209.
2. Garcia-Pino A, Buts L, Wyns L, Imberty A, Loris R. How a plant lectin recognizes high mannose oligosaccharides?. *Plant Physiol* 2007;144:1733-41.
3. Berg JM, Tymoczko JL, Stryer L. *Biochemistry* [Internet]. 5th edition. New York: W H Freeman; 2002. Section 11.4, Lectins Are Specific Carbohydrate-Binding Proteins. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK22545/>
4. Khan F, Khan RH, Sherwani A, Mohmood S, Azfer MA. Lectins as markers for blood grouping. *Med Sci Monit* 2002;8:RA293-300.
5. Sharon N, Lis H. History of lectins: from hemagglutinins to biological recognition molecules. *Glycobiology* 2004;14:53R-62R.
6. Michiels K, Van Damme EJM, Smagghe G. Plant-insect interactions: what can we learn from plant lectins?. *Arch Insect Biochem Physiol* 2010;73:193-212.
7. Regnier FE, Jung K, Hooser SB, Wilson CR. Chapter 8 – Glycoproteomics Based on Lectin Affinity Chromatographic Selection of Glycoforms. In: Nilsson CL (ed). *Lectins*. Amsterdam: Elsevier Science B.V., 2007;193-212.
8. Franco Fraguas L, Carlsson J, Lönnberg M. Lectin affinity chromatography as a tool to differentiate endogenous and recombinant erythropoietins. *J Chromatogr A* 2008;1212:82-8.
9. Alvarez-Manilla G, Warren NL, Atwood J 3rd, Orlando R, Dalton S, Pierce M. Glycoproteomic analysis of embryonic stem cells: identification of potential glycoproteomic biomarkers using lectin affinity chromatography of glycopeptides. *J Proteome Res* 2010;9:2062-75.
10. Qi Y-J, Ward DG, Pang C, Wang QM, Wei W, Ma J et al. Proteomic profiling of N-linked glycoproteins identifies ConA-binding procathepsin D as a novel serum biomarker for hepatocellular carcinoma. *Proteomics* 2014;14:186-95.
11. Ghazarian A, Oppenheimer SB. Microbead analysis of cell binding to immobilized lectin. Part II: Quantitative kinetic profile assay for possible identification of anti-infectivity and anti-cancer reagents. *Acta Histochem* 2014;116:1514-8.
12. Oliveira C, Teixeira JA, Domingues L. Recombinant lectins: an array of tailor-made glycan-interaction biosynthetic tools. *Crit Rev Biotechnol* 2013;33:66-80.
13. Lam SK, Ng TB. Lectins: production and practical applications. *Appl Microbiol Biotechnol* 2011;89:45-55.
14. Wang TH, Lee MH, Su NW. Screening of lectins by an enzyme-linked adsorbent assay. *Food Chem* 2009;113:1218-25.
15. Kuno A, Uchiyama N, Koseki-Kuno S, Ebe Y, Takashima S, Yamada M et al. Evanescent-field fluorescence-assisted lectin microarray: a new strategy for glycan profiling. *Nat Methods* 2005;2:851-6.
16. Harvey DJ. Analysis of carbohydrates and glycoconjugates by matrix-assisted laser desorption/ionization mass spectrometry: an update for the period 2005–2006. *Mass Spectrom Rev* 2010;30:1-100.
17. Nasi A, Picariello G, Ferranti P. Proteomic approaches to study structure, functions and toxicity of legume seeds lectins. Perspectives for the assessment of food quality and safety. *J Proteom* 2009;72:527-38.
18. Ahn YH, Ji ES, Oh NR, Kim YS, Ko JH, Yoo JS. Differential proteomic approach for identification and verification of aberrantly glycosylated proteins in adenocarcinoma lung cancer (ADLC) plasmas by lectin-capturing and targeted mass spectrometry. *J Proteomics* 2014;106:221-9.
19. Ahn YH, Shin PM, Kim YS, Oh NR, Ji ES, Kim KH et al. Quantitative analysis of aberrant protein glycosylation in liver cancer plasma by AAL-enrichment and MRM mass spectrometry. *Analyst* 2013;138:6454-62.
20. Souza MA, Carvalho FC, Ruas LP, Ricci-Azevedo R, Roque-Barreira MC. The immunomodulatory effect of plant lectins: a review with emphasis on ArtinM properties. *Glycoconj J* 2013;30:641-57.
21. Drickamer K. Evolution of Ca(2+)-dependent animal lectins. *Prog Nucleic Acid Res Mol Biol* 1993;45:207-32.
22. Kilpatrick DC. Animal lectins: a historical introduction and overview. *Biochim Biophys Acta* 2002;1572:187-97.
23. Parham P. *The immune system*. 2nd edition. New York: Garland Science, 2005;244-5.
24. Ley K. The role of selectins in inflammation and disease. *Trends Mol Med* 2003;9:263-8.
25. Rossi B, Constantin G. Anti-selectin therapy for the treatment of inflammatory diseases. *Inflamm Allergy Drug Targets* 2008;7:85-93.
26. Huo Y, Xia L. P-selectin glycoprotein ligand-1 plays a crucial role in the selective recruitment of leukocytes into the atherosclerotic arterial wall. *Trends Cardiovasc Med* 2009;19:140-5.
27. Wang K, Zhou X, Zhou Z, Tarakji K, Qin JX, Sitges M et al. Recombinant soluble P-selectin glycoprotein ligand-Ig (rPSGL-Ig) attenuates infarct size and myeloperoxidase activity in a canine model of ischemia-reperfusion. *Thromb Haemost* 2002;88:149-54.
28. Assinger A. Platelets and infection – an emerging role of platelets in viral infection. *Front Immunol* [Internet] 2014;5:689. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4270245/>
29. Ogata M, Uzawa M, Kazuya I, Hidari PJ, Suzuki T, Park EY et al. Facile Synthesis of Sulfated Sialoglycopolypeptides with a γ -Polyglutamic Acid Backbone as Hemagglutinin Inhibitors against Influenza Virus. *J Appl Glycosci* 2014;61:1-7.
30. Hartshorn KL, White MR. Influenza A virus up-regulates neutrophil adhesion molecules and adhesion to biological surfaces. *J Leukoc Biol* 1999;65:614-22.
31. Ferroni P, Mammarella A, Martini F, Paoletti V, Cardarello CM, Labbadia G et al. Increased soluble P-selectin levels in hepatitis C virus-related chronic hepatitis: correlation with viral load. *J Investig Med* 2001;49:407-12.
32. Vardareli E, Saricam T, Demirustu C, Gulbas Z. Soluble P selectin levels in chronic liver disease: relationship to disease severity. *Hepatogastroenterology* 2007;54:466-9.
33. Jiang Q-L, Zhang S, Tian M, Zhang S-Y, Xie T, Chen D-Y et al. Plant lectins, from ancient sugar-binding proteins to emerging anti-cancer drugs in apoptosis and autophagy. *Cell Prolif* 2015;48:17-28.
34. De Mejia EG, Prisecaru VI. Lectins as bioactive plant proteins: a potential in cancer treatment. *Crit Rev Food Sci Nutr* 2005;45:425-45.
35. De Mejia EG, Dia VP. The role of nutraceutical proteins and peptides in apoptosis, angiogenesis, and metastasis of cancer cells. *Cancer Metastasis Rev* 2010;29:511-28.

36. Kobayashi Y, Nakamura H, Sekiguchi T, Takanami R, Murata T, Usui T et al. Analysis of the carbohydrate binding specificity of the mushroom *Pleurotus ostreatus* lectin by surface plasmon resonance. *Anal Biochem* 2005; 336:87-93.
37. Wang H, Ng TB, Ooi VE, Liu WK. Effects of lectins with different carbohydrate-binding specificities on hepatoma, choriocarcinoma, melanoma and osteosarcoma cell lines. *Int J Biochem Cell Biol* 2000;32:365-72.
38. Gu Y-H, Sivam G. Cytotoxic Effect of Oyster Mushroom *Pleurotus ostreatus* on Human Androgen-Independent Prostate Cancer PC-3 Cells. *J Med Food* 2006;9:196-204.
39. Sarangi I, Ghosh D, Bhutia SK, Mallick SK, Maiti TK. Antitumor and immunomodulating effects of *Pleurotus ostreatus* mycelia-derived proteoglycans. *Int Immunopharmacol* 2006;6:1287-97.
40. Jordinson M, El-Hariry I, Calnan D, Calam J, Pignatelli M. *Vicia faba* agglutinin, the lectin present in broad beans, stimulates differentiation of undifferentiated colon cancer cells. *Gut* 1999;44:709-14.
41. Mikkat U, Damm I, Kirchoff F, Albrecht E, Nebe B, Jonas L. Effects of lectins on CCK-8-stimulated enzyme secretion and differentiation of the rat pancreatic cell line AR42J. *Pancreas* 2002;23:368-74.
42. Remani P, Nair RA, Sreelekha TT, Madhavan J, Vijayakumar T, Nair MK. Altered expression of jack fruit lectin specific glycoconjugates in benign and malignant human colorectum. *J Exp Clin Cancer Res* 2000;19:519-23.
43. Puri M, Kaur I, Perugini MA, Gupta RC. Ribosome-inactivating proteins: current status and biomedical applications. *Drug Discov Today* 2012;17:774-83.
44. Olsnes S, Kozlov JV. Ricin. *Toxicon* 2001;39:1723-8.
45. Olsnes S. The history of ricin, abrin and related toxins. *Toxicon* 2004;44:361-70.
46. Fang K. A toxin conjugate containing transforming growth factor- α and ricin A specifically inhibits growth of A431 human epidermoid cancer cells. *Proc Natl Sci Coun Repub China B* 1998;22:76-82.
47. Moriwaki S, Ohba H, Nakamura O, Sallay I, Suzuki M, Tsubouchi H et al. Biological activities of the lectin, abrin-a, against human lymphocytes, and cultured leukemic cell lines. *J Hematother Stem Cell Res* 2000;9:47-53.
48. Bantel H, Engels IH, Voelter W, Schulze-Osthoff K, Wesselborg S. Mistletoe lectin activates caspase-8/FLICE independently of death receptor signaling and enhances anticancer drug-induced apoptosis. *Cancer Res* 1999;59:2083-90.
49. Lee C, Kim J, Kim H, Park S, Lee S. Immunomodulating effects of Korean mistletoe lectin in vitro and in vivo. *Int Immunopharmacol* 2009;9:1555-61.
50. Beuth J, Stoffel B, Ko HL, Buss G, Tunggal L, Pulverer G. Immunostimulating activity of different dosages of mistletoe lectin-1 in patients with mammary carcinoma. *Arzneimittel-Forschung* 1995;45:505-7.
51. Maletzki C, Linnebacher M, Savai R, Hobohm U. Mistletoe lectin has a shiga toxin-like structure and should be combined with other Toll-like receptor ligands in cancer therapy. *Cancer Immunol Immunotherapy* 2013;62:1283-92.
52. Frantz M, Jung ML, Ribereau-Gayon G, Anton R. Modulation of mistletoe (*Viscum album* L.) lectins cytotoxicity by carbohydrates and serum glycoproteins. *Arzneimittelforschung* 2000;50:471-8.
53. Vervecken W, Kleff S, Pfuller U, Bussing A. Induction of apoptosis by mistletoe lectin I and its subunits. No evidence for cytotoxic effects caused by isolated A- and B-chains. *Int J Biochem Cell Biol* 2000;32:317-26.
54. Pae HO, Oh GS, Kim NY, Shin MK, Lee HS, Yun YG et al. Roles of extracellular signal-regulated kinase and p38 mitogen-activated protein kinase in apoptosis of human monoblastic leukemia U937 cells by lectin-II isolated from Korean mistletoe. *In Vitro Mol Toxicol* 2001;14:99-106.
55. Kim MS, So HS, Lee KM, Park JS, Lee JH, Moon SK et al. Activation of caspase cascades in Korean mistletoe (*Viscum album* var. *coloratum*) lectin-II-induced apoptosis of human yeloleukemic U937 cells. *Gen Pharmacol* 2000;34:349-55.
56. Park R, Kim MS, So HS, Jung BH, Moon SR, Chung SY et al. Activation of c-Jun N-terminal kinase 1 (JNK1) in mistletoe lectin II-induced apoptosis of human myeloleukemic U937 cells. *Biochem Pharmacol* 2000;60:1685-91.
57. Steele ML, Axtner J, Happe A, Kröz M, Matthes H, Schad F. Adverse Drug Reactions and Expected Effects to Therapy with Subcutaneous Mistletoe Extracts (*Viscum album* L.) in Cancer Patients. Evidence-Based Complement Altern Med Volume [Internet]. 2014;724258. Available from: <http://dx.doi.org/10.1155/2014/724258>.
58. Thies A, Pfuller U, Schachner M, Horny HP, Molls I, Schumacher U. Binding of mistletoe lectins to cutaneous malignant melanoma: Implications for prognosis and therapy. *Anticancer Res* 2001;21:2883-7.
59. Siegle I, Fritz P, McClellan M, Gutzeit S, Murdter TE. Combined cytotoxic action of *Viscum album* agglutinin-1 and anticancer agents against human A549 lung cancer cells. *Anticancer Res* 2001;21:2687-91.
60. Hostanska K, Vuong V, Rocha S, Soengas MS, Glanzmann C, Saller R et al. Recombinant mistletoe lectin induces p53-independent apoptosis in tumour cells and cooperates with ionising radiation. *Br J Cancer* 2003;88:1785-92.
61. Longhi A, Reif M, Mariani E, Ferrari S. A Randomized Study on Postrelapse Disease-Free Survival with Adjuvant Mistletoe versus Oral Etoposide in Osteosarcoma Patients. *Eur J Integrat Med* 2009;1:27-33.
62. Tröger W, Ždrale Z, Tišma N, Matijašević M. Additional Therapy with a Mistletoe Product during Adjuvant Chemotherapy of Breast Cancer Patients, Improves Quality of Life: An Open Randomized Clinical Pilot Trial. Evidence-Based Complement Altern Med Volume [Internet]. 2014;430518. Available from: <http://dx.doi.org/10.1155/2014/430518>.
63. Mansky PJ, Wallerstedt DB, Sannes TS, Stagl J, Johnson LL, Blackman MR et al. Erratum to "NCCAM/NCI Phase 1 Study of Mistletoe Extract and Gemcitabine in Patients with Advanced Solid Tumors". *Evid Based Complement Alternat Med* [Internet]. 2014;606348. Available from: www.ncbi.nlm.nih.gov/pmc/articles/PMC3944907/.
64. Steele ML, Axtner J, Happe A, Kröz M, Matthes H, Schad F. Adverse Drug Reactions and Expected Effects to Therapy with Subcutaneous Mistletoe Extracts (*Viscum album* L.) in Cancer Patients. *Evid Based Complement Alternat Med*. 2014;724258. Available from: <http://dx.doi.org/10.1155/2014/724258>.

65. Mansky PJ, Wallerstedt DB, Sannes TS, Stagl J, Johnson LL, Blackman MR et al. NCCAM/NCI Phase 1 Study of Mistletoe Extract and Gemcitabine in Patients with Advanced Solid Tumors, Evid Based Complement Alternat Med [Internet]. 2013;964592. Available from: <http://dx.doi.org/10.1155/2013/964592>.
66. Drees M, Berger DP, Dengler WA, Fiebig GH. Direct cytotoxic effects of preparations used as unconventional methods in cancer therapy in human tumor xenografts in the clonogenic assay and in nude mice. *In: Arnold W, Köpf-Maier P, Micheel B (eds). Immunodeficient Animals: Models For Cancer Research. Volume 51. Basel: Karger Verlag, 1996;115-22.*
67. Beuth J, Ko HL, Schneider H, Tawadros S, Kasper HU, Zimst H et al. Intratumoral application of standardized mistletoe extracts down regulates tumor weight *via* decreased cell proliferation, increased apoptosis and necrosis in a murine model. *Anticancer Res* 2006;26:4451-6.
68. Seifert G, Jesse P, Laengler A, Reindl T, Lüth M, Lobitz S et al. Molecular mechanisms of mistletoe plant extract-induced apoptosis in acute lymphoblastic leukemia *in vivo* and *in vitro*. *Cancer Lett* 2008;264:218-28.
69. Li YR, Liu QH, Wang HX, Ng TB. A novel lectin with potent antitumor, mitogenic and HIV-1 reverse transcriptase inhibitory activities from the edible mushroom *Pleurotus citrinopileatus*. *Biochim Biophys Acta* 2008;1780:51-7.
70. Zhang G, Sun J, Wang H, Ng TB. First isolation and characterization of a novel lectin with potent antitumor activity from a *Russula* mushroom. *Phytomedicine* 2010;17:775-81.
71. Ueno T, Ohtawa K, Kimoto Y, Sakurai K, Koderia Y, Hiroto M. Polyethylene glycol-modified concanavalin A as an effective agent to stimulate anti-tumor cytotoxicity. *Cancer Detect Prev* 2000;24:100-6.
72. Andrade CA, Correia MT, Coelho LC, Nascimento SC, Santos-Magalhães NS. Antitumor activity of *Cratylia mollis* lectin encapsulated into liposomes. *Int J Pharm* 2004;278:435-45.
73. Gerber B, Scholz C, Reimer T, Briese V, Janni W. Complementary and alternative therapeutic approaches in patients with early breast cancer: a systematic review. *Breast Cancer Res Treat* 2006;95:199-209.
74. Sadraeian M, Rasoul-Amini S, Mansoorkhani MJ, Mohkam M, Ghoshoon MB, Ghasemi Y. Induction of antitumor immunity against cervical cancer by protein HPV-16 E7 in fusion with ricin B chain in tumor-bearing mice. *Int J Gynecol Cancer* 2013;23:809-14.
75. García I, Marradi M, Penadés S. Glyconanoparticles: multifunctional nanomaterials for biomedical applications. *Nanomedicine (Lond)* 2010;5:777-92.
76. Adak AK, Li BY, Lin CC. Advances in multifunctional glycosylated nanomaterials: preparation and applications in glycoscience. *Carbohydr Res* 2015;405C:2-12.
77. Marín MJ, Schofield CL, Field RA, Russell DA. Glyconanoparticles for colorimetric bioassays. *Analyst* 2014;140:59-70.
78. Gorityala BK, Ma J, Wang X, Chen P, Liu X-W. Carbohydrate functionalized carbon nanotubes and their applications, *Chem Soc Rev* 2010;39:2925-34.
79. Rakus JF, Mahal LK. New Technologies for Glycomic Analysis: Toward a Systematic Understanding of the Glycome. *Ann Rev Anal Chem* 2011;4:367-92.
80. Katrlík J, Svitel J, Gemeiner P, Kozár T, Tkac J. Glycan and lectin microarrays for glycomics and medicinal applications. *Med Res Rev* 2010;30:394-418.
81. Campbell C, Gildersleeve J. Tools for Glycomics: Glycan and Lectin Microarrays. *In: Wang B, Boons G-J (eds). Carbohydrate Recognition: Biological Problems, Methods, and Applications. Hoboken: John Wiley & Sons, Inc., 2011;205-27.*
82. Sharma A, Ng TB, Wong JH, Lin P. Purification and characterization of a lectin from *Phaseolus vulgaris* cv. (Anasazi beans). *J Biomed Biotechnol*. [Internet]. 2009;929568. Available from: <http://dx.doi.org/10.1155/2009/929568>.
83. Xia L, Ng TB. A hemagglutinin with mitogenic activity from dark red kidney beans. *J Chromatogr B Analyt Technol Biomed Life Sci* 2006;844:213-6.
84. Castillo-Villanueva A, Caballero-Ortega H, Abdullaev-Jafarova F, Garfias Y, Jiménez-Martínez CM, Bouquelet S et al. Lectin from *Phaseolus acutifolius* var. escumite: chemical characterization, sugar specificity, and effect on human T-lymphocytes. *J Agric Food Chem* 2007;55:5781-7.
85. Fang EF, Lin P, Wong JH, Tsao SW, Ng TB. A lectin with anti-HIV-1 reverse transcriptase, antitumor, and nitric oxide inducing activities from seeds of *Phaseolus vulgaris* cv. extralong autumn purple bean. *J Agric Food Chem* 2010;58:2221-9.
86. Leung EH, Wong JH, Ng TB. Concurrent purification of two defense proteins from French bean seeds: a defensin-like antifungal peptide and a hemagglutinin. *J Pept Sci* 2008;14:349-53.
87. Lam SK, Ng TB. Isolation and characterization of a French bean hemagglutinin with antitumor, antifungal, and anti-HIV-1 reverse transcriptase activities and an exceptionally high yield. *Phytomedicine* 2010;17:457-62.
88. Ye XY, Ng TB, Tsang PW, Wang J. Isolation of a homodimeric lectin with antifungal and antiviral activities from red kidney bean (*Phaseolus vulgaris*) seeds. *J Protein Chem* 2001;20:367-75.
89. Kvennefors EC, Leggat W, Hoegh-Guldberg O, Degnan BM, Barnes AC. An ancient and variable mannose-binding lectin from the coral *Acropora millepora* binds both pathogens and symbionts. *Dev Comp Immunol* 2008;32:1582-92.
90. Pan S, Tang J, Gu X. Isolation and characterization of a novel fucose-binding lectin from the gill of bighead carp (*Aristichthys nobilis*). *Vet Immunol Immunopathol* 2010;133:154-64.
91. Ashraf GM, Rizvi S, Naqvi S, Suhail N, Bilal N, Hasan S et al. Purification, characterization, structural analysis and protein chemistry of a buffalo heart galectin-1. *Amino Acids* 2010;39:1321-32.
92. Ashraf GM, Banu N, Ahmad A, Singh LP, Kumar R. Purification, characterization, sequencing and biological chemistry of galectin-1 purified from *Capra hircus* (goat) heart. *Protein J* 2011;30:39-51.
93. Gowda NM, Goswami U, Khan MI. Purification and characterization of a T-antigen specific lectin from the coelomic fluid of a marine invertebrate, sea cucumber (*Holothuria scabra*). *Fish Shellfish Immunol* 2008;24:450-8.
94. Adhya M, Singha B, Chatterjee BP. Purification and characterization of an N-acetylglucosamine specific lectin from marine bivalve *Macoma birmanica*. *Fish Shellfish Immunol* 2009;27:1-8.

95. Imamichi Y, Yokoyama Y. Purification, characterization and cDNA cloning of a novel lectin from the jellyfish *Nemopilema nomurai*. *Comp Biochem Physiol B Biochem Mol Biol* 2010;156:12-8.
96. Upadhyay SK, Saurabh S, Rai P, Singh R, Chandrashekar K, Verma PC et al. SUMO fusion facilitates expression and purification of garlic leaf lectin but modifies some of its properties. *J Biotechnol* 2010;146:1-8.
97. Oliveira C, Costa S, Teixeira JA, Domingues L. cDNA cloning and functional expression of the alpha-d-galactose-binding lectin frutalin in *Escherichia coli*. *Mol Biotechnol* 2009;43:212-20.
98. Oliveira C, Felix W, Moreira RA, Teixeira JA, Domingues L. Expression of frutalin, an alpha-d-galactose-binding jacalin-related lectin, in the yeast *Pichia pastoris*. *Protein Expr Purif* 2008;60:188-93.
99. Adar R, Streicher H, Rozenblatt S, Sharon N. Synthesis of soybean agglutinin in bacterial and mammalian cells. *Eur J Biochem* 1997;249:684-9.
100. Lannoo N, Verveckens W, Proost P, Rougé P, Van Damme EJ. Expression of the nucleocytoplasmic tobacco lectin in the yeast *Pichia pastoris*. *Protein Expr Purif* 2007;53:275-82.
101. Branco AT, Bernabé RB, Santos FB, Oliveira MV, Garcia AB, Souza Filho GA. Expression and purification of the recombinant SALT lectin from rice (*Oryza sativa* L.). *Protein Expr Purif* 2004;33:34-8.
102. Stubbs ME, Carver JP, Dunn RJ. Production of pea lectin in *Escherichia coli*. *J Biol Chem* 1986;261:6141-4.
103. Tateno H, Winter HC, Goldstein IJ. Cloning, expression in *Escherichia coli* and characterization of the recombinant Neu5Acalpha2, 6Galbeta1, 4GlcNAc-specific high-affinity lectin and its mutants from the mushroom *Polyporus squamosus*. *Biochem J* 2004;382:667-75.
104. Huang LH, Yan QJ, Kopparapu NK, Jiang ZQ, Sun Y. *Astragalus membranaceus* lectin (AML) induces caspase-dependent apoptosis in human leukemia cells. *Cell Prolif* 2012;45:15-21.
105. Yan Q, Li Y, Jiang Z, Sun Y, Zhu L, Ding Z. Antiproliferation and apoptosis of human tumor cell lines by a lectin (AMML) of *Astragalus mongholicus*. *Phytomedicine* 2009;16:586-93.
106. Yu LG, Milton JD, Fernig DG, Rhodes JM. Opposite effects on human colon cancer cell proliferation of two dietary Thomsen-Friedenreich antigen-binding lectins. *J Cell Physiol* 2001;186:282-7.
107. Zhong RK, van de Winkel JG, Thepen T, Schultz LD, Ball ED. Cytotoxicity of anti-CD64-ricin A chain immunotoxin against human acute myeloid leukemia cells *in vitro* and in SCID mice. *J Hematother Stem Cell Res* 2001;10:95-105.
108. Van Horssen PJ, van Oosterhout YV, Evers S, Backus HH, van Oijen MG, Bongaerts R et al. Influence of cytotoxicity enhancers in combination with human serum on the activity of CD22-recombinant ricin A against B cell lines, chronic and acute lymphocytic leukemia cells. *Leukemia* 1999;13:241-9.
109. Sames K, Schumacher U, Halata Z, Van Damme EJ, Peumans WJ, Asmus B et al. Lectin and proteoglycan histochemistry of Merkel cell carcinomas. *Exp Dermatol* 2001;10:100-9.
110. Rimmelink M, Darro F, Decaestecker C, De Decker R, Bovin NV, Gebhart M et al. *In vitro* influence of lectins and neoglycoconjugates on the growth of three human sarcoma cell lines. *J Cancer Res Clin Oncol* 1999;125:275-85.
111. Lorea P, Goldschmidt D, Darro F, Salmon I, Bovin N, Gabius HJ et al. *In vitro* characterization of lectin-induced alterations on the proliferative activity of three human melanoma cell lines. *Melanoma Res* 1997;7:353-63.
112. Desilets DJ, Davis KE, Nair PP, Salata KF, Maydonovitch CL, Howard RS et al. Lectin binding to human colonocytes is predictive of colonic neoplasia. *Am J Gastroenterol* 1999;94:744-50.
113. Deepa M, Sureshkumar T, Satheeshkumar PK, Priya S. Purified mulberry leaf lectin (MLL) induces apoptosis and cell cycle arrest in human breast cancer and colon cancer cells. *Chem Biol Interact*; 2012;200:38-44.
114. Takamatsu H, Kawajiri H, Takahashi Y, Ali AM, Yoshimoto T. Continuous antibody production by phytohemagglutinin-L-aggregated hybridoma cells. *J Immunol Methods* 1999;223:165-70.
115. Suzuki O, Nozawa Y, Kawaguchi T, Abe M. *Phaseolus vulgaris* leucoagglutinating lectin-binding reactivity in human diffuse large B cell lymphoma and its relevance to the patient's clinical outcome: Lectin histochemistry and lectin blot analysis. *Pathol Int* 1999;49:874-80.
116. Cochran AJ, Wen DR, Berthier-Vergnes O, Bailly C, Dore JF, Berard F et al. Cytoplasmic accumulation of peanut agglutinin-binding glycoconjugates in the cells of primary melanoma correlates with clinical outcome. *Hum Pathol* 1999;30:556-61.
117. Miyoshi N, Koyama Y, Katsuno Y, Hayakawa S, Mita T, Ohta T et al. Apoptosis induction associated with cell cycle dysregulation by rice bran agglutinin. *J Biochem Tokyo* 2001;130:799-805.
118. Hu R, Zhai Q, Liu W, Liu X. An insight into the mechanism of cytotoxicity of ricin to hepatoma cell: Roles of Bcl-2 family proteins, caspases, Ca(2+)-dependent proteases and protein kinase C. *J Cell Biochem*, 2001;81:583-93.
119. Ghosh S, Majumder M, Majumder S, Ganguly NK, Chatterjee BP. Saracin: Alectin from *Saraca indica* seed integument induces apoptosis in human T-lymphocytes. *Arch Biochem Biophys* 1999;371:163-8.
120. Dall'Olio F, Chiricolo M, Mariani E, Facchini A. Biosynthesis of the cancer-related sialyl-alpha 2,6-lactosaminyl epitope in colon cancer cell lines expressing beta-galactoside alpha 2,6-sialyltransferase under a constitutive promoter. *Eur J Biochem* 2001;268:5876-84.
121. Žarković N, Vuković T, Lončarić I, Miletić M, Žarković K, Borović S et al. An overview on anticancer activities of the *Viscum album* extract Isorel. *Cancer Biother Radiopharm* 2001;16:55-62.
122. Siegle I, Fritz P, McClellan M, Gutzeit S, Murdter TE. Combined cytotoxic action of *Viscum album* agglutinin-1 and anticancer agents against human A549 lung cancer cells. *Anticancer Res* 2001;21:2687-91.
123. Lyu SY, Choi SH, Park WB. Korean mistletoe lectin-induced apoptosis in hepatocarcinoma cells is associated with inhibition of telomerase via mitochondrial controlled pathway independent of p53. *Arch Pharm Res* 2002;25:93-101.
124. Stein G, Henn W, von Laue H, Berg P. Modulation of the cellular and humoral immune responses of tumor pa-

- tients by mistletoe therapy. *Eur J Med Res* 1998;3:194-202.
125. Schwarz RE, Wojciechowicz DC, Picon AI, Schwarz MA, Paty PB. Wheatgerm agglutinin-mediated toxicity in pancreatic cancer cells. *Br J Cancer* 1999;80:1754-62.
 126. Schumacher U, Feldhaus S, Mengs U. Recombinant mistletoe lectin (rML) is successful in treating human ovarian cancer cells transplanted into severe combined immunodeficient (SCID) mice. *Cancer Lett* 2000;150:171-5.
 127. Elsässer-Beile U, Ruhnau T, Freudenberg N, Wetterauer U, Mengs U. Antitumoral effect of recombinant mistletoe lectin on chemically induced urinary bladder carcinogenesis in a rat model. *Cancer* 2001;91:998-1004.
 128. Elsässer-Beile U, Voss M, Schuhle R, Wetterauer U. Biological effects of natural and recombinant mistletoe lectin and an aqueous mistletoe extract on human monocytes and lymphocytes *in vitro*. *J Clin Lab Anal* 2001;14:255-9.
 129. Pryme IF, Bardocz S, Pusztai A, Ewen SW. Dietary mistletoe lectin supplementation and reduced growth of a murine non-Hodgkin lymphoma. *Histol Histopathol* 2002;17:261-71.
 130. Pryme IF, Pusztai AJ, Grant G, Bardocz S. The effect of switching between a phytohemagglutinin-containing and a control diet on the growth and lipid content of a Krebs II lymphosarcoma tumor. *Exp Ther Oncol* 1996;1:273-7.
 131. Mitchell BS, Brooks SA, Leatham AJ, Schumacher U. Do HPA and PHA-L have the same binding pattern in metastasizing human breast and colon cancers? *Cancer Lett* 1998;123:113-9.
 132. Mengs U, Schwarz T, Bulitta M, Weber K, Madaus AG. Antitumoral effects of an intravesically applied aqueous mistletoe extract on urinary bladder carcinoma MB49 in mice. *Anticancer Res* 2000;20:3565-8.
 133. Park WB, Lyu SY, Kim JH, Choi SH, Chung HK, Ahn SH et al. Inhibition of tumor growth and metastasis by Korean mistletoe lectin is associated with apoptosis and antiangiogenesis. *Cancer Biother Radiopharm* 2001;16:439-47.
 134. Zalatnai A, Lapis K, Szende B, Raso E, Telekes A, Resetar A et al. Wheat germ extract inhibits experimental colon carcinogenesis in F-344 rats. *Carcinogenesis* 2001;22:1649-52.
 135. Jakab F, Mayer A, Hoffmann A, Hidvegi M. First clinical data of a natural immunomodulator in colorectal cancer. *Hepatogastroenterology* 2000;47:393-5.
 136. Fang EF, Zhang CZ, Ng TB, Wong JH, Pan WL, Ye XJ et al. Momordica Charantia lectin, a type II ribosome inactivating protein, exhibits antitumor activity toward human nasopharyngeal carcinoma cells *in vitro* and *in vivo*. *Cancer Prev Res (Phila)* 2012;5:109-21.
 137. Panda PK, Mukhopadhyay S, Behera B, Bhol CS, Dey S, Das DN et al. Antitumor effect of soybean lectin mediated through reactive oxygen species-dependent pathway. *Life Sci* 2014;111:27-35.
 138. Mukhopadhyay S, Panda PS, Das DN, Sinha N, Behera B, Maiti TK. Abrus agglutinin suppresses human hepatocellular carcinoma *in vitro* and *in vivo* by inducing caspase-mediated cell death. *Acta Pharmacol Sin* 2014:814-24.